

75001.842

DEPARTMENT OF THE INTERIOR.
UNITED STATES ENTOMOLOGICAL COMMISSION.

BULLETIN No. 6.

GENERAL INDEX

AND

SUPPLEMENT

TO THE

NINE REPORTS

ON THE

INSECTS OF MISSOURI.

BY

CHARLES V. RILEY, M. A., Ph. D.



WASHINGTON:
GOVERNMENT PRINTING OFFICE.

MARCH 24, 1881.

DEPARTMENT OF THE INTERIOR.
UNITED STATES ENTOMOLOGICAL COMMISSION.

BULLETIN No. 6.

GENERAL INDEX

AND

SUPPLEMENT

TO THE

NINE REPORTS

ON THE

INSECTS OF MISSOURI.

BY

CHARLES V. RILEY, M. A., Ph. D.

WASHINGTON:
GOVERNMENT PRINTING OFFICE.

MARCH 24, 1881.

C. V. RILEY, *Chief.*

A. S. PACKARD, JR., *Secretary.*

CYRUS THOMAS, *Disbursing Agent.*

ORDER OF MATTER.

1. INTRODUCTION.
2. TABLE OF CONTENTS.
3. CORRECTIONS.
4. NOTES AND ADDITIONS.
5. DESCRIPTIONS OF NEW SPECIES AND VARIETIES.
6. LIST OF DESCRIPTIONS OF ADOLESCENT STATES.
7. LIST OF DESCRIPTIONS, MOSTLY AMPLIFIED, OF SPECIES NOT NEW.
8. LIST OF ILLUSTRATIONS BY REPORTS.
9. CLASSIFIED LIST OF ILLUSTRATIONS.
10. GENERAL INDEX.
11. INDEX TO FOOD-PLANTS.

STORY OF THE

History of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

the story of the

INTRODUCTION.

The present Bulletin has been prepared in response to frequent suggestions from those having occasion to use the nine Annual Reports on the Insects of Missouri, made by me, as State Entomologist, to the State Board of Agriculture, during the years 1868 to 1877, inclusive. These Reports contain a good deal of matter anent the Cotton Worm, the Chinch Bug, the Rocky Mountain Locust, and other insects which the Commission has studied, and were published, as required by law, in the Annual Reports of said State Board of Agriculture for the years mentioned. That method of publication was always regretted by myself and by many others, inasmuch as the reports of the Board were generally volumes of such bulk as to delay publication and render mailing expensive. By virtue of the fact that they were distributed only to members of the State legislature and to State societies, access to them by persons outside the State of Missouri was extremely difficult; while the State printing and press-work were, as a rule, of a very unsatisfactory character. To avoid some of these difficulties it was my habit to have about 300 separate copies of the entomological portion printed on better paper, at my own expense, for distribution to correspondents both at home and abroad, and it is through these, principally, that the Reports have been accessible outside the State.

The demand for the Reports and the manner in which they have been used and commended by subsequent writers can but be gratifying to the author, who feels that whatever of commendation they deserve is due to the fact that they embody results of original investigation. They contain some matter that, with present light, he would expunge, and the earlier volumes, more particularly, contain imperfections which no one appreciates more fully than himself. Many of these are attributable to isolation from other working entomologists at the time, as well as to the almost absolute dearth of entomological works of reference in any of the libraries of Saint Louis.

The general plan of the Reports, which were addressed to the intelli-

gent cultivator of the soil rather than to the naturalist, is set forth in the following passage from the introduction to the first:

Fully aware that I write for those who, as a rule, are unversed in entomology, I have endeavored to treat of each insect with as little of the nomenclature of science as is consistent with clearness of expression. Yet, as much that is of scientific interest, such as descriptions of new species, must necessarily be inserted, I have had such descriptions printed in a type of smaller size than the text, so that it can be skipped if desirable, at the time of reading, and easily referred to for comparison, with specimens which one is desirous of naming. I have also endeavored to illustrate, as far as possible, the insects of which this report treats, believing that good illustration forms the basis of successful teaching in a science with which the general husbandman is not expected to be acquainted; for the eye conveys to the mind, in an instant, what the ear would fail to do in an hour. The practical man cares little to what genus or family an insect belongs, so long as he can tell whether it be friend or foe. He must become familiarized with the insects about him without having necessarily to overcome scientific detail and technicality.

I have made no effort at a systematic arrangement of the insects treated of. Indeed, that were useless for the purpose in view; but, in order that the reader may refer the more readily to any particular insect which interests him, I have separated them into three series—NOXIOUS, BENEFICIAL, and INNOXIOUS—and attached a very full index. For the benefit of those who are making a study of entomology, I have also given, with each species, the Order and Family to which it belongs, in parenthesis under each heading.

So far as possible, I have used a common name for each insect, knowing that the scientific name is remembered with greater difficulty, and is, consequently, distasteful to many. But as popular names are very loosely applied, and the same name often refers to different insects, in different localities, a great deal of confusion would ensue without the scientific name, which is, therefore, invariably added, for the most part, in parentheses, so that it may be skipped without interfering in any way with the sense of the text.

In order to add value to this general index, I have brought together tables of contents of the nine volumes and given corrections and some notes and additions. I have also reproduced the descriptions of new species, and added a list of descriptions of adolescent states, of descriptions of species not new, of illustrations by reports, of illustrations by classification, and of food-plants.

The Reports were independently paginated, and the separate copies were often distributed before the Agricultural Report was off the press. The date of publication and distribution is given for each in the tables of contents. The nomenclature of the Reports is retained in this Bulletin, the synonymy being indicated in the notes and additions and with the reproduced descriptions. The name of the author of the species and not of the genus was always given as authority, and in the later Reports I endeavored to indicate whether or not the insect was described under the generic name which it bears, by adding the authority without a comma when the specific name is coupled with the generic name under which it was first published — *e. g.*, *Phycita nebulo* Walsh — but placed it in parentheses when a different generic name was used than that under which the insect was first described — *e. g.*, *Acrobasis nebulo* (Walsh) — except where the whole name was already in parentheses when a comma

was used for the same purpose — *e. g.* (*Acrobasis nebulo*, Walsh). The same plan is adopted throughout this Bulletin.

It had always been my intention to publish a tenth volume and to end the decade with a review of, and general index to, the whole series. Indeed, an appropriation for the tenth year's work was made and the tenth report would have been duly issued had I not been called at the time to my present work for the General Government. This Bulletin is, in a measure, the fulfillment of that intention, and is issued in the hope that it will render the Reports more serviceable to the student of insect life and to those having to deal with insects injurious to agriculture.

My thanks are due to Messrs. E. A. Schwarz and W. H. Patton, agents of the Commission, for aid in its preparation.

C. V. R.

WASHINGTON, D. C., *March 1, 1881.*

TABLES OF CONTENTS.

Neither of the first five volumes contained a table of contents, the plan of giving such having been adopted with the sixth. Most of these tables are, therefore, prepared for this Bulletin, while those of the Sixth and Seventh Reports are amplified. Those of the last two volumes are reproduced as they were originally made.

REPORT I.

[Submitted December 2, 1868; published March, 1869.]

	Page.
Introductory	3
NOXIOUS INSECTS.	
The Bark-lice of the Apple-tree	7
Two species known to occur in the United States, 7 — Harris's Bark-louse not numerous enough to do material damage, 7.	
THE OYSTER-SHELL BARK-LOUSE	
Difference in the scales of the two species, 7 — Introduction of the Oyster-shell Bark-louse from Europe and its spread in the United States, 8 — Precautionary measures to prevent its introduction into Missouri, 8 — Its habits studied by different observers in 1867, 9 — Seasonal notes on the development of the insect, 10 — Only one annual brood in Missouri, 12 — Formation and nature of the scale, 12 — Rare occurrence of males, 14 — Difference of opinion among observers as to the mode of growth of the scale, 14 — Difficulty of explaining the spread of the insect from one tree to another, 15 — Its occurrence upon other kinds of trees, 15 — Natural enemies, 16 — Artificial remedies, 16 — Examination of young trees before planting, 16 — Washing with alkalies, etc., 17 — Scrubbing the branches with a stiff brush, 17 — Fumigating, 17 — Application of sheep-manure, 17 — Washes in general ineffective, 17 — The insect can most successfully be fought during three or four days of the year only, 18.	
The Periodical Cicada	18
Its singular life history, 18 — Seventeen and thirteen year races, 19 — The two races not distinct species, 19 — Two distinct forms occurring in both broods, 20 — Season of their appearance and disappearance, 22 — Natural history and transformations, 22 — Elevated chambers of the pupa, 22 — The larvæ frequently occurring at great depth in the ground, 24 — The operation of emerging from the pupa, 24 — Only the males are capable of singing, 24 — Trees in which the females deposit their eggs, 24 — Mode of oviposition, 24 — The newly hatched larva, 25 — The W on the wings of the Cicada, 25 — Enemies, 26 — Fungus infesting the imago, 26 — The sting of the Cicada, 26 — Wide-spread fear of the insect on ac-	

The Periodical Cicada—Continued.

count of its supposed stinging-powers, 26—Explanations of the sting, 27—Injury caused by the insect, 29—by the larva, 29—by the imago, 29—Fruitless attempts to stop the injury, 30—Chronological table of all well-ascertained broods in the United States, 30—The insect will appear during the next 17 years somewhere in the United States every year except in 1873, 41—Number of broods that will appear in the next 17 years in the different States, 42.

Apple-tree Borers..... 42**THE ROUND-HEADED APPLE-TREE BORER..... 42**

It is more numerous in trees on high land than on low ground, 42—Extent of its injury, 43—Its larva, 43—Appearance of the imago, 43—The hole made by the young larva, 44—It remains nearly three years in the larva state, 44—Its pupa state, 44—Remedies, 45—Alkaline washes, 45—Killing the larva by hot water, 45—Cutting out the larva, 46.

THE FLAT-HEADED APPLE-TREE BORER..... 46

Differences between it and the foregoing species, 46—Habits of the beetle, 47—Amount of injury caused by it, 47—Parasite attacking it, 47—Remedies, 47.

The Peach Borer..... 47

Its nature, 47—Differences in the sexes, 48—Remedies, 48—The mounding system the best remedy, 48—Testimony as to the value of the mounding system, 48—Other remedies, 49.

The Plum Curculio..... 50

Difference of opinion among authors on some points in its natural history, 50—Reasons for this difference of opinion, 51—Facts in its natural history, 52—It causes the spread of the peach-rot, 52—Fruit trees attacked and those not attacked by it, 53—It may hibernate as larva or pupa, but does generally as imago, 53—Mode of egg-laying, 54—It has one annual brood, 55—Walsh's experiments to show that it is two-brooded, 55—Natural remedies, 56—No parasites known to infest it, 56—Enemies, 57; The Pennsylvania Soldier-beetle, and its larva, 57; Lacewing-larva, 57; The Subangular Ground-beetle 58; Ground-beetle larva, probably of the Pennsylvania Ground-beetle, 59—Hogs as Curculio destroyers, 59—Artificial remedies, 60—Jarring the trees the most effectual method, 60—Dr. Hull's Curculio catcher 60—Lessons for the fruit-grower from the account of the Curculio, 62.

The Codling Moth or Apple Worm..... 62

It is common wherever apples are grown, 62—Description of the insect in its different states, 63—Its life-history, 63—Other fruits attacked by it, 64—Remedies, 65—Picking up the fallen fruit, 65—Entrapping the worms the best remedy, 66—Trimble's hay band system and how to apply it, 66—Attracting the moth by fires, 67.

Cut-worms..... 67

The natural history of twelve distinct species, 67—Definition of the term "Cut-worm", 67—Habits of Cut-worms, 67—Their natural history briefly given, 68—Difficulty of breeding them in captivity, 69—Climbing Cut-worms, 69—Injury done by them to orchards, 69—Fruit trees and shrubs they attack, 70—They attack large trees, 71.

THE VARIEGATED CUT-WORM..... 72

The full-grown larva, 72—The eggs, 72—Habits of the larva, 72—Cut-worm moths deposit their eggs on the leaves and not on the ground, 73—The imago, 73—Description of the insect as larva, pupa, and imago, 74.

	Page.
Cut-worms—Continued.	
THE DARK-SIDED CUT-WORM.....	74
General characters of the larva, 74 — Habits of, and injury done by it, 75 — Description of the imago, 75; of the larva and chrysalis, 76.	
THE CLIMBING CUT-WORM	76
Injury done by the larva, 77 — General characters of the larva, 77; of the moth, 78 — Description of the larva, 78; of the imago, 78.	
THE W-MARKED CUT-WORM.....	79
General characters of the larva, 79 — Plants it attacks, 79 — Characters of the moth, 79 — Description of the larva and chrysalis, 79.	
THE GREASY CUT-WORM	80
The larva very variable in coloration, 80 — Its injury to tomato and tobacco plants, 80 — General characters of the moth, 80 — Description of larva, chrysalis, and imago, 81.	
THE WESTERN STRIPED CUT-WORM	81
Resemblance of its larva to that of the Corn Rustic, 81 — General characters of the worm and moth, 82 — Description of the larva, 82.	
THE DINGY CUT-WORM	82
Difference between it and the foregoing species, 82 — General characters of pupa and imago, 82 — At least three species of our Cut Worms are difficult to distinguish, 83 — Description of larva, chrysalis, and imago, 83.	
THE GLASSY CUT-WORM	83
Habits and general characteristics of the larva, 83 — Characteristics of the moth, 84 — Description of larva and chrysalis, 84.	
THE SPECKLED CUT-WORM	84
Characteristics and habits of the insect, 84 — Description of larva, chrysalis, and imago, 85.	
THE SMALL WHITE BRISTLY CUT-WORM.....	86
Habits of the worm, 86 — Characteristics of the moth, 86 — Description of larva, chrysalis, and imago, 86.	
OTHER CUT-WORMS	87
Fitch's account of the Corn Cut-worm and the Yellow-headed Cut-worm, 87.	
THE WHEAT CUT-WORM	87
Injury caused by it, 87 — Description of the larva, 88.	
REMEDIES AGAINST CUT-WORMS	89
Natural enemies, 89; <i>Microgaster militaris</i> , 89; <i>Paniscus geminatus</i> , 89; The Spined Soldier-bug, 89; The Cut-worm Lion, 89 — Other enemies, 90 — Artificial remedies for climbing cut-worms, 90; for common field cut-worms, 91.	
Insects infesting the Potato	91
General remarks, 91 — Number of species affecting the Potato, 92.	
THE STALK-BORER	92
Habits of the larva and imago, 92 — Remedy, 93.	
THE POTATO STALK-WEEVIL	93
Its geographical distribution, 93 — Its habits, 93 — Remedy, 95.	
THE POTATO- OR TOMATO-WORM	95
It cannot sting with its horn, 95 — Its chrysalis, 95 — How the imago differs from the Tobacco-worm Moth, 95 — Remedies and parasites, 96.	
BLISTER-BEETLES	97
The Striped Blister-beetle, 96 — The Ash-gray Blister-beetle, 97 — The Black-rat Blister-beetle, 98 — The Black Blister-beetle, 98 — The Margined Blister-beetle, 98 — Synonymical remarks, 98 — Remedies for Blister-beetles, 99.	

	Page.
Insects infesting the Potato—Continued.	
THE THREE-LINED LEAF-BEETLE	99
Merdigerous habit of the larva, 99—It has two annual broods, 100— Other notes on the habits of the insect, 100.	
THE CUCUMBER FLEA-BEETLE	101
THE COLORADO POTATO-BEETLE	101
Its past history and future progress, 101—Its native home, 101—Its gradual spread eastward, 102—Its confusion with the Bogus Colorado Potato-beetle, 103—How the two species differ in habits, 104; in their larval states, 104; in the egg-state, 105—Description of the larva of <i>Doryphora juncta</i> , 106—Differences in the imagos of the two species, 106 —Habits of the Colorado Potato-beetle, 107—When it appears and dis- appears, 107—Number of eggs laid by each female, 107—Food-plants, 107—Singular fact that <i>D. juncta</i> has not acquired the habit of attack- ing the Potato, 108—Natural remedies, 109—Complicated economy of nature, 109—Decrease in the number of Potato-beetles on account of increase in the number of parasites, 109—The Colorado Potato-beetle parasite, 111—Its general character and habits, 111—Description of <i>Lydella doryphoræ</i> , 111—Ladybirds and their larvæ, 112—The Spined Soldier-bug, 113—The Common Squash-bug erroneously considered an enemy of the Potato-beetle, 113—The Bordered Soldier-bug, 114—The Many Banded Robber, 114—The Rapacious Soldier-bug, 114—The Virginia Tiger-beetle, 115—The Fiery Ground-beetle, 115—Blister- beetles, 115—The larvæ not touched by fowl, 115—Artificial remedies, 116—Ineffectiveness of mixtures tried, 116—Killing the beetle early in spring, 116—Pincers for crushing the insect, 116—Benson's machine, 116—Proper choice of varieties of potatoes, 117—The pest will over- run the Eastern States, 117—Carelessness in transmitting specimens of the beetle, 117.	
The Apple-root Plant-louse	118
Three distinct kinds of rots affecting the roots of Apple-trees, 118—The Root-louse the cause of one of these rots, 118—The cause of the other rots still hidden, 119—The Root-louse especially injurious in southerly latitudes, 119—It occurs also on other parts of the tree besides the root, 120—Description of the winged louse, 120—Fitch's description of the winged form refers to another species, 120—The Root-louse belongs to the genus <i>Eriosoma</i> , 121—Natural enemies, 121; Chalcis-fly, 121; The Root-louse Syphus-fly, 121; <i>Scymnus cervicalis</i> , 122—Artificial remedies, 123.	
The Woolly Elm-tree Louse	123
Its general appearance and habits, 123—Description of the winged form, 124,	
Insects Injurious to the Grape-vine	124
THE NEW GRAPE-ROOT BORER	124
Reports on the damage caused by it, 124—Description of the larva, 126— It belongs probably to the Cylindrical Orthosoma, 126—Former ac- counts of the natural history of this beetle, 127—Its injury known for several years, 127—Remedies, 128.	
THE GRAPE CURCULIO	128
Nature of the damage done by it, 128—Its larva, 128—The perfect beetle, 129—No injury done by it in 1868, 129.	
THE GRAPE-SEED CURCULIO	129
General appearance of the maggot, 129—Mr. Saunders' account of the damage done by it, 130.	

Insects Injurious to the Grape-vine—Continued.

THE GRAPE-CANE GALL-CURCULIO	131
The Gall caused by it, 131 — The larva, 131 — Its transformation, 131 — Description of the beetle, 132 — Differences between it and a closely allied species, 132 — The Gall caused by the punctures of the female beetle, 132 — Remedy, 132.	
THE GRAPE-VINE FIDIA	132
It is very injurious in Missouri, 132 — Habits of the beetle, 132 — Remedies, 133.	
THE GRAPE FRUIT-WORM	133
Amount and extent of the injury caused by it, 133 — Characteristics of the larva, 134 — Transformations, 134 — Description of larva, chrysalis and imago, 135 — Remedies, 135.	
THE EIGHT-SPOTTED FORESTER	136
Characteristics of the larva, 136 — It is not numerous enough to cause serious injury, 136 — Other caterpillars resembling it, 136.	
THE GRAPE-VINE PLUME	137
Work of the larva, 137 — Its habits and characteristics, 137 — The moth, 137 — Remedy, 138.	
THE SNOWY TREE-CRICKET	138
Characteristics of the insects, 138 — It is injurious, 138 — Nature of the injury caused by it, 138 — Remedy, 139.	
The Raspberry Geometer	139
Habits of the larva, 139 — Parasite attacking it, 139 — Characteristics of the moth, 139 — Description of the larva, 139; of the imago, 140.	
The Gooseberry Fruit-worm	140
Accounts of the injury caused by it, 140 — Habits of the worm, 140 — The moth, 141 — Remedies, 141 — Description of larva, chrysalis, and imago, 141.	
The Strawberry Leaf-roller	142
Extent and nature of the damage caused by it, 142 — Habits of the insect, 142 — Accounts of its injury in Indiana and Illinois, 142 — Remedy, 143 — Description of the imago and larva, 143.	
The White-marked Tussock-moth	144
The egg-mass, 144 — The larva and larval changes, 144 — The full-grown larva, 145 — Habits of the larva, 145 — Mode of casting off the larval skin, 145 — The cocoon, 146 — The imago, 146 — Two annual broods, 146 — Food-plants, 146 — Remedies, 147.	
The Bag-worm, <i>alias</i> Basket-worm, <i>alias</i> Drop-worm	147
Its geographical distribution, 148 — Injury caused by it, 148 — The egg, 148 — The larva and its growth, 148 — Habits of the larva, 149 — The chrysalis, 149 — The sex distinguishable in the chrysalis state, 149 — The imago, 149 — Food-plants, 150 — Parasites, 150; <i>Cryptus inquisitor</i> , 150; <i>Hemiteles thyridopterygis</i> , n. sp., 150 — Remedies, 151.	
The Ailanthus-worm	151
Injury done to the Ailanthus tree, 151 — Habits of the larva, 151 — The chrysalis, 151 — The imago, 152 — Geographical distribution, 152 — Remedy, 152 — Description of larva and chrysalis, 152; of the imago, 153.	
The Walnut Tortrix	153
Habits of the larva, 153 — General appearance of the moth, 153 — Phytophagic form of the insect on Snowberry, 153 — Description of larva, chrysalis, and imago, 154; of the variety <i>symphoricarpi</i> , 154.	

	Page.
The Seed-corn Maggot	154
Accounts of damage caused by it, 154 — The maggot, 155 — Transformation 155 — Description of the imago, 155 — Remedy, 155 — Habits of <i>Anthomyia</i> larvæ, 156.	
The White Grub	156
Account of the damage caused by it, 156 — Injury done by the perfect insect, 157 — <i>Résumé</i> of its life-history, 157 — Remedies, 157 — Regularity in the appearance of the beetle, 158 — Accounts of the fungus infesting the White Grub, 158.	
The American Meromyza	159
Nature of the damage caused by it, 159 — Characteristics of larva, chrysalis and imago, 160 — European Diptera with similar habits, 160 — Remedies, 161.	
The Sheep Bot-fly or Head-maggot	161
The insect in its different states, 161 — Its larva, 162 — Pupa, 162 — Characteristics of the imago, 162 — Fatal results of the presence of the maggot in the head of the sheep, 163 — Rabbits attacked by gad-fly, 164 — Testimony regarding the viviparous habits of the Bot-fly, 164 — Remedies, 165.	
Insect enemies of the Honey-bee	166
THE BEE-MOTH OR WAX-WORM	166
General appearance of the moth, 166 — There are no moth-proof bee hives, 166 — Habits of the worm, 167 — How its presence in the hive may be recognized, 167 — Prevention and remedy, 167.	
THE BEE-KILLER	168
It is an Asilid fly, 168 — Mr. Thompson's account of the fly, 168 — How it captures and kills bees, 168 — No remedy known, 168.	

BENEFICIAL INSECTS.

The Rear-horse, <i>alias</i> Camel-cricket, <i>alias</i> Devil's Riding-horse	169
Its food, 169 — How it grasps its prey, 169 — Difference in the sexes, 170 — The larva, 170 — The egg-mass, 170 — The mode of egg-laying, 170 — Voracious disposition of the Mantis, 171 — Its beneficial influence, 171 — Tachina-parasite of the Mantis, 171.	

INNOXIOUS INSECTS.

The Solidago Gall-moth	173
Gall caused by <i>Trypeta solidaginis</i> , 173 — Gall produced by the Solidago Gall-moth, 173 — Its natural history, 173 — Provision of the larva for its protection within the gall, 174 — Previous account of the gall, 174 — Gall caused by <i>Cochylis hilarana</i> on <i>Artemisia campestris</i> , 175 — Description of the Solidago Gall-moth as larva, chrysalis, and imago, 175 — Parasites attacking it, 175; the Inflating Chalcis, 176; <i>Eurytoma bolteri</i> , n. sp., 176; <i>Hemiteles</i> (?) <i>cressonii</i> , n. sp., 177; <i>Microgaster gelechiæ</i> n. sp., 177; other parasites, 178 — <i>Oberea</i> larvæ intruding the gall, 178.	
The Knotweed Geometer	179
Its natural history, 179 — Description of larva and chrysalis, 179.	
The Thistle Plume	180
Work of its larva on thistle-heads, 180 — Description of the larva, chrysalis, and imago, 180.	

REPORT II.

[Submitted December 2, 1869; published March, 1870.]

Preface	3
---------------	---

NOXIOUS INSECTS.

Report of the Committee on Entomology of the State Horticultural Society	5
--	---

Noxious insects less injurious in Missouri in 1869 than usual, 5 — The Army Worm and the Grain Plant-louse considerably injurious in Missouri in 1869, 5 — The Chinch Bug and the Codling Moth less injurious, 6 — A species of *Thrips* destroying great numbers of the Curculio, 6 — Eggs of the Apple-tree Plant-louse destroyed by insect foes and birds, 6 — According to Dr. Hull the "scab" in apples is caused by the Apple-tree Plant-louse, 7 — The Pickle Worm doing great damage during 1869, 7 — Importance of preventing the introduction of injurious insects, 7 — Cultivation causes insects to multiply unduly, 8 — More attention paid in Europe to injurious insects than in this country, 8.

IMPORTED INSECTS AND NATIVE AMERICAN INSECTS

The imported Currant-worm much more injurious than the native, 8 — Other instances showing the greater destructiveness of imported insect enemies than of their native representatives, 9 — Almost all our worst insect pests and pernicious weeds have been introduced from Europe, 10 — Few American insects and plants have become naturalized in Europe, 11 — The American fauna and flora not so highly improved and developed as in Europe, 12 — Australian fauna still more "old-fashioned" than America, 12 — The parasites of injurious insects are not imported with the insects themselves, 13 — Government aid should be solicited to exterminate recently imported injurious insects, 13 — But little attention given so far by our Government to assist the study of economic Entomology, 14 — Danger of introducing the Oyster-shell Bark-louse into Missouri, 15 — Immunity of the Pacific States from many of our fruit insects, 15.

The Chinch Bug	15
----------------------	----

It is the most injurious of all insects infesting grain, 16 — Its past history, 17; it was known in South Carolina in Revolutionary times, 17; it was injurious in Missouri as early as 1854, 17; noticed in Illinois in 1840, 17; it was very injurious in Missouri in 1868, 17; but hardly noticed in 1869, 17 — Probable reason why it was not noticed in Missouri in former times, 18 — Why it is not injurious in Massachusetts and New York, 18 — Its natural history, 18 — The pupa state in the different insect Orders, 18 — Time required for different insects to complete the cycle of development, 19 — The Chinch Bug is two-brooded in Missouri, 19 — Its winter quarters, 20 — Its rapid multiplication, 20 — Dr. Shimer's account of its nuptial flights, 21 — It deposits the eggs underground on the roots of the plant, 21 — The egg, 22 — Dimorphous forms of the Chinch Bug, 22 — Its destructive powers, 22 — Account of its appearance in immense numbers, 23 — Heading off the marching Bugs by a barrier of pine boards, 23 — Heavy rains destructive to the Chinch Bug, 24 — Moisture injurious to the egg, 24 — The Chinch Bug is always worse in a dry season than in a wet one, 24 — Dr. Shimer's theory on epidemic disease affecting the Chinch Bug, 25 — Cannibal foes of the Chinch Bug, 25; several species of Ladybirds, 25; the Weeping Lacewing, 26; How the

The Chinch Bug—Continued.

Lacewing larva seizes its prey, 26; The Insidious Flower-bug, 27; the Common Quail, 28—Amount of damage done by the Chinch-bug, 28—Remedies, 28—Burning in winter the old corn-stalks and other dead stuff on and near the fields, 29—Mixing winter rye among spring wheat, 29—Intercepting the marching Bugs by fence-boards, 29—Sowing gas-lime, 30—Other remedies, 30—Bogus Chinch Bugs, 31—Several species of Heteroptera confounded with the true Chinch Bug, 31—The smell emitted by the Half-winged Bugs, 32—The Insidious Flower-bug, 32—The Ash-gray Leaf-bug and its injury to grape-vines, 33—The Flea-like Negro-bug, 33; injury caused by it to raspberry, strawberry, and garden flowers, 34—Two other species of Negro-bug, 35—Recapitulation of the natural history of the Chinch Bug, 36.

The Army Worm 37

Four distinct caterpillars designated as Army Worms in this country, 37.

THE TENT-CATERPILLAR OF THE FOREST 37

It cannot properly be called an Army Worm, 37.

THE COTTON WORM 37

Historical data on the injury caused by it, 38—The egg, 38—The worm and its habits, 39—Mr. Lyman's incorrect account of its development, 39—The moth and its habits, 40—Its hibernation, 40—Remedies, 41.

THE SOUTHERN GRASS-WORM 41

It resembles in habits the veritable Army Worm, 41.

THE TRUE ARMY WORM 41

Its past history, 41—Rev. Powers' account of its invasion in the New England States in 1770, 42—Accounts of later invasions previous to 1861, 43—Years of its appearance in Illinois, 43—The invasion of the year 1861, 44—Its appearance in Missouri in 1869, 44—Its sudden appearance and disappearance, 45—Reason for the apparently sudden appearance, 45—Army Worm years are wet with the preceding year dry, 46—Reason for the increase and decrease of the number of worms, 46—Its natural history, 47—Previous accounts of its natural history, 47—When the eggs are laid, 47—Where they are laid, 48—Misdirected instinct in insects and birds, 48—Exceptions to the normal habit of the Worm, 48—Color of the Worm, 49—The chrysalis and imago, 49—Parasites, 50; The Red-tailed Tachina-fly, 50; Its beneficial work, 50; It infests also other insects, 50; Walsh's description of the fly, 51; It has been re-described as *Exorista OstenSackenii*, 51; The Yellow-tailed Tachina-fly, 51; Description of the fly, 51; The Glassy Mesochorus, 52; The Diminished Pezomachus, 52; The Military Microgaster, 52; The Purged Ophion, 53; The Army-Worm Ichneumon-fly, 53—Habits of the Army Worm and suggestions for its destruction, 53—Burning grass meadows in winter or early spring, 54—Plowing late in the fall, 54—The marching of the Worms, 54—Plants they prefer, 54—They become beneficial by devouring the chaff in the fields, 55—Ditching, 55—Description of the insect as larva and imago, 56.

Insects infesting the Sweet-potato 56**TORTOISE-BEETLES 57**

The Clubbed Tortoise-beetle affects the Irish Potato, 56—Its general appearance, 57—Characteristics of Tortoise-beetles, 57—Merdigerous habits of Tortoise-beetles and others of the same family, 58—General appearance of the larvæ, 58—Their dung parasol, 59—Larval molts, 59—Egg of Tortoise-beetles, 60—The chrysalis, 60—Habits of and injury done by the beetles, 60—Remedies, 61.

	Page.
Insects infesting the Sweet-potato—Continued.	
THE TWO-STRIPED SWEET-POTATO BEETLE	16
It seems to be confined to that plant, 61 — The larva and the use of its fork, 61 — Its pupa and imago, 61.	
THE GOLDEN TORTOISE-BEETLE	62
Food-plants and characteristics of the larva, 62 — Brilliant color of the beetle, 62.	
THE PALE-THIGHED TORTOISE-BEETLE	62
It is hardly distinguished from the foregoing species, 62.	
THE MOTTLED TORTOISE-BEETLE	63
Characteristics of the beetle, 63 — of the larva, 63.	
THE BLACK-LEGGED TORTOISE-BEETLE	63
Characteristics of imago and larva, 63.	
The Pickle Worm	64
OTHER INSECTS INFESTING CUCURBITACEOUS VINES	64
The Squash Borer, 64 — It seems to be confined to the Eastern States, 64 —	
The Striped Cucumber-beetle, 64 — Injury done by the beetle, 64; by the larva, 65 — The larva and pupa, 65 — Number of annual generations, 65 — Remedies, 66 — Extent of the injury caused by it, 66 — The 12-Spotted Diabrotica, 66.	
THE PICKLE WORM	67
Characteristics and description of the worm, 67 — Its habits, 67 — Characteristics of the moth, 68 — Accounts of injury done by the worm in Missouri and Illinois, 69; in other portions of the country, 70 — It was not known before as injurious, 70 — Remedy, 70.	
Insects injurious to the Grape-vine	71
THE HOG CATERPILLAR OF THE VINE	71
The egg, 71 — Characteristics of the larva, 71 — Its habits when about to transform, 72 — The chrysalis and imago, 72 — It is one-brooded North and two-brooded further South, 72 — It is very injurious, 73 — The Microgaster parasite and its development, 73 — Habits of caterpillar infested with the parasite, 73.	
THE ACHEMON SPHINX	74
Appearance and habits of the larva, 74 — The chrysalis, 75 — The insect is single-brooded, 75 — The moth and its issuing from the pupa shell, 75 — No parasites known, 76.	
THE SATELLITE SPHINX	76
How to distinguish its larva from that of the foregoing species, 76 — Development of the larva, 76 — Variations in color of the larva, 77 — Its position when at rest, 77 — The moth, 78.	
THE ABBOT SPHINX	78
Its distribution, 78 — The larva varies much in color, 78 — The chrysalis and imago, 79.	
THE BLUE CATERPILLARS OF THE VINE	79
The Eight-spotted Forester, 80 — Larva previously mistaken for it, 80 — habits and characteristics of the larva, 80 — Harris's description of the larva, 81 — The moth, 81 — Mr. Andrews' account of its ravages, 81 — Remedies, 82.	
The Beautiful Wood Nymph, 83 — Characteristics of the moth, 83 — Close resemblance between the larva of this and the foregoing species, 83 — The differences pointed out, 83 — Development of the insect, 83.	
The Pearl Wood Nymph, 83 — It greatly resembles the Beautiful Wood Nymph, 83 — Its probable larva, 84 — Practical importance of distinguishing these closely allied species, 84.	

	Page.
Insects injurious to the Grape-vine—Continued.	
THE AMERICAN PROCRIS.....	85
Work of its larva, 85—Description of full-grown larva, 86—The moth, 86—It is not very destructive, 86—Two annual broods of the insects, 86—Parasite of the American Procris, 87.	
THE NEW GRAPE-ROOT BORER.....	87
Correction of opinion formerly expressed, 87.	
The Broad-necked Prionus, 87—Duration of the larva state, 87—Its transformation, 88—It bores also in Apple roots, 88—Great damage done by the borer, 88—No good remedy known, 88.	
The Tile-horned Prionus, 89—How it differs from the foregoing, 89—Its occurrence on prairie land, 90—Small dimorphous male form, 90—The larva subsists also upon the roots of herbaceous plants, 90—Practical considerations, 91.	
THE GRAPE-SEED MAGGOT	92
The Grape-seed Curculio larva of the first report is that of a Hymenopterous insect, 92—The perfect insect is closely allied to the Joint-worm Fly, 92—Mr. Saunders' account and description of the imago, 93.	
The Canker Worm	94
The eggs, 94—The larva and larval changes, 95—Importance for the orchardists to recognize the true Canker Worm, 95—The Imported Elm leaf-beetle mistaken for it, 95—Description of the larva, 96—Its food-plants, 96—The chrysalis. 96—Only one annual brood in the latitude of Saint Louis, 97—The moth and its varieties, 97—It is less injurious in Missouri than in the Eastern States, 97—Remedies, 98—Classification of remedies proposed, 98—The trough and bandage systems, 99—Muriate of lime as remedy, 100—Jarring the tree, 101—Late fall plowing, 101—Summer plowing, 102—Efficiency of hogs, 102—Enemies, 102; Birds, mite and parasites, 102; Ground-beetles, 103; The Fraternal Potter-wasp, 103.	
Cabbage Worms	104
THE SOUTHERN CABBAGE BUTTERFLY.....	104
Its geographical range, 104—Injury caused by it in Missouri, 104—Description of the larva, 105—The chrysalis and imago, 105—Habits and other food-plants, 105.	
The Potherb Butterfly, 105—It is a Northern species, 105—It will very likely never occur in Missouri, 106—Geographical range of insects principally influenced by temperature, 106—Isentomic lines, 106—Southern insects found near Saint Louis, 106.	
The Imported Cabbage Butterfly, 106—Amount of damage caused by it in Canada, 107—Its spread westward, 107—It will undoubtedly spread to St. Louis, 107—The insect in Europe, 107—History of its introduction, 107—The insect in its different stages, 108—Its food, 108—Remedies, 109—Parasites, 109.	
THE CABBAGE PLUSIA	110
Characteristics and habits of the larva, 110—Its transformations, 111—Remedies, 111—Description of larva, chrysalis and imago, 111—A similar worm occurring on thistles, 112.	
THE ZEBRA CATERPILLAR	112
Habits and characteristics of the larva, 112—The chrysalis and the moth, 113—Two annual broods, 113—Food-plants, 113.	
The Tarnished Plant-bug	113
Injury caused by it to various trees and plants, 114—It is a very variable species, 114—Its development, 114—No effective remedy known, 115—Preventive measures, 115.	

	Page.
The Philenor Swallow-tail	116
Its food-plant, 116 — Damage done by it, 116 — Characteristics and development of the larva, 116 — Description of the larva, 117 — The pupa, 117 — The imago, 117 — Prevention, 118.	
The Cottonwood Dagger	119
General appearance of the larva, 119 — Two annual broods, 119 — Chrysalis and moth, 119 — Larvæ of other species belonging to the genus <i>Acronycta</i> , 119 — Parasites, 120 — Description of larva and imago, 120 — Characters and habits of other species of the same genus, 121.	
The Missouri Bee-killer	121
The true scientific name of the Nebraska Bee-killer, 121 — Wing-veins of the genera <i>Asilus</i> , <i>Promachus</i> and <i>Erax</i> , 122 — Description of the Missouri Bee-killer, 122 — How to destroy the flies, 123 — Habits and life-history of <i>Asilus</i> -flies, 123 — Description of larva and pupa of <i>Erax Bastardi</i> (?), 124 — Synonymical notes on the imago, 124.	

INNOXIOUS INSECTS.

The Goat-weed Butterfly	125
Its geographical distribution and position in classification, 125 — Its food-plant, 125 — Habits of the larva, 126 — Larval changes, 126 — Conformity in the color of the larva with that of the leaves, 127 — Description of the full-grown larva, 127 — Transformation of the larva to chrysalis, 127 — The two sexes of the imago, 127 — Hibernation, 128.	
The Black Breeze-fly	128
Breeze-flies beneficial in the larva state, 128 — Tormenting power of Breeze-flies, 128 — Their mode of flight, 129 — Our knowledge of their larval character and habits, 129 — General characters of the larva of the Black Breeze-fly, 129 — It is semi-aquatic, 129 — Walsh's description of the larva, 130 — Habits and food of the larva, 130 — Its transformations, 131 — Description of the pupa, 131 — Probable habits of Breeze-fly larvæ on the Western prairies, 132.	
Galls made by Moths	132
THE FALSE INDIGO GALL-MOTH	132
The gall and its structure, 132 — General appearance of the larva and the moth, 133 — Description of larva and imago, 133.	
THE MIS-NAMED GALL-MOTH.....	134
Is it a true gall-maker or an inquiline?, 134 — Walsh's description of the larva, 134 — Description of the imago, 134 — Generic characters, 134 — Reasons why the insect is an intruder and not a gall-maker, 134 — Enumeration of the known gall-making moths, 135 — How the gall is formed, 135.	

REPORT III.

[Submitted December 2, 1870; published April, 1871.]

Preface	3
----------------------	---

NOXIOUS INSECTS.

Snout-beetles	5
The whole vegetable kingdom and every part of each plant serve as food for insects, 5 — Enumeration of insects affecting the different parts of the Apple-tree, 5 — Other food-habits of insects, 7 — Vast extent of the science of entomology, 8 — Beauty and simplicity of classification in entomology, 8 — Each family distinguished by its general appearance, 8 — Unity of habits in each family, 9 — Distinguishing characters of Snout-beetles, 9 — Their larvæ, 10 — They are among the most injurious beetles, 10 — Injurious Snout-beetles in Europe, 11.	

	Page.
Snout-beetles—Continued.	
THE COMMON PLUM CURCULIO	11
It is single-brooded, 11 — Experiments to prove this fact, 12 — It hibernates as beetle, 13 — Form of the egg, 13 — Feeding habits of the beetle, 13 — Creaking noise produced by it, 14 — Stridulation in other insects, 14 — It is nocturnal rather than diurnal, 14 — Habits of the beetle at night, 14 — Remedies, 15 — The Ransom Chip-trap process, 15; explanation of the process, 15; it is not so successful as anticipated, 15; it is not a new discovery, 16; number of Curculios caught by it, 16; its success dependent on the character of the soil, 17; directions for using the process, 17; more experiments needed, 17 — Offering premiums for collecting specimens, 17 — Absurdity of the application of Paris Green for the Curculio, 18 — Jarring by machinery, 18 — The Hull Curculio-catcher defective in several respects, 18 — The Ward Curculio-catcher, 20; how it could be improved, 20; rules for using the machine, 21 — Curculio-catcher invented by Claxton & Stevens, 22 — Hooten's Curculio-catcher, 22; its advantages and mode of operation, 24 — Two true parasites of the Plum Curculio, 24 — The Sigalphus Curculio parasite, 24; Fitch's account of it, 25; Walsh's doubt about its being parasitic on the Curculio, 25; experiments and observations proving that it is a parasite of the Curculio, 25; its development and frequent occurrence around St. Louis, 25; it attacks also other soft-bodied larvæ, 26; points in its natural history, 26; its position in the system, 27; description of the imago, larva, pupa, cocoon and of the var. <i>rufus</i> , 27 — The Porizon Curculio parasite, 28; how it differs from the foregoing, 28; description of the imago, 28 — Importance of the work of these parasites of the Curculio, 29 — Artificial propagation and distribution of parasites, 29.	
THE APPLE CURCULIO	29
How it differs from the Plum Curculio, 30 — Its natural history, 30 — Food-plants, 30 — The punctures it drills into the fruit, 31 — The egg, 31 — The larva and its habits, 31 — The larva transforms within the fruit, 32 — Amount of damage it does, 33 — How its work in the fruit can be distinguished from that of other insects, 33 — It is very injurious in Southern Illinois and parts of Missouri, 33 — It is less injurious to apples than the Plum Curculio, 33 — Injury done by it to pears, 33 — The rot in apples principally produced by it and by the Plum Curculio, 34 — Season of its appearance, 34 — Remedies and preventive measures, 34 — Description of larva and pupa, 35.	
THE QUINCE CURCULIO	35
Its food-plants, 35 — It is very injurious to the quince in the East, 36 — Dr. Trimble's account of the damage done by it, 36 — How the beetle differs from the two preceding Curculios, 36 — Its transformations and habits, 37 — Its puncture on the fruit, 37 — It hibernates in the larva state, 37 — Its larva mistaken by Dr. Fitch for that of the Plum Curculio, 38 — The imago state lasts only two months, 38 — It does not attack the Apple, 38 — Remedies, 38 — Description of the larva and pupa, 39.	
THE PLUM-GOUGER	39
How it differs from the preceding species, 39 — Injury caused by it, 39 — Food-plants, 40 — It is often mistaken for the Plum Curculio, 40 — Season of its appearance, 40 — Holes bored by it in the fruit, 41 — The larva lives within the kernel of the fruit, 41 — Remedies, 41.	
THE STRAWBERRY CROWN-BORER	42
Distribution of the insect, 42 — Injury done by it, 42 — Habits and characters of the larva, 43 — Habits of the beetle, 43 — Remedies, 43 — Parasite, 44 — How the larva differs from that of the Grape-vine Colaspis, 44 — Description of the imago and larva, 44.	

Snout-beetles—Continued.

THE PEA-WEEVIL	44
Insect enemies of the garden pea, 44 — Characters of the <i>Bruchides</i> , 45 — Habits of other species of <i>Bruchidæ</i> , 45 — Frequent occurrence of its larva in green peas, 45 — Characteristics of the beetle, 46 — It is in all probability an indigenous North American insect, 46 — The beetle does not sting the peas, 46 — The eggs are fastened by the female beetle on the outside of the pod, 47 — The larva and its habits, 47 — Its transformations, 47 — Remedies and preventives, 48 — Examination of peas intended for seed, 48 — Concerted action necessary to exterminate the insect, 48 — Mr. Saunders' account of the occurrence of the Pea-weevil in Canada, 49 — Other preventive measures, 49 — Birds destroying the insect, 50.	
THE GRAIN BRUCHUS	50
Its introduction from Europe, 50 — How it differs from the Pea-weevil, 50 — Curtis' account of its habits in Europe, 51.	
THE AMERICAN BEAN-WEEVIL	52
Food-plants, 52 — Its geographical distribution, 52 — Accounts of damage done by it in New York and Pennsylvania, 52 — It has only lately become injurious, 53 — Habits of the larva and beetle, 54 — The proper nomenclature of the species, 54 — Description of the imago, 55 — Its differences from allied species, 54 — Note on descriptions based upon individual variations, 56.	
THE NEW YORK WEEVIL	57
Damage caused by it to fruit trees, 57 — Former descriptions of the insect, 57 — Its breeding habits, 57 — The Pear Blight not caused by the beetle, 58.	
THE IMBRICATED SNOUT-BEETLE	58
Injury done by it to vegetation, 58 — Its natural history still unknown, 58 — General appearance of the beetle, 58.	
THE CORN SPHENOPHORUS	59
Damage caused by it to corn plants, 59 — Characteristics of the beetle, 59 — Its larval history still unknown, 59 — Probable habits of the larva, 59 — Walsh's description of the imago, 59.	
THE COCKLEBUR SPHENOPHORUS	60
It is not injurious, 60 — Coloration of the beetle and its probable identity with <i>S. 13-punctatus</i> , 60 — The larva, 60 — Enumeration of other injurious Snout-beetles, 60.	
Insects injurious to the Grape-vine	61
THE GRAPE LEAF-FOLDER	61
Its geographical distribution, 61 — Generic characters, 61 — Characters of the moth, 61 — Sexual differences, 62 — Habits of the larva, 62 — Remedy and prevention, 62 — Natural enemies, 62 — Description of the larva, 62.	
THE GRAPE-VINE EPIMENIS	63
Its larva formerly mistaken for that of the Pearl Wood Nymph, 63 — Characters of the moth and sexual differences, 63 — Habits of the larva when about to transform, 64 — Its grape-vine feeding habits formerly unknown, 64 — Description of the larva, 64; of the chrysalis, 65.	
THE GRAPE-VINE PLUME	65
Work of its larva and of that of the Grape-vine Epimenis, 65 — Both larvæ rather beneficial when not too numerous, 65 — Characters of the larva, 66 — Peculiar form of the pupa, 66 — Protective mimicry of the pupa, 67 — Habits and appearance of the moth, 67 — Is it single or double-brooded, 67.	

	Page.
Insects injurious to the Grape-vine—Continued.	
THE COMMON YELLOW BEAR.....	68
How the young larvæ differ from the mature larva of the Grape-vine Plume, 68—Food-plants, 68—Color variations in the larvæ, 69—The chrysa- lis, 69—The imago, 69—Parasites, 69—Remedy, 69.	
THE SMEARED DAGGER.....	70
The larva is polyphagous, 70—Characters of larva, pupa, and imago, 70—Remedy, 70—Parasites, 71—Description of imago and larva, 71; of the pupa, 72.	
THE PYRAMIDAL GRAPE-VINE WORM.....	72
Distinguishing characters of the Worm, 72—Its food-plants, 72—Its trans- formations, 72—It is single or double-brooded according to latitude, 73—Its closely allied congener in Europe, 73—Remedies, 73—Descrip- tion of the larva, 73; of pupa and imago, 74—How it differs from <i>Amphipyra pyramidea</i> , 74—Description of larva and imago of the Spat- tered Copper Underwing, 75.	
THE GRAPE-ROOT BORER.....	75
Its distribution, 75—Distinguishing characters of the larva, 76—Its transformation, 76—Characters of the moth, 76—Description of the imago and its sexual differences, 76—Work of the larva on grape- roots, 77—Remedies, 77.	
THE SPOTTED PELIDNOTA.....	77
It is usually not injurious, 77—The larva and its habits, 78—The beetle, 78—Description of the larva, 78.	
THE GRAPE-VINE FLEA-BEETLE.....	79
It is well-known to the grape-grower in Missouri, 79—Its distribu- tion and food-plants, 80—Hibernation, 80—Damage done by the beetle in spring, 80—The eggs, 80—Damage caused by the larvæ, 80; their transformation, 80—Remedies, 80—Description of the larva and pupa, 81.	
THE GRAPE-VINE COLASPIS.....	81
Fitch's account of the insect, 82—Varieties of the beetle, 82—The larva in all probability attacks sometimes that of the Leaf-folder, 82—It lives on the roots of the Strawberry, 82—Larva of the European <i>Colaspis</i> <i>barbara</i> , 82—Difference in habits of larvæ of allied species, 83—Singu- lar characters of the larva of the Grape-vine Colaspis, 83—Work of the larva on Strawberry roots, 83—Remedy, 84—Description of the larva, 84.	
THE GRAPE-LEAF GALL-LOUSE.....	84
Its life-history not yet fully studied, 84—Previous accounts of the insect by Fitch, Shimer, and Walsh, 85—The root disease in France, 85—The <i>Phylloxera vastatrix</i> recognized as the cause of this disease, 85—Identity of the gall-louse with the root-inhabiting insect, 86—The American and European insects are identical, 86—Remedies tried in France, 86—The disease directly caused by the Phylloxera, 87—Injury done by the Phyl- loxera in Missouri, 87—Forming of the gall, 87—Propagation of the lice and multiplication of the galls, 88—The gall-lice descend in the latter part of the season to the roots, 88—Change of the insect after passing from the leaves to the root, 88—Questions still to be settled in the life-history of the Phylloxera, 88—Rare occurrence of the winged form, 89—The insect can be transported from one place to another on roots, 89—It hibernates on the roots, 89—Grape-vines that should be planted, 89—Number of indigenous species of the Grape-vine, 90— Grape-vines which are most seriously infested with the Grape leaf- louse, 90—Danger in planting the Clinton among other grapes, 91—In-	

Insects injurious to the Grape-vine—Continued.**THE GRAPE-LEAF GALL-LOUSE.**

sects acquiring different food-habits as illustrated in the Apple-maggot and the Pine-leaf Scale, 92—The different forms of the Grape leaf-louse, 93—Discussion on the proper place of the insect in the classification, 93—On Dr. Shimer's proposed new families *Dactylosphaeridæ* and *Lepidosaphidæ*, 93—Objections to Fitch's specific name *vitifoliæ*, 95—Identity of the European with the American insect, 95—The Apple-root louse is identical with the Woolly Aphis, 95—The Gall-inhabiting form of the Phylloxera identical with the root-inhabiting type, 96—Characters of the genus *Phylloxera* and its place in the system, 96.

The Colorado Potato-beetle again..... 97

Its onward march, 97—It invades the Dominion of Canada, 97—How it crossed Lake Michigan, 97—It will probably spread through Ontario unless preventive measures are taken, 98—Excellent chance to prevent its spread in Canada, 98—The Paris green remedy, 99—It is efficient if judiciously applied, 99—It does not affect the tuber, 99—Natural checks to the increase of the Potato-beetle 100—The Great Lebia destroying the larvæ, 100—Bogus experiments, 100—The true Remedy, 101—How to prevent the insect from becoming too numerous, 101—Planting early varieties of potatoes, 101.

The Codling Moth again..... 101

Hay-bands around the trunk of the tree more effectual than rags placed in the fork, 102—The Codling Moth is single-brooded in the more northern countries, but double-brooded in the latitude of St. Louis, 102—Sexual differences of the moth, 103—Sexual characters in the genera *Argynnis* and *Grapta*, 103—The Codling Moth also infests peaches, 103.

The Corn Worm alias Boll Worm..... 104

Its geographical range, 104—Injury done by it to corn, 104—It attacks tomatoes and other plants, 105—Food-plants of the Stalk-borer, 105—The egg of the Corn Worm, 105—Mr. Glover's account of the habits of the Boll Worm, 106—The larva is very variable in color, 107—Its transformations, 107—Number of annual broods, 107—Amount of damage done by it, 107—Remedies, 108—Attracting the moth by sweets, 108—Heard's moth-trap, 109.

The Fall Army Worm..... 109

Reports of its appearance in 1870, and how it was generally mistaken for the True Army Worm, 109—It was also mistaken for the Boll Worm, 111—Injury caused by it, 111—How it differs from the True Army Worm, 112—It is a very variable species in the imago state, 113—The Spiderwort Owlet-moth, and how it differs from the Fall Army Worm moth, 113—Number of annual broods and time of appearance of the Fall Army Worm, 114—The eggs and how they are deposited, 114—Preventive measures, 115—It is never injurious during two consecutive years, 115—Parasitic checks, 116—Description of the imago, 116; of the varieties and the earlier states, 117.

The Apple-tree Tent-caterpillar, or American Lackey-moth..... 11

The web-nests of the caterpillar and importance of their destruction, 118—The egg-mass, 118—The caterpillar and its habits, 119—Transformations of the insect, 119—The imago very variable in color, 119—Food-plants of the caterpillar, 120—Remedies, 120—Parasites and enemies, 120.

	Page.
The Tent-caterpillar of the Forest	121
The egg-mass and how the eggs are deposited by the female moth, 121 — Development of the larva, 122 — Fitch's descriptions of the full-grown larva, 123 — Confusion arising from want of uniform rule in describing larvæ, 123 — The cocoon, 124 — The chrysalis and the moth, 124 — The web spun by the caterpillar, 124 — Mr. Ferris's observations on differences in habits, appearance, and food-plants of the caterpillar, 125 — Phytophagic varieties or species, 127 — Food-plants of the caterpillar, 127 — Its destructive powers, 127 — Remedies, 128 — Natural enemies and parasites, 128 — Summary, 129.	
The Fall Web-worm	130
It is often mistaken for the Tent-caterpillars, 130 — It feeds upon almost every kind of trees and shrubs, 130 — The web spun by the worm 130 — General appearance of the worm, 130 — The chrysalis and imago, 131 — Number of annual broods, 131 — Plants it prefers, 131 — How it differs from the Tent-caterpillar, 132 — Remedies, 132 — Description of the larva, 132.	
The Blue-spangled Peach Worm	132
Winter retreat of the larva, 132 — General appearance of the larva, 133 — Chrysalis and imago, 133 — <i>Callimorpha vestalis</i> Packard synonymous with <i>C. fulvicosta</i> , 133 — Food plants, 134 — Description of the larva, 134.	
The Ash-gray Pinion	134
Food plants of the larva, 135 — Transformations of the insect, 135 — Characters of the moth, 135 — Description of larva and imago, 135 — Remarks on allied species, 136.	
BENEFICIAL INSECTS.	
The Glassy-winged Soldier-bug	137
It preys upon different species of leaf-hoppers, 137 — Its larva and pupa, 138 — How it seizes its prey, 138 — Coloration of the insect, 138 — It was never observed before to attack the leaf-hoppers of the Grape-vine, 139 — Habits of the <i>Phytocoridae</i> , 139.	
INNOXIOUS INSECTS.	
The White-lined Morning Sphinx	140
Resemblance of the Moth to a Humming bird, 140 — Interest attached to the study of Lepidopterous larvæ, 140 — Larval variations in the same species, 141 — Food plants of the larva, 141 — The light and dark colored forms of the larva, 142 — Its geographical range, 142 — Parasite, 142.	
Two of our Common Butterflies	142
THE ARCHIPPUS BUTTERFLY	143
Synonymy of its specific name, 143 — Its geographical distribution, 143 — Characters of the <i>Danaidæ</i> , 143 — Sexual differences, 143 — Food plants, 144 — Hibernation, 144 — Two annual broods, 144 — Description of the egg, 144 — The larva and its larval changes, 145 — How the horns of the larva become longer at each moult, 145 — The full grown larva, 146 — Interest attached to the metamorphoses of insects, 146 — How the larva becomes a chrysalis, 147 — The hardened chrysalis, 147 — Duration of the chrysalis state, 147 — The issuing of the butterfly, 148 — Protective colors of insects, 148 — Nauseous odor of the Archippus Butterfly in all its stages, 149 — The Tachina-parasite of the Archippus larva, 149 — Action of the parasitized larva, 149 — How the Tachina larva and other insects prepare the lid of their puparia, 149 — Characters of this Tachina-fly, 150 — Difficulties of the study of the <i>Tachinariæ</i> , 150 — Two forms occurring in <i>Tachina archippivora</i> , 150 — The Butterfly often congregating in immense swarms, 151 — Probable reasons of this assembling in swarms, 152.	

Page.

Two of our Common Butterflies—Continued.

THE DISIPPUS BUTTERFLY	153
------------------------------	-----

Distinguishing characters of the *Nymphalidæ*, 153 — Food plants and geographical range, 153 — The egg, 153 — Description of the egg, 154 — Development of the larva, 154 — Description of the mature and young larva, 154 — The insect hibernates as young larva, 155 — Case prepared by the larva for its winter quarters, 155 — Modifications of the case, 156 — Peculiar habit of the autumnal larvæ, 156 — Parasites, 157 — *Tachina*-fly, 157 — The *Disippus* egg-parasite, 157 — The *Disippus* *Microgaster*, 158.

MIMICRY AS ILLUSTRATED BY THESE TWO BUTTERFLIES, WITH SOME REMARKS ON THE THEORY OF NATURAL SELECTION	159
---	-----

Conformity of color between animals and their surroundings, 159 — Definition of the term "mimicry," 160 — Pungent odor possessed by the *Danaidæ*, 160 — Their mode of flight, 160 — Protection they derive from their peculiar odor, 161 — *Pieridæ* and *Danaidæ* in the Valley of the Amazon, 161 — Mimetic forms of *Pieridæ*, 161 — Explanation of the origin of mimetic forms, 162 — Mimicry between the *Archippus* and *Disippus* Butterflies, 163 — The *Ursula* Butterfly, 163 — Its caterpillar and chrysalis undistinguishable from those of the *Disippus*, 163 — Its imago has no protective color, 164 — Mr. Bennett's objections to the theory of Natural Selection producing mimetic forms, 165 — Mr. Scudder's objections, 166 — Discussion of Mr. Scudder's arguments, 167 — Mr. Murray's objections to the connection between mimetic resemblances and Natural Selection, 170 — Natural Selection not the only power producing mimicry, 171 — Reasons for discussing in this Report the theory of Natural Selection, 172 — Natural Selection involves belief in the doctrine of Evolution, 173 — Darwinism is neither irreligious nor atheistic, 174.

REPORT IV.

[Submitted December 2, 1871; published April, 1872.]

Preface	3
---------------	---

NOXIOUS INSECTS.

Notes of the Year	5
-------------------------	---

THE COLORADO POTATO-BEETLE	5
----------------------------------	---

Its injuries in 1871, 5 — Its appearance in great numbers in early spring, 5 — Exorbitant price of Paris green, 6 — Natural enemies of the beetle very abundant, 6 — Diminution in numbers of the beetle later in the season, 7 — Causes of such diminution, 7 — Damage caused by the Potato-beetle in Missouri, 7 — It invaded Canada in 1870, 8 — The Three-lined Potato-beetle mistaken for it in New York and Massachusetts, 8 — Its further spread eastward irresistible, 8 — Slow spread of the insect in the South, 9 — Its present extent northward, 9 — It spreads but does not leave the districts already invaded, 9 — It is not injurious to potatoes in Colorado at a certain altitude, 10 — New food-plants, 10 — It feeds upon cabbage, 11 — Its hibernation, 11 — Objections raised against the use of Paris green, 11 — Paris green is an efficient remedy and now in general use, 12 — Box for dusting Paris green, 12 — Mixing the poison with diluents, 12 — No serious cases of poisoning have come to knowledge, 13 — Antidote for Paris green, 13 — Other applications, 13 — Messrs. Saunders's and Reed's experiments with various substances, 14 — Experiments with decoctions of various plants, 15 — Air-slacked lime as a remedy, 15 — Mechanical means, 15 — Squire's Brushing machine, 15 — Creighton's

Notes of the Year—Continued.**THE COLORADO POTATO-BEETLE.**

Improved Patent Insect Destroyer, 15—Disadvantage of all mechanical means, 16—A simple and effective way of brushing off and killing the bugs, 16—Natural enemies increasing, 16—Chickens acquiring a taste for eggs and larvæ of the beetle, 16—Spiders are among its enemies, 17—The 15-Spotted Ladybird and its larva, 18—The Icy Ladybird, 18—The Ring-banded Soldier-bug, 19—The Dotted-legged Plant-bug, 19—The Spined Soldier-bug and its earlier states, 20—The Nebraska Bee-killer, 21—The Kansas Bombardier-beetle, 21—Rove-beetles of the genera *Philonthus* and *Quedius*, 21.

THE CODLING MOTH AGAIN 22

Time of year that the first moths appear, 22—Time required for development, 22—Proper time to apply the bandages around the tree, 22—It attacks peaches, 22—Best kind of bandages, 23—Wier's Apple-worm Trap, 23—Advantages and disadvantages of the trap, 24—Overestimating the value of Wier's trap, 25—Jarring, 25—Mr. Chapin's method of knocking down the wormy apples, 26—When this operation should begin, 26—Fires, lights and bottles of liquid not to be recommended as a remedy, 27—Worthlessness of Todd's book "The Apple Culturist," 28—Natural enemies, 28—The Pennsylvania Soldier-beetle and its larva, 28—The Two-lined Soldier-beetle and its larva, 29—Description of the larva of *Telephorus bilineatus*, 30—Summary, 30.

THE PERIODICAL CICADA 30

Reproduction, with corrections and additions, of the Chronological table of Broods I to VI from Report I, 31.

THE GRAPE-VINE COLASPIS AGAIN 34

Redescription of the larva from well-preserved specimens, 34.

The Harlequin Cabbage-bug 35

It was not known in Missouri prior to 1870, 35—Its geographical range and color variations, 35—Insect enemies of the cabbage plant, 35—Dr. Lynceum's account of its habits and injury caused by it in Texas, 36—Its appearance in Missouri in 1870, 36—The egg, 37—The larva and pupa, 37—Several annual broods, 37—The mature bug, 37—Injury caused by it, 37—Its congener in Europe, 38.

The Rascal Leaf-crumpler 38

Its proper scientific name, 38—It is hardly noticed in summer time, 38—Injury caused by it, 39—It hibernates as larva, 39—Habits of the larva, 39—The larval case, 39—Characteristics of the moth, 39—Food plants, 39—Remedies, 40—Natural enemies, 40—LeBaron's description of *Tachina phycita*, 40—*Limneria fugitiva*, 41—Description of imago, larva and chrysalis of the Rascal Leaf-crumpler, 41—Description of the variety *nebulella*, 42.

The Walnut Case-bearer 42

Other case-bearers enumerated, 42—The case of the Walnut case-bearer, 42—Differences between the moth and that of the Rascal Leaf-crumpler, 43—Natural enemies, 43—Description of *Perilitus indagator*, 43.

The Apple-leaf Skeletonizer 44

Its work on the leaves of Apple trees, 44—The worm and the chrysalis, 44—Appearance of the moth, 45—Mr. Hammond's account of the injury caused by the worm, 45—The European *Acrobasis consociella*, 45—Remedies and parasites, 45—Description of imago, larva and pupa, 46.

The Green Apple Leaf-tyer 46

It occurs almost always in company with the foregoing, 46—Characters and habits of the worm, 46—The Chrysalis, 47—Description of imago, larva and chrysalis, 47.

	Page.
The Lesser Apple Leaf-folder	47
Its larva and pupa closely resembles those of the foregoing species, 47 — Mr. Wier's account of its habits, 48 — Remedy, 49.	
The Apple-leaf Bucculatrix	49
It is not very injurious in Missouri, 49 — Account of damage caused by it in New York, 50 — The worm and its habits, 50 — Its transformations, 50 — Season of the appearance of the moth, 50 — Remedies, 50 — Hab- its of <i>Bucculatrix thuiella</i> , 51 — Description of larva and pupa of the Apple-leaf Bucculatrix, 51.	
The Apple-twigg Borer	51
Its frequent occurrence in Missouri, 51 — Characters of the beetle, 52 — The holes made by it in the twigs, 52 — The holes are made only for food and protection, 52 — The insect breeds probably in the sap-wood of forest trees, 52 — The larva of <i>Sinoxylon bassillare</i> mistaken for that of the Apple-twigg Borer, 52 — Remedy, 53.	
Insects injurious to the Grape-vine	53
THE RED-SHOULDERED SINOXYLON	53
Characteristics of the insect as imago, larva and pupa, 54 — Damage done by it to grape-vines, fruit trees and Hickory, 54 — Description of the larva and pupa, 54.	
Grape Disease	55
THE GRAPE-LEAF GALL-LOUSE	55
Its scientific name, 55 — The law of priority, 55 — European grape-vines valueless in the eastern U. S., 55 — Deterioration of some of our native vines, 56 — Climatic reasons for the failure of European vines, 56 — The principal cause of this failure is the Phylloxera, 56 — Further proof of the identity of the American with the European insect, 57 — Reasons for the identity of the Gall-louse with the Root-louse, 57 — Further facts respecting the habits of the Root-louse, 58 — Underground forms, 58 — The young lice and their habits, 59 — Hibernation, 59 — The pupa, 59 — The winged female, 59 — Susceptibility of different vines to the attacks of the louse, 60 — Classification of the North Amer- ican grape-vines according to their practical importance, 60 — Synop- sis of the True Grape-vines of the U. S., by Dr. George Engelmann, 60 — Difficulty of separating the cultivated varieties of grape-vine, 61 — Importance of a proper classification of cultivated grape-vines, 62 — Importation into Europe of resisting American vines, 62 — Enum- eration of the cultivated varieties and their susceptibility to the dis- ease, 63 — No variety is entirely exempt from the attacks of the root- louse, 64 — American vines which resisted the root-louse in Europe, 64 — Means of contagion from one vine to another, 64 — The spread of the root-lice from one vine to another, 64 — Transportation of the louse upon the roots of seedlings and cuttings, 64 — Spread of the disease in France, 64 — The winged female of the root-louse and her function, 65 — The vernal leaf-gall, 65 — Preference of the winged Phylloxera for the Clinton grape and its allies, 65 — Power of flight of <i>Phylloxera ca- ryæfolia</i> , 66 — Deposition of the eggs upon the leaf, 66 — Probable rea- sons why the injuries of the Phylloxera are greater in Europe than in America, 66 — Outward and more visible effects of the root-disease, 67 — Practical suggestions, 67 — No need to destroy the Clinton vines, 67 — Influence of soil on the intensity of the disease, 67 — Remedies, 68 — Destruction of the gall-lice, 68 — Destruction of the root-lice, 68 — Results of experiments in France with various substances, 68 — Irriga- tion and submersion, 69 — Résumé of the insect's history, 69 — No need of unnecessary alarm, 70.	

BENEFICIAL INSECTS.

	Page.
Silkworms	72
INTRODUCTORY	72
The <i>Morus multicaulis</i> fever and its reaction, 72—Increasing attention lately given to silk-culture in America, 72—North America well adapted to the raising of silk, 73—General outlines of the natural history of the eight species of Silkworms treated of in this Report, 74.	
THE MULBERRY SILKWORM	75
Its past history, 75—Earliest silk-culture in China and India, 75—Its introduction into Europe, 76—Value of silk produced in France, 76—The “Silk Supply Association” in England and its objects, 76—Countries exporting raw silk, 77—The name given to silk by different nations, 77—History of the Mulberry Silkworm in America, 77—Its introduction and failure of earlier efforts, 77—Renewal of silk-culture within the past decade, 77—Silk manufacture in the United States, 78—Favorable prospects for raising silk in this country, 78—Silk-growing in California, 79—Mr. Prevost’s “California Silk-grower’s Manual,” 79—False statements and exaggerations in Prevost’s book, 79—Sale of Californian Silkworm eggs, 80—Disastrous effect of the Franco-Prussian war on the egg trade, 81—Success of silk-culture in California dependent on the ability to reel the silk, 81—Silk-culture in Kansas, 82—Mons. Boissière’s silk establishment and its chances of success, 82—Silk-growing in Missouri, 83—The fall season in Missouri eminently propitious for rearing silkworms, 84—Natural history of the Silkworm, 84—Races of the Silkworm produced by domestication, 84—Effects produced on the insect by domestication, 85—The egg, 86—Larva and larval changes, 86—Cocoon, chrysalis and moth, 87—No insect parasite of the silkworm in Europe, 87—The “Uji” disease in China and Japan, 87—Diseases of the Silkworm, 88; The <i>Muscardine</i> , its effect and cause, 88; The <i>Pébrine</i> disease, its symptoms and cause, 89; Nature and origin of these diseases, 90—Other diseases of the Silkworm, 91—Best varieties or races, 90—Different forms of cocoons produced by different races, 92—How best to rear silkworms, 92—Rearing a very simple process, 92—Character of climate of the Japanese silk districts, 93—Keeping the eggs during the winter, 93—Hatching of the eggs, 93—Room and building for the rearing of the worms, 93—The feeding net or fillet, 94—Importance of carrying all the worms simultaneously through their moults, 94—Regularity of feeding, 95—Regulating the temperature, 95—Cocoonery, 96—Choking the chrysalis, 96—Egg-laying, 97—Selecting and fastening the cocoons for breeding purposes, 97—Treatment of the female moths after copulation, 97—Preservation of the eggs, 97—Reeling, 98—Great skill required to accomplish the work properly, 98—Classification of raw silk, 98—Preparing the cocoons for reeling, 98—Objects of and manipulations in reeling, 99—Best food for the worms, 100—Varieties of the Mulberry, 100—Cultivation of the Mulberry, 100—Osage Orange as silkworm food, 100—Introduction of the Osage Orange into France, 100—Experiments in America with feeding silkworms on Osage Orange, 101—Advantages and disadvantages of the Osage Orange, 102.	
THE CECROPIA SILKWORM	103
Changes made in its scientific generic name, 103—General appearance of the moth, 103—Fitch’s explanation of the specific name, 104—Food plants, 104—The cocoon, 104—Value of its cocoon as compared with that of the <i>Polyphemus</i> moth, 105—How the moth issues from the co-	

Silkworms—Continued.**THE CECROPIA SILKWORM.**

coon, 105 — The moth immediately after hatching, 106 — The egg, 106 — Description of the larval changes, 106 — Strange habit of birds of using the empty cocoon as a storehouse, 107 — The Cecropia worm cannot be classed as an injurious insect, 107 — *Samia columbia* ought to be considered a variety of *cecropia*, 107 — Parasites, 107 — The Long-tailed Ophion, 107; its mode of oviposition, 108; habits of its larva, 108 — The Cecropia Tachina-fly, 108; how it affects its victim, 108; its larva and imago, 109 — The Mary Chalcis-fly, 109; how it escapes from the cocoon of the moth, 109; description of the imago, 110 — The Cecropia Cryptus, 110; its habits, 110; description of the two sexes, 111; how it differs from the allied species, 111.

THE ALIANTHUS SILKWORM..... 112

Difference between the silkworms of the Castor-bean and Ailanthus, 112 — Introduction of the Ailanthus silkworm into Europe, 113 — Ailanthiculture in Europe, 113 — Its introduction into America, 114 — Disadvantages of the Ailanthus silkworm, 115 — Its acclimatization in America and Australia, 115 — Value of the cocoon, 115 — Mr. Forgemol's device for unwinding the cocoon, 116 — Natural history of the Ailanthus silkworm, 117 — The egg and mode of egg-laying, 117 — Larval changes, 117 — Pupation, 118 — The imago and its variations, 118 — Food plants, 118 — Directions for raising the worms, 119 — A good word for the Ailanthus tree, 120.

THE PROMETHEA SILKWORM..... 121

Value of the cocoon, 121 — Mode of egg-laying, 121 — Larval changes, 121 — How the cocoon is fastened to the twig, 122 — Striking sexual difference in the moth, 122 — *Callosamia angulifera* ought not to be considered as a different species, 122 — Food plants of the worm, 123 — Natural enemies, 123.

THE LUNA SILKWORM..... 123

The beautiful appearance of the moth, 123 — The cocoon, 124 — Food-plants, 124 — Larval changes, 124 — Season when the imago issues, 125 — No parasites of the worm known, 125 — *Actias selene* probably identical with *luna*, 125.

THE POLYPHEMUS SILKWORM..... 125

Mode of egg-laying, 125 — The egg, 126 — Food-plants, 126 — Larval changes, 126 — The cocoon, 126 — Mr. Trouvelot's account of the issuing of the moth, 127 — Wonderful vitality of the chrysalis, 127 — Characters of the moth and its variations, 128 — The broad antennæ of the male moth mistaken for a third pair of wings, 128 — Difficulty of reeling the cocoon, 129 — Number of annual broods, 129 — Parasites, 129.

THE YAMA-MAI SILKWORM..... 130

Its native home and food-plant, 130 — Its introduction into Europe and Australia, 130 — Attempts at raising it in America, 130 — The egg and how it should be kept over winter, 131 — Climate most favorable for the worm, 131 — The worm and its resemblance to the leaf, 132 — Food plants, 132 — Larval changes, 132 — The cocoon and the value of its silk, 133 — The moth and its habits, 133 — Difficulties of rearing the worm, 133 — Mr. Adams's account of rearing the Yama-mai Worm in Japan, 134 — The Uji parasite, 136.

THE PERNY SILKWORM..... 137

Its native home, 137 — How it differs from the preceding, 137 — Larval changes, 137 — The cocoon and its silk, 137 — The moth, 137 — Its culture in China, 138 — The Tusseh Silkworm, 138.

	Page.
Silkworms—Continued.	
SUMMARY.....	138
Successful silk culture possible in this country, 138 — Comparative value of the different species of silkworms, 138.	
INNOXIOUS INSECTS.	
The Horned Passalus	139
Its frequent occurrence in old logs, 139 — The noise produced by the beetle, 139 — It occurs only in decaying wood, 140 — The larva and its exceptional character, 140 — Description of the larva, 140 — Previous descriptions of <i>Passalus</i> -larvæ, 141 — Description of the pupa, 141.	
The Great Leopard Moth	141
Characters of the larva, 141 — Food-plants, 142 — Hibernation and transformations, 142 — The imago, 142 — The larva able to resist extreme cold, 143 — Description of the larva, 143.	
The Isabella Tiger Moth	14
Characters of the larva, 143 — Food-plants, 143 — Cocoon and chrysalis, 143 — The moth, 144 — The popular name "Fever-worm" in the South, 144 — No parasite known of this and the preceding species, 144.	
The Acorn Moth	144
The egg laid in acorns that have been ruined by <i>Balaninus rectus</i> , 144 — Habits of the larva, 144 — Characters of larva and moth, 145 — Description of imago, larva and pupa, 145.	

REPORT V.

[Submitted December 2, 1872; published April 18, 1873.]

Preface	3
ENTOMOLOGY, ITS RELATIONS TO AGRICULTURE AND ITS ADVANCEMENT...	5
Definition of Entomology, 5 — The term "insect," 5 — The four Subkingdoms of the Animal Kingdom, 6 — Characteristics of insect's, 7 — Classification of insects, 8 — Hymenoptera, 9 — Coleoptera, 10 — Lepidoptera, 11 — Hemiptera, 12 — Diptera, 13 — Orthoptera, 14 — Neuroptera, 14 — Osculant and aberrant groups, 15 — Importance of entomology as a study, 17 — Economic importance of entomology, 18 — Relation of insects to Agriculture, 18 — Damage inflicted by insects to Agriculture and Horticulture, 19 — Progress of economic entomology, 19 — Economic entomology in Europe, 19 — Insects in America much more injurious than in Europe, 21 — State entomologists and entomologists in public positions in the United States, 21 — How to counterwork noxious insects, 23 — Prevention, 24 — Checking the spread of injurious insects, 24 — Cure, 25 — Modification of Dr. Hull's Curculio-catcher, 25 — Encouragement of the natural enemies of injurious insects, 26 — Duties of a State Entomologist, 27 — Annual report, 27 — Correspondence and other duties, 28 — How to collect, preserve and study insects, 29 — Appliances for collecting, 29 — Use of the umbrella, 29 — The knapsack, 30 — The hand-net, 30 — The frame of the net, 30 — The bag of the net, 31 — Use of the sieve, 31 — Sugaring, 32 — Attracting moths by light, 32 — How to kill insects, 32 — The cyanide bottle, 32 — Use of chloroform, 33 — Entomotaxy, 34 — Insect pins, 34 — How to pin insects, 34 — How to mount small insects, 34 — Spreading board, 35 — Drying box, 36 — Insect boxes and cabinet, 37 — Mr. Lintner's boxes for Lepidoptera, 38 — Substances for lining insect boxes, 40 — Preserving insects in the collection, 41 — Relaxing specimens, 41 — Breeding insects, 41 — Breeding-cage, 42 — Note-book of the collector, 43 — How to transmit insects, 44 — Text-books, 44.	

NOXIOUS INSECTS.

	Page.
Notes of the Year	46
THE CODLING MOTH.....	46
Experiments with Wier's Apple-worm trap, 46 — Value of different materials for bandages, 47 — Jarring, 48 — Occurrence of the Apple-worm in California, 49 — Enemies and parasites, 44 — The Ring-legged Pimpla, 49 — The Delicate Long-sting, 50 — Other enemies, 51 — Efficiency of the Spined Soldier-bug, 51 — False doctrines for exterminating the Codling Moth, 51.	
THE COLORADO POTATO-BEETLE.....	52
Its comparative harmlessness in 1872, 52 — New food-plants, 52 — Its progress eastward up to 1872, 52 — Experiments with Paris green, 53 — New enemies, 53 — The Rust-red Social Wasp, 54 — The Rose-breasted Grosbeak, 54.	
THE APPLE-TWIG BORER.....	54
It attacks also pear twigs, 54 — It has been bred from grape-canes by Dr. Shimer, 54.	
EGG OF THE HORNED PASSALUS	55
The egg, 55 — The newly hatched larva, 55 — Rapid development of the insect, 55.	
EGG OF THE COMMON MAY BEETLE	55
Description of the eggs and how they are deposited, 55.	
EGG OF THE BROAD-NECKED PRIONUS.....	56
Characteristics of the eggs and where they are deposited, 56.	
EGGS OF AMERICAN TENT-CATERPILLAR	56
Correction of the figure given in Report III of the egg-belt, 56.	
COUNTERWORKING THE TOBACCO WORM	56
Mr. White's method of planting the Jamestown Weed among the potatoes and poisoning the blossoms thereof, 56.	
The Grape Phylloxera	57
Its popular name, 57 — Accounts of the unusual mortality among grape-vines in the spring of 1872, 57 — Causes of this mortality given in the accounts, 59 — Excessive drouth and overbearing, 60 — The Phylloxera is the true cause, 60 — Actual proof of the Phylloxera having caused the mortality among grape-vines, 61 — Influences that favored the increase of the lice in 1872, 61 — Importance of a full understanding and management of the Phylloxera, 62 — Its range in North America, 62 — Its spread in Europe, 63 — Inconstancy in the habits of the gall-lice, 63 — The leaves of Clinton vine no longer affected by the gall-lice since 1871, 63 — Method of formation of the leaf-gall, 64 — Relative immunity of American vines in Europe, 64 — Propagating American varieties from cuttings, 65 — Appreciation of American vines in Europe, 66 — Careless statement published as to the immunity of <i>Labrusca</i> -vine, 66 — Grafting the grape-vine, 67 — New theories, 67 — The Phylloxera is the true cause and not the effect of the disease, 67 — Mr. Laliman's theory that the insect has always existed in Europe, 68 — <i>Oidium Tuckeri</i> of Europe and America identical, 69 — Mr. Saunders's account of the presence of <i>Oidium Tuckeri</i> in America, 70 — Means of contagion of the disease from one vine to another, 69 — Flying capacity of the winged Phylloxera, 70 — The male louse, 71 — Remedies, 71 — Efficacy of carbolic acid and soot, 71 — Value of submersion or irrigation, 72 — Sprinkling with quick-lime, ashes, etc., 72 — Mr. Lichtenstein's experiments to allure the lice, 72 — Experiments with carbolic acid, 73.	

	Page.
The Oyster-shell Bark-louse of the Apple	73
Its occurrence in Missouri, 74 — Its appearance in Luray County, Missouri, 74 — Mr. Hanan's account of its spread, 75 — Its occurrence in Southern Missouri, 76 — Its occurrence in Mississippi and Georgia, 78 — Its appearance in Kansas, 79 — It is double-brooded in the South, 79 — The waxy secretion of Homoptera, 80 — The newly hatched louse, 80 — The larval scale, 81 — Development of the female scale, 81 — Growth of the male scale, 82 — The male louse, 83 — Rare occurrence of the winged male, 84 — Agamic multiplication, 85, — Mode of spreading, 85 — Food-plants, 86 — Varieties of the Apple tree preferred by it, 86 — Enemies and parasites, 87; Mites, 87; <i>Aphelinus mytilaspidis</i> and Dr. Le Baron's account of its habits, 88 — Easy transportation and introduction of the Aphelinus, 90 — Remedies, 90 — Application of oily substances, 90 — Bibliographical and Descriptive, 91 — The generic name, 91 — Signoret's classification of the <i>Coccidæ</i> , 92 — Specific name, 92 — Characteristics and habits of three allied species, 93 — A new name necessary for our apple-tree species, 94 — Description of the eggs and the winged male, 95; of the male and female scale and of the female louse, 96.	
The Pine-leaf Scale-insect	97
Nature of the malady caused by it, 97 — Natural history of the insect, 98 — The male, 99 — There are two annual broods, 99 — It is confined to the Pines proper, 100 — Natural enemies, 100; The Twice-stabbed Ladybird, 100; The Painted Ladybird and description of its larva, 101 — Remedies, 101 — Stripping the old leaves, 102 — Application of powdered and liquid substances, 102.	
The Hickory Bark-borer	103
Accounts of the damage caused by it in Missouri, 103 — Habits of the European <i>Scolytus destructor</i> , 104 — Various kinds of Hickory attacked by the Hickory Bark-borer, 105 — Its natural history, 105 — Natural enemies, 106 — Descriptions of the Three-banded Spathius, 106 — Mr. Cresson's description of <i>Bracon scolytivorus</i> , 106 — Remedies, 107 — Description of the imago, 107 — Is it different from <i>Scolytus 4-spinosus</i> ? 107.	
The Rose Chafer	108
Great damage caused by the beetle in 1872, 108 — Remedy, 109 — Harris's account of its natural history, 109.	
The False Chinch-bug	111
It was not known as injurious before 1872, 111 — Accounts of injury done by it, 111 — How it differs from the true Chinch-bug, 112 — Its probable natural history, 112 — Description of imago, larva, and pupa, 113 — Variation in the imago, 113 — Its great abundance in the Fall of 1872, 114.	
Insects injurious to the Grape-vine	114
THE GRAPE-VINE APPLE-GALL.....	114
The breast bone of Gall-gnat larvæ, 114 — The gall mistaken for an apple, 114 — Form of the gall, 115 — Habits of the larva, 116.	
THE GRAPE-VINE FILBERT-GALL.....	116
Appearance of the gall, 116 — Larva of the gall-maker, 117.	
THE GRAPE-VINE TOMATO-GALL.....	117
Its curious resemblance to a tomato, 117 — Various shapes assumed by it, 118 — The larva, 118 — Enemies of the larva, 118.	
THE GRAPE-LEAF TRUMPET-GALL.....	118
Characteristics and occurrence of the gall, 118.	

	Page.
Eggs in and on canes and twigs	119
Probable eggs of the Jumping Tree-cricket, 119 — The egg-punctures on grape-cane, 119 — Characters of the egg, 119 — General appearance of the cricket, 120.	
Eggs of the Snowy Tree-cricket, 120 — Trees and shrubs attacked by this cricket, 120 — The egg, 120 — Habits and natural history, 120 — Injury done by it, 121 — The eggs mistaken by Fitch for those of the Buffalo Tree-hopper, 121.	
Egg-punctures of the Buffalo Tree-hopper, 122 — Development of the larva, 121 — Characters and habits of the perfect insect, 122 — Egg-punctures of some unknown Tree-hopper, 122.	
Egg-punctures of the Frosted Lightning-hopper, 122 — Development and habits of the insect, 122.	
Egg-punctures probably of <i>Orchelimum glaberimum</i> , 123 — The egg, 123 — Characters and habits of the imago, 123.	
Eggs of the Oblong-winged Katydid, 123 — They have been mistaken by Harris for those of the Broad-winged Katydid, 123 — Ovipositors and modes of egg-laying in the three Katydids occurring in Missouri, 124 — Increase in thickness of Katydid eggs before hatching, 124.	
Stinging larvæ	125
The Potato-worm falsely considered as a stinging caterpillar, 125 — General harmlessness to man of insect larvæ, 125 — Stinging caterpillars in Europe and North America, 126 — List of such larvæ in U. S., 126 — Description of larva and chrysalis of <i>Acronycta xylinoides</i> , 126.	
THE BUCK MOTH OR MAIA MOTH	127
Flight and characteristics of the moth, 127 — The egg and mode of egg-laying, 128 — Appearance of the full-grown larva, 129 — Larval changes, 129 — Color variations of the larva, 130 — Habits of the young larva, 130 — The sting of the larva, 131 — The pupa, 131 — Issuing of the moth, 132 — Irregularity and retardation in development, 132; the species benefits thereby, 132 — Food plants, 132 — Natural enemies, 132.	
THE IO MOTH	133
The male and female moth, 133 — The egg, 134 — The larva and its urticating power, 134 — Larval changes, 135 — Pupation, 135 — Food plants, 136 — Parasites, 136.	
The Green-striped Maple-worm	137
Account of its occurrence in great numbers in Kansas, 137 — Injury done by it to Soft maples, 138 — The egg, 138 — Larval changes, 138 — The chrysalis, 139 — The imago, 139 — Natural enemies, 139 — Description of <i>Belvoisia bifasciata</i> , 140 — Remedies, 141.	

INNOXIOUS INSECTS.

The Hellgrammite Fly	143
The eggs, 143 — Respiratory apparatus of the larva, 143 — Habits of the larva when about to transform, 143 — The pupa, 144 — Sexual difference in the imago, 145.	
The Goat-weed Butterfly	145
The egg, 146 — Habits of the larva, 146 — Larval changes, 146 — Transformation, 147 — New food-plant, 147 — Simulation of the color of the food-plant by the larva, 147 — Two annual broods, 148 — Hibernation of the imago, 148 — Natural enemies, 149.	

	Page.
On a new genus in the Lepidopterous Family Tineidæ with remarks on the fertilization of <i>Yucca</i>	150
Generic characters of <i>Pronuba</i> , 150 — Description of <i>Pronuba yuccasella</i> , 151 — Plants requiring the aid of insects for fertilization, 152 — Fertilization of Orchids, 152 — Fructification of <i>Yucca</i> , 153 — Yuccas must rely on insects for fertilization, 153 — Insects frequenting Yuccas, 154 — Diurnal and nocturnal habits of <i>Pronuba</i> , 154 — How the female moth fertilizes the plant, 154 — Oviposition, 155 — The larva within the young fruit, 155 — Description of the larva, 155 — Only a small percentage of fruit not infested by the larva, 156 — The larva leaves the capsule and enters the ground for hibernation, 156 — Mutual adaptation of plant and insect, 156 — The moth doubtless occurs wherever Yuccas grow wild, 157 — Easy transportation of the cocoon, 158 — Further facts regarding the fructification of <i>Yucca filamentosa</i> and <i>gloriosa</i> , 158 — Yuccas seeding in Europe, 159 — Range of the insect, 159 — Mr. Stainton's opinion on the characters of the <i>Yucca</i> moth, 160.	

REPORT VI.

[Submitted December 2, 1873; published March, 1874.]

Table of Contents	3
Preface	6

NOXIOUS INSECTS.

Notes of the Year	9
THE CODLING MOTH	9
Failure of the apple crop in 1873, 9 — Dr. LeBaron's observations on the habits of the worm, 9 — Proportion of worms leaving the fruit before it falls, 10 — How it affects pears, 10 — Time and method of using bandages, 10 — Westward spread of the insect, 10.	
THE COLORADO POTATO-BEETLE	11
How it has affected the price of potatoes, 11 — New food-plants, 11 — Its progress eastward during 1873, 12 — Improved methods of using Paris green, 13 — Device for jarring off the bugs, 14 — European publications on the insect, 15 — Danger of its introduction into Europe, 15 — Precautionary measures to be taken in Europe, 16.	
THE COTTON WORM	17
Paris green suggested as remedy, 17 — Address before the National Agricultural Congress, 17 — Mr. Glover's summary on experience with Paris green, 19 — Experiments with the poison, 19 — Johnson's Sprinkling machine, 20 — Patents on Paris green, 20 — The Royall mixture, 21 — Hibernation of the insect, 22 — Natural enemies, 23 — Geographical range, 23 — Position of the moth when alighting, 24.	
THE CANKER WORM	24
Dr. LeBaron's summary of remedies, 24 — Mr. Milliken's experience with the rope and tin trap, 25 — The Paris green remedy, 26 — A new trough, 26 — Birds which destroy the worm, 27 — Mr. Mann's observations on the insect, 28 — Two species have hitherto been confounded, 28 — The English Sparrow and the increase of the White-marked Tussock-moth, 29.	
Insects injurious to the Grape-vine	30
THE GRAPE PHYLLOXERA	30
The term "Phylloxera", 30 — Bibliographical history, 30 — Characters of the genus and its position in the system, 33 — Biological history, 33 — Different forms which the insect assumes, 33 — The gall-inhabiting form, 34 — The root-inhabiting form, 38 — Balbiani's discovery of the	

Insects injurious to the Grape-vine—Continued.**THE GRAPE PHYLLOXERA.**

true sexual individuals and the winter egg of *Phylloxera quercus*, 41 — The winter egg not essential to the winter life of the Grape Phylloxera, 42 — Polymorphism in Phylloxera, 43 — Conclusive proof of the identity of the root- and gall-lice, 44 — Practical considerations, 44 — The more manifest and external effects of the disease, 44 — Mode of spreading, 45 — Swarming of winged specimens of *Phylloxera caryæfoliæ*, 45 — Power of flight in the Grape Phylloxera, 45 — Spread of the disease through the winged females, 46 — Susceptibility of different vines to the disease, 46 — Practical importance of knowing the more resisting and more susceptible vines, 46 — List of the cultivated species and varieties of vines indicating their relative susceptibility to Phylloxera, 47 — Prophylactic means of coping with the disease, 48 — Grafting the more susceptible onto the roots of resisting vines, 48 — Other preventive measures, 50 — Natural enemies, 50 — Insect enemies of the gall-louse, 50 — Enemies of the root-louse, 52 — The Phylloxera mite, 52 — Different forms assumed by mites in their development, 52 — *Hoplophora arctata*, 53 — Peculiarities of the *Oribatidæ*, 54 — Direct remedies, 55 — Submersion a perfect remedy, 55 — Application of pure insecticides without satisfactory results, 56 — Range of the insect in America, 57 — Injury caused by it in America, 58 — Reasons why the insect is more injurious in Europe, 59 — False theories, 60 — The Phylloxera is the cause and not the effect of the disease, 60 — It is a native American insect and has been imported into Europe, 62 — It is peculiar to the Grape-vine, 63 — The American Oak Phylloxera and its natural history, 64 — Conclusion, 65.

APPENDIX AND NOTES TO THE ARTICLE ON GRAPE PHYLLOXERA.....

66

Diagnosis of *Phylloxera vastatrix*, 66 — Influence of the insect's puncture, 67 — The supposed male of the gall-louse, 67 — The non-cultivation of the Clinton grape, 67 — Transient nature of the leaf-galls, 68 — Constancy of the differences between the forms assumed by the insect, 68 — Supposed sexual individuals, 68 — Number of generations annually produced, 69 — Number of molts, 69 — Transplanting root-lice on to the leaves, 69 — Nature of the swelling on the roots produced by Phylloxera, 70 — The true Grape-vines of the United States, by Dr. G. Engelmann, 70 — Exceptional instances where the European Vine has succeeded in America, 76 — Grafting the more susceptible onto the roots of the resisting varieties, 78 — Descriptions of *Tyroglyphus phylloxeræ* and *Hoplophora arctata*, 81 — Efficacy of inundating the vineyards, 82 — Facts showing that the disease of Grape-vines in America is principally caused by Phylloxera, 82 — Description of *Phylloxera rileyi*, 86 — The true sexual individuals and the winter egg, 86.

THE BLUE CATERPILLARS OF THE VINE.....

87

1. The Grape-vine Epimenis, 87 — Habits and characters of the larva, 87 — Description of larva and chrysalis, 88.
2. The Beautiful Wood Nymph, 88 — Characters and food plants of the larva, 88 — The moth and the egg, 89 — Description of the egg and larva, 89 — Of the chrysalis, 90.
3. The Pearl Wood Nymph, 90 — Its larva almost undistinguishable from that of the foregoing species, 90 — Food-plant, 91 — Distinguishing characters of the moth, 91 — Mr. Lintner's description of the larva, 92 — Mr. Lintner on the difference between the larva of *Eudryas grata* and *E. unio*, 93.
4. The Eight-spotted Forester, 94 — Description of the larva, 94; Chrysalis and imago, 95.

Insects injurious to the Grape-vine—Continued.**THE BLUE CATERpillars OF THE VINE—Continued.**

Summary, 95 — Comparison of the larvæ of these four species, 95 — Remedies, 96.

- The Red-legged Ham-beetle** 96
 Its popular name, 96 — It has been the cause of an interesting law-suit, 96; Entomological information necessary to a just verdict, 98 — Another case before a jury requiring botanical knowledge, 98 — Injury caused by the beetle in St. Louis, 99 — The eggs, and how they are deposited, 99 — The larva and its habits, 99 — The pupa, 99 — Prevention, 100 — Other species associated with it, 100 — Habits of other species of *Cleridæ*, 101 — How *Mecrobia ruficollis* saved the life of Latreille, 101 — Description of the larva of *Corynetes rufipes*, 101 — Description of the pupa, 102.
- The Clover-hay Worm** 102
 Its geographical distribution, 102 — It has probably been imported from Europe, 101 — Its past history and accounts of damage caused by it, 102 — Its natural history, 105 — Remedies, 105 — Description of larva and chrysalis, 103; of the imago, 107 — Its allied congener, *Asopia olinalis*, 107.
- The Legged Maple Borer** 107
 The burrows of the larva, 108 — Appearance of the moth, 108 — Remarks on the nomenclature of the species, 108 — Injury caused by it to maples, 109 — It attacks only trees already injured by some cause or another, 109 — Remedies, 110 — Description of imago, larva and pupa, 110.
- The Raspberry Root-borer** 111
 Work of *Oberea perspicillata* in canes of Raspberry and Blackberry, 111 — Injury caused by the Root-borer, 111 — General appearance and habits of the worm, 112 — Description of imago and larva, 113.
- The Northern Brentthian** 113
 Its occurrence and distribution, 113 — Characteristics of the beetle, 114 — Fighting habits of the males, 114 — How the male assists the female in ovipositing, 115 — How much time is required for the transformations of the insect, 115 — Habits of the larva, 115 — Description of the larva, 115; of the pupa, 116 — Real position of the Brentthians in the system, 116 — The specific and generic names, 116 — How another larva has been mistaken for that of the Northern Brentthian, 117 — Description of this larva, which evidently belongs to the Tenebrionidæ, 118.
- The Jumping Sumach-beetle** 118
 Sumach industry in Europe and America, 118 — Species of sumach possessing economic value, 119 — Appearance and habits of the beetle, 119 — The egg-masses, 120 — Development and habits of the larva, 120 — Transformation, 120 — The two annual broods, 120 — Injury caused by it to the sumachs, 121 — Geographical range, 121 — Remedies, 121 — Description of larva and pupa, 121 — Variations of the beetle, 122.

BENEFICIAL INSECTS.

- The Unadorned Tiphia or White Grub Parasite** 123
 Other enemies of the White Grub, 123 — Cocoon and larva of *Tiphia*, 123 — It is undoubtedly a parasite of the White Grub, 124 — Characters of the genus *Tiphia*, 124 — Habits of *Tiphia femorata*, 124 — Habits of the *Tiphia* larva, 124 — Characters of the Unadorned *Tiphia*, 125 — Beetle parasitic upon it, 125 — Description of larva and imago, 126 — The species has been described under three different names, 126.

INNOXIOUS INSECTS.

	Page.
The Dominican Case-bearer	127
Characters of the beetle, 127 — Mode of oviposition, 128 — Oviposition of <i>Chlamys plicata</i> , 128 — The young larva and its food habits, 129 — The case of the larva, 129 — Characters of the larva, 130 — Time required for its development, 130.	
The Yucca Moth	131
Its natural history completed, 131 — Description of the chrysalis, 131 — Hatching of the chrysalis, 132 — Method of oviposition, 133 — Oviposition is followed by pollination, 134 — The egg in the young fruit and the embryo larva, 134 — It is the only insect that can well fertilize the Yuccas, 135.	
Hackberry Butterflies	136
The European Purple Emperor, 136 — Insufficient former account of their earlier states, 136 — Species of <i>Celtis</i> in Missouri, 137.	
THE EYED EMPEROR	137
The full-grown larva, 137 — Habits of the larva, 138 — Pupation, 138 — The imago, 139 — The egg and the young larva, 139 — Two annual broods, 139 — Hibernation of the young larva, 140.	
THE TAWNY EMPEROR	140
How it differs from the Eyed Emperor, 140 — The egg, 141 — Larval changes and habits, 141 — Egg parasite, 142.	
BIBLIOGRAPHICAL	142
The generic name, 142 — The specific names of the two species, 143 — Fabricius's original description of <i>Lycaon</i> and <i>Herse</i> , 144 — Other species of the genus in the United States, 145 — On the validity of <i>Alicia</i> , 145 — Descriptions of the earlier states of both species, 146 — Their popular names, 148 — The scientific specific names, 149.	
Katydids	150
General considerations, 150 — Spring in Europe and America, 151 — Stridulating noise produced by crickets and grasshoppers, 152 — Sounds inaudible to man, 152 — Grasshoppers, Katydids, Locusts, 153 — Habits and general appearance of our Katydids, 154 — They oviposit above ground, 154, — Their ovipositors not rudimental, 155.	
THE ANGULAR-WINGED KATYDID	155
It is the most common species in Missouri, 155 — General character of the insect, 155 — The eggs, 155 — Erroneous statement concerning the egg, 156 — How the female deposits the eggs, 156 — Number of eggs laid by the female, 158 — Hatching of the larva, 158 — Food of the larva, 158 — Changing from the pupa to the perfect state, 159 — Its song, 159 — It is capable of domestication to a certain degree, 160 — Description of the immature states, 161 — Natural enemies, 162 — The Back-rolling Wonder, an egg-parasite, 162 — Curious habit of the female <i>Antigaster</i> , 162 — Description of <i>Antigaster mirabilis</i> , 163 — Striking sexual differences, 163.	
THE NARROW-WINGED KATYDID	164
Distinguishing characters, 164 — Method of oviposition, 165 — The egg, 165 — Its song, 166 — Description of larva and pupa, 166.	
THE BROAD-WINGED KATYDID	167
It is the true Katydid, 167 — Distinguishing characters, 167 — Mode of oviposition as observed in confinement, 167 — Mr. Jaeger's erroneous statement regarding the oviposition, 168 — Its song, 168.	
THE OBLONG-WINGED KATYDID	169
It has not yet been found in Missouri, 169 — Distinguishing characters, 169 — The eggs not yet known, 169 — Abnormal female specimen, 169.	

REPORT VII.

[Submitted at time of publication, April, 1875.]

Page.

Preface.....	III
Table of Contents.....	VII

NOXIOUS INSECTS.

The Colorado Potato-beetle	1
Its gradual spread eastward, 1—It reached the Atlantic during the year 1874, 1—Injuries done during the year 1874, 2—Alarm about it in Europe, 3—Prohibiting the importation of American potatoes by European governments, 3—The insect probably introduced into Europe in the perfect form, 3—It would doubtless thrive in Europe if imported, 4—Its ravages exaggerated and underrated, 5—On the safety and advisability of the use of Paris green, 8—Past experience with the poison, 10—Influence of Paris green on the plant and on the soil, 11—Influence of the green on man indirectly through the soil or through the plant, 13—The beetle eats as well as the larva, 14—It passes the winter as imago, 14—New food-plants, 14—New means of destruction, 15—The Gray Sprinkler, 15—The proper scientific name of the beetle, 16—Mr. Carrière's ridiculous statements, 17.	
The Chinch Bug	19
Its disastrous work in 1874, 19—Circular distributed among farmers, 19—Appearance and transformations of the Chinch Bug, 20—The short-winged form, 20—Description of the Chinch Bug and its earlier states, 21—Its past history in America, 22—Its past history in Missouri, 22—Destructive powers of the Chinch Bug, 24—Its injuries in 1874, 24—Its injuries in Missouri in 1874, 25—Food-plants, 26—Time required for the complete development of various insects, 27—Number of annual broods, 27—Its rapid increase, 28—Where the eggs are laid, 28—Flight of the Chinch Bug, 29—Its migrations on foot, 30—Heavy rains destructive to it, 30—Direct remedies, 31—Irrigation, 31—Preventive measures, 32—Burning, 32—Rolling, 33—Manuring and early sowing, 34—Mixing seed, 34—Preventing the migration of the bugs from one field to another, 35—Importance of winter work and combined action, 36—Other possible remedies, 37—Abstaining from the cultivation of grains, 38—Natural enemies, 38; Ladybirds, 39; Lace-wing fly and habits of its larva, 40; The Insidious Flower-bug and the Many-banded Robber, 41—Birds destructive to the Chinch Bug, 41—Discussion of other proposed remedies and preventive measures, 41—The Chinch Bug injurious to stock, 43—Prognosticating, 44—Unnecessary fears, 44—Bogus Chinch Bugs, 45—The False Chinch Bug, 46—The Insidious Flower-bug and the Ash-gray Leaf-bug, 47—The Flea-like Negro-bug, 48—Recapitulation, 49.	
APPENDIX TO THE ARTICLE ON THE CHINCH BUG	51
List of correspondents who replied to the circular, 51—Questions answered by correspondents, 52—Answers given by correspondents, 53.	
The Flat-headed Apple-tree Borer	71
Extent of its ravages in fruit and shade trees, 71—Its natural history, 72—Natural enemies, 73—Chalcid larva, 74—The Cherished Bracon, 75—The Useful Labena, 75—Ants, 76—Remedies, 76—Keeping young trees vigorous and healthy, 77—Coating the trunks and larger branches with soap and other greasy substances, 77—Scraping, 78—Wrapping wire gauze around the trunk, 79—Cutting out the newly-hatched larva, 79.	

Canker Worms	80
Confusion regarding the two species of Canker Worms, 80 — Distinguishing characters of the Spring Canker-worm, 80 — Description of egg, larva and chrysalis of <i>Anisopteryx vernata</i> , 82 — How the Fall Canker-worm differs from the preceding, 83 — Description of egg, larva, and chrysalis of <i>Anisopteryx pometaria</i> , 84 — Practical importance of distinguishing the two species, 85 — Comparative description of the Spring and Fall Canker-worms, 86 — Conclusion, 88 — Extracts from the original essay on the Canker Worm by W. D. Peck, 89.	
The Grape Phylloxera	90
Completion of its natural history, 90 — The true sexual individuals discovered, 91 — Epitome of the life-history of the Grape Phylloxera, 91 — Different forms presented by the species, 93 — Its power to change its habit, 93 — Specific identity of the gall- and root-louse, 94 — Untrustworthy experiments made by the Department of Agriculture, 95 — Proof of the identity of the two forms, 95 — The gall-louse is but a transient form, 96 — Where do the winged females lay their eggs? 96 — The winged female lays the egg wherever she is carried by the wind, 97 — Particular part of the vine chosen by the winged female for laying her eggs, 98 — The true sexual individuals, 98 — Injury done by Phylloxera in America during the year 1874, 99 — Range of the insect in America, 101 — Does it occur in South Carolina and Georgia? 102 — The Phylloxera in California, 103 — Injury done during the year 1874 in France, 103 — Its spread in Europe, 104 — Direct remedies, 105 — Natural enemies, 106 — Susceptibility of different varieties of grape-vine, 103 — Grafting as a means of counteracting the work of Phylloxera, 108 — Underground grafting, 109 — Methods of grafting above ground, 112 — Roots to use as stock, 115 — Varieties to graft, 116 — American grape-vines abroad, 116.	
APPENDIX TO THE ARTICLE ON GRAPE PHYLLOXERA	117
Synopsis of the American species of the genus Phylloxera, 117 — The American Oak Phylloxera, 118; Its natural history, 119 — Description of <i>Phylloxera rileyi</i> and the different forms presented by it, 119 — Further points in its life-history, 120.	
The Rocky Mountain Locust	121
Its natural history, 121 — Method of egg-laying, 121 — The egg, 122 — The newly-hatched locust and its development, 122 — Where the eggs are laid by preference, 123 — The invading swarms are formed by a single species, 124 — Differences between the Rocky Mountain and the Red-legged Locusts, 125 — Mr. Thomas's description of the Red-legged Locust, 126 — Variations, 126 — Measurements of <i>Caloptenus femur-rubrum</i> , 127 — Description of <i>Caloptenus spretus</i> and of its larva and pupa, 129 — Measurements of <i>Caloptenus spretus</i> , 130 — Summing up the difference between the two species, 132 — Chronological history, 132 — Locust invasions in the Old World, 132 — Accounts of earlier locust invasions in America, 133 — Chronological history of the Rocky Mountain Locust, 134 — Earlier invasions, 135 — Data regarding the invasion of 1867, 137 — The invasion of 1873, 141 — The invasion of 1874, 143 — The invasion of 1874 in Missouri, 144 — Questions addressed to correspondents in each county in Missouri regarding the locust, 144 — Summary of the answers given by correspondents, 145 — The invasion of 1874 in Kansas, 148 — in Nebraska, 151 — in Iowa and Minnesota, 153 — in Colorado, 154 — in Dakota and Manitoba, 155 — Flight and ravages of a locust swarm, 156 — Food plants, 158 — Time of appearance of invading	

The Rocky Mountain Locust—Continued.

swarms, 160 — Eastern limits of locust invasions, 161 — Native home of the species, 162 — Explanation of the migratory instinct, 164 — This locust cannot thrive in the Mississippi Valley, 164 — It is a subalpine insect, 165 — What injury may be expected in Missouri in 1875, 166 — Ravages of migratory locusts in the Atlantic States, 167 — Description of the Atlantic Migratory Locust, 169 — Differences between *Caloptenus spretus*, *differentialis* and *atlanis*, 170 — Injury from other, non-migratory, locusts, 171 — The Differential and Two-striped Locusts, 173 — Enemies and parasites, 174 — Birds destroying locusts or their eggs, 174 — The Silky Mite, 175 — The Locust Mite and Dr. LeBaron's description thereof, 176 — The Anonymous Tachina-fly, 178 — The Common Flesh-fly, 180 — Remedies, 181 — Classification of protective measures, 181 — Natural agencies, 181 — Destruction of the eggs, 181 — Destruction of the young wingless locusts, 182 — Driving off the winged locusts, 184 — Further suggestions, 185 — Locusts as food for man, 186 — The popular and scientific names of the insect, 187 — Prairie fires vs. the Rocky Mountain Locust, 189.

APPENDIX TO THE ARTICLE ON THE ROCKY MOUNTAIN LOCUST	191
Letters of three correspondents from Texas and Kansas regarding Locust invasions, 191.	

REPORT VIII.

[Published May, 1876.]

Preface	III
Table of Contents	V

NOXIOUS INSECTS.

The Colorado Potato-beetle	1
---	---

Damage during the year, 1 — Abundant in Atlantic States, 1 — Swarming on Coney Island, 2 — Injuring Egg-plant, 2 — Its scientific name, 2 — Additional enemies, 3 — Eaten by the Crow, 3 — Remedies, 3 — Cost of applying Paris green, 3 — Preparing the poison, 3 — Use of straw as a protection, 4 — Machine for sprinkling, 4 — Machine for brushing off the insects, 4 — Experience with Paris green, 5 — Experiments of Profs. R. C. Kedzie and Wm. McMurtrie show that it may be used with safety, 6 — Trial of other remedies, 6 — The insect's native home, 8 — The theory that it came from the Rocky Mountain Region essentially correct, 10 — Poisonous qualities of the insect discussed, 10.

Canker Worms	12
---------------------------	----

Two species long confounded, 12 — They differ generically; new genus (*Paleacrita*) proposed for one, 13 — The two compared in all stages, 13, 14 — Characters of the genus *Paleacrita*, 17 — Distinguished as Spring and Fall Canker-worms, 17 — Practical considerations from their differences of habit, 18 — Stunting the larvæ does not produce male moths, 19 — Traps recommended, 20, 21.

The Army Worm	22
----------------------------	----

Its generic name, 22 — The term "Army-Worm" applied to various insects, 23 — Past history of the Army Worm, 24 — Known since 1854 in Missouri, 27 — It followed the 1871 conflagration around Peshtigo, Wis., 28 — Its history in 1875; very general all over the country, 28, 29 — Its history in Missouri in 1875, 30 — Sexual differences, 30 — Sexual organs illustrated, 30, 32 — Natural history of the species, 32 — Illustrated in all states, 32, 33 — It occurs in Europe, Asia, New Zealand, and Australia.

The Army Worm—Continued.

lia, 34 — Description of the egg, 34 — Where the eggs are laid, 34 — Conclusions drawn from structure, 36, 37 — When the eggs are laid, 40 — In what state does the insect hibernate?, 43 — Habits of the Worm, 45 — Why it escapes detection when young, 45 — Why it travels in armies, 46 — Time of its appearance, 46 — Are there one or two broods?, 47 — The Fall Army-worm, 48 — How distinguished from the real Army Worm, 48 — Plants preferred by the Army Worm, 49 — Its sudden appearance and disappearance, 50 — It swarms during wet preceded by very dry seasons, 51 — Its natural enemies, illustrated, 52 — Remedies, 54 — Philosophy of winter burning, 54, 55 — Prevention, 55 — Summary of the leading facts concerning it, 56.

The Rocky Mountain Locust.....

57

Previous experience in spring 1867, 57 — Predictions verified, 58 — General outlook in spring of 1875, 60 — Extent of country ravaged, 60 — The outlook in Missouri, 61 — Country ravaged often as bare as in midwinter, 61 — Account by counties, 62 — Atchison County, 62 — Andrew County, 62 — Benton County, 63 — Barton County, 63 — Bates County, 63 — Buchanan County, 64 — Caldwell County, 64 — Cass County, 64 — Clay County, 67 — Clinton County, 68 — Dade County, 68 — DeKalb County, 69 — Gentry County, 69 — Hickory County, 69 — Holt County, 69 — Henry County, 69 — Jackson County, 69 — Johnson County, 72 — Lafayette County, 73 — Nodaway County, 73 — Newton County, 73 — Pettis County, 73 — Platte County, 73 — Ray County, 74 — Saint Clair County, 75 — Vernon County, 76 — Condition of things in other States, 76 — Kansas, 76 — Nebraska, 79 — Iowa, 81 — Minnesota, 81 — Colorado, 84 — Dakota, 85 — Montana, 87 — Wyoming, 88 — Texas, 88 — Indian Territory, 88 — Manitoba, 89 — Damage done in Missouri, 89 — Destitution in Missouri, 91 — Address of Relief Committee from Saint Louis Merchants' Exchange, 93 — Cases of starvation, 94 — The Governor's proclamation, 95 — The locusts not a divine visitation, 97 — Natural history; Mode of molting illustrated, 98 — Habits of the unfledged young, 100 — Directions in which the young travel, 101 — Rate at which they travel, 102 — They reached but a few miles east of where they hatched, 102 — Not led by "Kings" or "Queens," 103 — The species taken for such, illustrated, 103, 104 — The exodus in 1875, 104 — Time of leaving of the winged insects, 104 — Direction taken by the winged insects, 105 — Destination of the departing swarms, 106 — Native home of the species, 109 — Views previously expressed confirmed, 110 — Conditions of migration, 112 — Conditions which prevent the permanent settlement of the species in Missouri, 113 — Modification of the species by climatic conditions, 114, 155 — Definition of the species, 114 — How distinguished in all stages from species most nearly allied, 117 — Experience in spring of 1875, 118 — Contrast in summer and fall, 119 — No evil without some compensating good, 120 — Injury to fruit and fruit trees, 121 — Food plants, 121 — Only one kind of plant not touched under all circumstances, 121 — Changes that followed the locusts, 121 — The widespread appearance of a new grass, ordinarily unnoticed, 122 — Appearance of large Worms, 123 — The Locusts did not return in the fall, 124 — Natural enemies, 124 — Remedies against the unfledged insects, 125 — Artificial means of destroying the eggs, 125 — Various means of destroying the unfledged young, 126 — They are within man's control, 126, — The proper ditch to make, 128 — Machines used in Colorado, 129 — Best means of protecting fruit trees, 130 — How to avert locust injuries, 131 — Pre-

The Rocky Mountain Locust—Continued.

vention, 131 — Legislation, both national and local, 132 — Bills before the Forty-fourth Congress, 133 — Need of a National Entomological Commission, 133 — The bounties offered in Minnesota, 138 — The requisites of a good bounty law, 138, 139 — How a bounty law would work, 140 — Suggestions, 140 — Lessons of year, 142 — Locusts as food for man, 143 — They have been used from time immemorial, and are used extensively at the present day, 145 — The Rocky Mountain species quite palatable, 146 — Mode of preparation, 147 — False opinions and predictions, 148 — Unnecessary alarm caused by other species, 148 — Injuries of native species in 1875, 150 — Locust flights in Illinois in 1875, 151 — They were composed of local species, 152, 153 — Explanation of these flights, 154 — Locust prospects in 1876, 155 — No danger from them in Missouri, 156.

The Grape Phylloxera 157

The injuries not great in Missouri in 1875, 157 — Completion of its natural history, 157 — Where the winged female lays her eggs, 157, 161 — The sexed individuals illustrated, 158 — Description of the true female, 159 — Description of the impregnated egg, 159, 162 — Practical considerations growing out of these latest discoveries, 163 — Decortication of the bark to destroy the impregnated egg, 163 — The insect may be imported from one country to another on cuttings as well as rooted plants, 163 — Best time to attack the root-lice, 163 — Phylloxera ravages in California, 163 — Great destruction around Sonoma, 164 — Need of action by the State authorities, 164 — Occurrence of Phylloxera in the Southern States, 164 — Report of Committee appointed by the American Pomological Society, 165 — Its occurrence in Georgia, 166, 167 — American Grapevines in Europe, 167 — Large demand for our vines, 167 — The American vines flourishing in Southern France where the European varieties perish, 167 — The orders for some varieties exceeded the supply, 168 — Probable future demand, 168.

INNOXIOUS INSECTS.

The Yucca Borer 169

The only North American Butterfly whose larva has the boring habit, 169 — The arbitrary nature of classificatory divisions, 170 — Butterflies and Moths not easily separated, 170 — Biological history of the species, 171 — Illustrations of all states, 171, 172 — Habits of the larva, 171, 172, 181 — Mode of pupation, 172, 180 — Flight of the imago, 173, 181 — Position of wings when the imago rests or walks, 173 — Bibliographical notes, 173 — Detailed descriptions of the different states, 174, 175, 181 — Structural characters illustrated, 175 — Affinities of the species, 176 — It is a true butterfly, belonging to the Hesperians, 178 — Characters of the Castnians contrasted with those of Hesperians, 176, 177, 178 — In classification it is better to widen than restrict in the higher groups, 179 — Enemies of the Yucca Borer, 179 — Concluding remarks, 179 — Unsafe to describe species from mere drawings, 179.

NOXIOUS INSECTS—Continued.

Supplementary Notes on the Army Worm 182

Completion of its natural history, 182 — Oviposition of the Moth described, 183 — Eggs described, 183 — Conclusions previously arrived at verified, 184 — Description of the different larval stages, 184.

REPORT IX.

[Published March, 1877.]

Page.

Preface	III
Table of Contents	V

NOXIOUS INSECTS.

Currant and Gooseberry Worms	1
There are several species having different habits, 1 — Three which may be destroyed by similar methods, 1 — Botanical details as to the Currant and Gooseberry, 2.	
The Gooseberry Span-worm	3
Its natural history, 3 — Most destructive gooseberry insect in Missouri, 3 — Generic nomenclature, 3 — Characters of the moth, 4 — Description of the egg, 4 — Where the eggs are laid, 4 — The insect single-brooded, 5 — How it spreads, 5 — A native species, 5 — Its past history, 5 — It prefers the Gooseberry to the Currant, 6 — The moth is closely imitated by one which greatly differs structurally, 6 — Parasites, 6 — Remedies, 7 — Other currant Span-worms, 7.	
The Imported Currant Worm	7
Belongs to the "False-caterpillars", 7 — The different specific names it has received, 7 — An imported species, 8 — Its introduction and spread, 8 — Independently imported at several eastern points, 9 — Its natural history, 9 — How the eggs are laid, 10 — Nature and habits of the Worm, 11 — Characters of the parent flies, 12 — Preventive measures, 13 — Remedies, 13 — White hellebore the best, 13 — How best used, 14, 15 — The worm is not poisonous, 16 — Natural enemies, 17 — It furnishes a forcible example of Arrhenotoky or the power of producing male offspring without impregnation, 18 — Results of Parthenogenesis in different insects, 18 — It also furnishes an interesting instance of Defunctionation of special parts, 19 — The saw of the female imperfect, compared with others, 20 — Evolutionary bearings of this fact, 21 — Descriptive, 21 — Variation in the antennæ and wing veins, 22.	
The Native Currant Worm	23
Wherein it differs from the imported species, 23 — Its habits, 24 — Where the eggs are laid, 25 — How the winter is passed, 25 — Its occurrence in Missouri, 26 — Remedies, 26 — Descriptive, 26.	
The Strawberry Worm	27
Also a False-caterpillar, 27 — It has a wide range, 27 — How the eggs are deposited, 28 — Character and habits of the Worm, 28 — Remedies, 28 — Descriptive, 28.	
Abbot's White Pine Worm	29
Destructive power of the insects of its genus in Germany, 29 — Evergreens which it prefers, 30 — Habits and characters of the Worm, 30 — Variation in the antennæ, 30 — Characters of the perfect flies, 30, 31 — How the eggs are laid, 31 — Natural enemies, 31 — Remedies, 32 — Descriptive, 32.	
LeConte's Pine Worm	32
A more general feeder than Abbot's species, 32 — The close resemblance of the Worms, 33 — How they differ, 33 — Descriptive, 33 — Other species of the genus, 34.	
The Colorado Potato-beetle	34
Injury in the West in 1876, 34 — Spread of the insect during the year, 34, 35 — Its great abundance on the Atlantic coast, 35 — Rate at which it traveled since 1859, 37 — An average of 88 miles a year, 37 — How it	

The Colorado Potato-beetle—Continued.

traveled, 37; principally in the beetle state, and greatly assisted by man, 37 — Its migrating habit, 38 — Area invaded by it nearly 1,500,000 square miles, 38 — Causes which limit its spread, 38 — Will it reach the Pacific slope?, 39 — How it affected the price of potatoes, 39 — The modification it has undergone, 40 — A mite parasite added to its natural enemies, 41 — Its introduction to Europe, 42 — A living specimen found last summer in the Bremen dock yards, 42 — Could it live and multiply in Europe?, 43 — Action taken by European governments to prevent its introduction, 44 — Consideration of the Kearney "Potato Pest Poison", 45.

The Army Worm.....

47

Further notes and experiments thereon, 47 — Two generations produced annually at Saint Louis, and a probable third generation, exceptionally, 48 — Summary of its natural history, 49.

The Wheat-head Army-worm

50

A new enemy to wheat, 50 — First complaint of it in the East, 51 — First appearance in Kansas, 51 — Habits and natural history, 52 — The egg differs from that of the Army Worm, 53 — Wherein the worm is distinguished from its destructive congener, 54 — Two broods each year, 54 — Natural enemies, 54 — Remedies, 55 — Descriptive, 55.

The Rocky Mountain Locust.....

57

It continues to interest the people of the West, 57 — Previous opinions justified, 57 — The invasion of 1876, 59 — Few in British America, 59 — Condition of things in Montana, 59 — In Wyoming, 59 — In Dakota, 59 — In Minnesota, 60; locusts and alkali soil, 61; good done by Governor Pillsbury, of Minnesota, 61 — In Colorado, 62 — In Iowa, 63 — In Nebraska, 64 — In Kansas, 65 — In Missouri, 66 — Flights in opposite directions at the same time, 66 — Counties in Missouri that were overrun, 67 — Red-legged Locust troublesome in East Missouri, 68 — Detailed reports from counties in Missouri, 68 — Andrew County, 68 — Atchison County, 68 — Barry County, 68 — Barton County, 69 — Bates County, 69 — Benton County, 69 — Buchanan County, 69 — Cass County, 69 — Cedar County, 70 — Caldwell County, 70 — Clay County, 70 — Dade County, 70 — DeKalb County, 70 — Gentry County, 70 — Greene County, 71 — Harrison County, 71 — Henry County, 71 — Hickory County, 71 — Holt County, 71 — Jasper County, 72 — Jackson County, 73 — Johnson County, 73 — Lafayette County, 73 — Lawrence County, 73 — McDonald County, 74 — Newton County, 74 — Nodaway County, 74 — Pettis County, 74 — Platte County, 74 — Polk County, 75 — Ray County, 75 — Saint Clair County, 75 — Vernon County, 75 — In Indian Territory, 76 — In Texas, 76 — In Arkansas, 76 — Destination of the departing swarms of 1875, 77 — They reached into British America, 78 — Source of the swarms of 1876, 79 — Eastern line reached, 80 — Rate at which the insects spread, 80 — Direction of flight, 81 — Influence of wind in determining the course of Locust swarms, 81 — Locust flights east of the Mississippi, 81 — Geographical range of species, 82 — Causes which limit the spread of the Rocky Mountain Locust, 83 — Flights of *Acridium Americanum*, 84 — Does the female of the Rocky Mountain Locust lay more than one egg-mass, 85 — How the eggs are laid, 86 — Philosophy of the egg-mass, 87 — How the young locust escapes from the egg, 88 — How it escapes from the ground, 90 — Additional natural enemies, 91 — Animals which destroy the eggs, 91 — The *Anthomyia* Egg-parasite, 92 — The Common Flesh-fly, 95 — Other undetermined enemies of the eggs, 96 — Insects which destroy the active locusts, 98 — Experi-

The Rocky Mountain Locust—Continued.

ments with the eggs and conclusions therefrom, 99 — Experiments to test the effects of alternately freezing and thawing, 99 — Experiments to test the influence of moisture upon the eggs, 104 — Experiments to test the effects of burying at different depths and of pressing the soil, 104 — Experiments to test the effects of exposure to the free air, 104 — The Omaha Conference, 106 — Remedies and suggestions, 108 — Destruction of the young or unfledged locusts, 108 — Protection of fruit trees, 110 — Legislation, 111 — Act passed by the Missouri legislature, 111 — Acts passed by the Kansas legislature, 112 — Act providing for the destruction of locusts in Minnesota, 114 — Area in which eggs were laid, 116 — Condition of eggs, 117 — Temperature of the winter of 1876-'77, 120 — Prospects for 1877, 121.

INNOXIOUS INSECTS.

The Hellgrammite..... 125

Its curious egg-mass described, 126 — Resembling bird-dung at a distance, 126 — Where laid, 127 — The Egg-burster, 127 — Characters and habits of the newly hatched larva, 127 — Difficulty of rearing it in still water, 128 — The eggs that have been hitherto mistaken for those of *Corydalus*, 128 — They are probably those of *Belostoma grandis*, 128.

The Yucca Borer..... 129

It is single-brooded, 129 — Will thrive in the latitude of Saint Louis, 129 — The larva molts quite often, 129.

CORRECTIONS.

A list of errata is given for each volume, and they are here reproduced with such additional ones as were previously omitted. Where foreign terms were not properly accentuated in the Reports, it was often due to the imperfect "plant" possessed by the State printer. In counting lines the running page title is omitted.

REPORT I.

- Page 8, line 21, *for* being *read* were.
Page 10, line 1, *for* Figure 3,³ *read* Figure 3,².
Page 12, line 20, *for* last *read* 1866.
Page 12, line 3 from bottom, *after* February *add* (1867).
Page 12, line 13 from bottom, *for* verter *read* venter.
Page 14, line 24, *for* hermaphrodite *read* agamic.
Page 14, line 32, *for* females *read* males.
Page 15, line 10 from bottom, *for* muscle-shaped *read* mussel-shaped.
Page 22, line 2 from bottom, *for* pupas *read* pupæ.
Page 30, note, *for* F. *read* T.
Page 31, line 15, *for* 37° *read* 38°.
Page 32, line 4, *for* Kreitz *read* Kreutz.
Page 32, line 14 from bottom, *for* III *read* V.
Page 32, line 7 from bottom, *for* XIII *read* VIII.
Page 38, line 5, *for* Tredeim *read* Tredecim.
Page 47, line 16, *for* far *read* for.
Page 50, line 7, *for* none the less *read* no more.
Page 53, line 28, *for* laid *read* lain.
Page 54, line 4 from bottom, *for* hatch *read* are deposited.
Page 56, lines 5 and 12, *for* to *read* at.
Page 58, line 15 from bottom, *for* *Aspidiglossa* *read* *Aspidoglossa*.
Page 64, line 26, *omit* again.
Page 67, line 11 from bottom, *for* class *read* branch.
Page 76, line 48, *for* Climbing Rustio *read* Climbing Cut-worm Moth.
Page 78, line 46, *for* *unipunctata* *read* *unipuncta*.
Page 86, line 21, and wherever they occur, *for* Guénée *read* Guenée; and *for* Guén. *read* Guen.
Page 87, line 11 from bottom, *for* F. *read* T.
Page 96, note, line 4, *for* West. *read* Wesm.
Page 112, line 3, *for* abbreviated *read* abbreviated.
Page 114, line 1, *after* "insect" *read* (*Stiretrus fimbriatus*, Say).
Page 120, line 30, *after* "Cottonwood" *read* (*Pemphigus vagabundus*, Walsh).
Page 123, last line, *for* *eriosoma* *read* *eriosomatis*.
Page 132, line 16, *for* *ampelopsis* *read* *ampelopsidis*.
Page 133, in heading, *for* Codling *read* Berry Moth.
Page 133, line 24 from bottom, *for* preceding insect *read* Grape Curculio.
Page 134, line 3 from bottom, *for* Part V *read* Part VI.

- Page 142, under the heading, *add* (LEPIDOPTERA TORTRICIDÆ).
 Page 150, line 26, *for* THYRIDOPTERYX *read* THYRIDOPTERIGIS.
 Page 150, line 37, *for* ferruginuous *read* ferruginous.
 Page 154, in the heading, *for* zeas *read* zeæ.
 Page 155, line 13, *for* ZEAS *read* ZEÆ.
 Page 161, line 38, *for* Trallien *read* Trallian.
 Page 166, under heading, *add* (LEPIDOPTERA, PYRALIDÆ).
 Page 171, line 3 from bottom, *for* transformation *read* transformations.
 Page 173, line 3 from bottom, *for* it *read* the more liquid parts.
 Page 174, line 3 from bottom, *for* Solidaga *read* Solidago.
 Page 175, line 32, *add* front *before* wing.
 Page 176, line 21, *for* through *read* into.
 Page 177, line 26, *strike out* in.
 Page 177, line 13, after *coxæ* *read* trochanters.
 Page 178, lines 2 and 3, *for* GELECHIA *read* GELECHLÆ.
 Page 179, line 32, *for* assimilating *read* assimilating.
 Page 179, in heading and line 12, *for* CHICKWEED *read* KNOTWEED.
 Page 179, lines 12, 13, *for* (*Stellaria media*) *read* (*Polygonum aviculare*).
 Page 180, line 7, *for* *Cersium lanceolata* *read* *Cirsium lanceolatum*.

REPORT II.

- Page 8, line 14 from bottom, *for* I have *read* has been.
 Page 8, line 13, from bottom, *before on* *read* largely from Mr. Walsh's previous writings.
 Page 13, line 25, *for* cupable *read* culpable.
 Page 16, line 13, *for* lava *read* larva.
 Page 23, line 6 from bottom, *for* hole *read* holes.
 Page 32, line 17, *for* insect *read* insects.
 Page 35, line 24, *for* *Corimelæna* *read* *Corimelæna*.
 Page 40, line 23, *for* *Ophinsa* *read* *Ophiusa*.
 Page 41, line 25, *for* *Laphrygma* *read* *Laphygma*.
 Page 50, line 5 from bottom, *for* leuca[i]æ *read* leucan[i]æ.
 Page 53, line 12, *for* *perpulera* *read* *perpulchra*.
 Page 56, line 7 from bottom, *for* *Salanum* *read* *Solanum*.
 Page 58, line 19, *for* *copalina* *read* *copallina*.
 Page 59, line 9 from bottom, *for* varigated *read* variegated.
 Page 76, line 4 from bottom, *for* I *read* V.
 Page 76, line 5 from bottom, *for* *Daphni* *read* *Daphne*.
 Page 82, line 25, *for* one *read* our.
 Page 92, line 3, *for* 125-131 *read* 129-131.
 Page 107, line 12, *for* *Naturalista* *read* *Naturaliste*.
 Page 111, line 34, *for* *crysalis* *read* *chrysalis*.
 Page 116, line 4 from bottom, *for* month *read* molt.
 Page 118, line 2, *for* carved *read* curved.

REPORT III.

- Page 6, line 3 from bottom, *for* Rosa *read* Rose; and *for* rosa *read* rosæ.
 Page 7, line 31, *for* *Hyleæctus* *read* *Hylecæctus*.
 Page 25, line 8 from bottom, *for* finely *read* finally.
 Page 28, line 3 from bottom, *for* Holmgreu's *read* Holmgren's.
 Page 30, line 16, *for* the *read* the.
 Page 30, line 16, *for* characterize *read* distinguish.
 Page 47, line 3, *for* Feunde *read* Feinde.
 Page 55, line 50, *for* that *read* than.
 Page 57, line 18 from bottom, *add* c *before* the first h.
 Page 58, line 3 from bottom, *for* *fornudolosus* *read* *formidolosus*.

- Page 64, line 19, for *Bignonio* read *Bignonia*.
 Page 78, note, for I read II.
 Page 95, line 26, for belongs read belonging.
 Page 117, line 5 from bottom, for Harr. read Fabr.
 Page 123, last line, for an read and.
 Page 129, lines 12 and 17, for *Colosoma* and *Calosoma*.
 Page 131, line 13 from bottom, for fauns read fauna.
 Page 135, line 33, for dints read dents; and line 21, for a read b.
 Page 136, line 22, for Guenée read Guenée.
 Page 136, line 33, for Furtsenthum Waldeck read Fürstenthum Waldeck.
 Page 145, line 35, strike out second the.
 Page 146, line 24, add s to transformation.
 Page 150, line 14, at end, add from an.
 Page 151, line 12, for *Cnythia* read *Cynthia*.
 Page 166, line 16 from bottom, strike out first comma.
 Page 166, line 6 from bottom, for *phalangea* read *phalanga*.
 Page 169, line 33, for first i read e.
 Page 170, line 10, for Nnaural read natural.

REPORT IV.

- Page 6, first verse, for grow read grows.
 Page 19, line 8, for 5 read 6.
 Page 20, last line but one, for R read U.
 Page 22, last line but one, for *Aleochora* read *Aleochara*.
 Page 40, line 9 from bottom, for ocular read ocular.
 Page 41, line 59, for Vt. read Ct.
 Page 41, line 15 from bottom, after "Larva" read Length 0.5 inch.
 Page 42, line 5, add a comma after Lepidoptera.
 Page 43, line 6 from bottom, for claud read cloud.
 Page 46, line 29, for edgae read edge.
 Page 46, line under heading, add a comma after Lepidoptera.
 Page 47, line 30, for rhomboidally read trapezoidally.
 Page 53, line 25, and page 54, line 27, for *basillare* read *basilare*.
 Page 53, strike out all after for many in the note.
 Page 59, line 10 from bottom, add winged before female.
 Page 67, line 4 from bottom, for *Cordifolia* read *Riparia*.
 Page 68, line 2 from bottom, for *Oid* read *Oil*.
 Page 75, third line in heading, add a comma after Lepidoptera.
 Page 103, third line of note, for inest read insert.
 Page 105, line 8 from bottom, for *chrysallis* read *chrysalis*.
 Page 110, line 3, for CHALSIS read CHALCIS.
 Page 110, line 29, for *extramatis* read *extrematis*.
 Page 112, in the heading, for Hübn read Drury.
 Page 132, line 19, for Chesnut read Chestnut.
 Page 137, line 1, for Pernyi Silkworm read Perny Silkworm.
 Page 137, under fig. 60, for Pernyi read Perny.

REPORT V.

- Page 7, line 22, for stage read state.
 Page 7, second line from bottom, strike out second the.
 Page 8, explanation of Fig. 1, first line, for and read the.
 Page 9, under Fig. 2, for BEMBEX FASCIATA read VESPA MACULATA.
 Page 9, line 11, for last and read with the.
 Page 11, line 9 from bottom, after worm add moth.
 Page 11, line 3, from bottom, for four read eight.
 Page 12, line 24, for ἐτεροῖς read ἑτερος.

- Page 12, Fig. 5, for *EUSCHISTES* read *EUSCHISTUS*.
 Page 13, line 3 from bottom, for *larvæ* are read *larva* is.
 Page 14, under Fig. 8, for *ÆDIPODA DIFFERENTIALE* read *CALOPTENUS DIFFERENTIALIS*.
 Page 18, line 10 from bottom, for *pollenation* read *pollination*.
 Page 19, line 30, for *Lymexilon* read *Lymexylon*.
 Page 21, line 8 from bottom, for *Townsend* read *Townend*.
 Page 24, line 13, for *ærial* read *aërial*.
 Page 33, in Fig. 15, for *cloroform* read *chloroform*.
 Page 43, line 6 from bottom, after or add in.
 Page 51, line 17, for *J. S* read *S. J.*
 Page 56, line 24, for *how* read *that*.
 Page 52, line 21, for *peteolaris* read *petiolaris*.
 Page 58, line 16 from bottom, for *decrepid* read *decrepit*.
 Page 61, line 18, for *hypertrophized* read *hypertrophied*.
 Page 66, line 13, for *Cordifolio* read *Cordifolia*.
 Page 67, line 27, for *with* read *to*.
 Page 67, line 28, after and add to offer.
 Page 83, line 13 from bottom, for *who* read *as*.
 Page 85, line 17 from bottom, after *fecundation* add either the.
 Page 85, line 18 from bottom, strike out either and after *female* add would.
 Page 86, line 2 from bottom, for *and* read *und*.
 Page 90, line 17, for *had* read *has*.
 Page 100, last line, add a comma before *say*.
 Page 101, line 10, for *nole* read *noli*.
 Page 103, line 9, for *Caryæ* read *caryæ*.
 Page 113, line 40, for 19 read 41.
 Page 115, for *exerted* read *exserted*.
 Page 120, line 25, for *regulary* read *regularly*.
 Page 126, line 4 in note, for *quercili* read *querceti*.
 Page 126, line 5 in note, for *pithicium* read *pithecium*.
 Page 129, line 14, omit color of the.
 Page 139, for *Papineau* read *Popenoe*.
 Pages 140 and 141, wherever *Belvosia* occurs read *Belvoisia*.
 Page 156, line 6, for *consumes* read *has consumed*.

REPORT VI.

- Page 8, last line, for 1874 read 1873.
 Page 12, line 13, for *Sisimbrium* read *Sisymbrium*.
 Page 12, line 25, for *osciamus* read *oscyamus*.
 Page 12, lines 13, 14, for *Poligonum* read *Polygonum*.
 Page 27, line 2 from bottom, for *peorus* read *pecoris*.
 Page 27, line 15 from bottom, for *vireus* read *virens*.
 Page 28, last line, for XV read XVI.
 Page 35, line 3, for three read four.
 Page 37, line 16, for *first* by read *be*.
 Page 42, line 11, for the read *certain*.
 Page 43, line 26, strike out to be presently treated of.
 Page 47, remove "Telegraph" from "Summer grape" § to that of "Northern Fox."
 Page 51, line 7 from bottom, for *insidious* read *insidiosus*.
 Page 53, line 18, for *Maguin* read *Mégnin*.
 Page 82, line 5 from bottom, for *New* read *West*.
 Page 87, line 3 from bottom, for *Bignonio* read *Bignonia*.
 Page 92, line 6, for *Callimorpa* read *Callimorpha*.
 Page 94, line 14 from bottom, for *point* read *joint*.
 Page 100, line 31, omit comma after *lardarius*.

- Page 108, line 28, *for orage read orange.*
 Page 111, line 6, *for perspicillata read tripunctata.*
 Page 118, last line, *for Phytopoga read Phytophaga.*
 Page 136, line 15, *for Rosel von Rösenhof read Röscl von Rosenhof.*
 Page 141, line 9, *after found add that.*
 Page 150, line 9, *for pictures of read imprints on.*
 Page 154, line 6, *strike out †.*
 Page 154, line 16, *for it is read they are.*
 Page 156, line 8, *after and add more.*
 Page 162, line 10, *for elytram read elytrum.*

REPORT VII.

- Page IV, line 9, *for contemptibly read contemptuously.*
 Page 1, line 10, *after and read invaded the.*
 Page 5, line 16, *for State read state.*
 Page 7, line 7, *for calubrine read colubrine.*
 Page 11, line 32, *for stoma read stomata.*
 Page 11, line 33, *for dilutent read diluent.*
 Page 11, line 37, *for J read S.*
 Page 12, line 13, *for W. K read R. C.*
 Page 17, last line, *for Dep. de l'Hérault read Dép. de l'Hérault.*
 Page 21, line 14 from bottom, *for LENCOPTERUS read LEUCOPTERUS.*
 Page 39, under Fig. 6, *for TTIM read TRIM.*
 Page 52, line 14, *for McWallie read McNallie.*
 Page 52, line 46, *for Princeton read Purinton.*
 Page 75, line 32, *for breed read bred.*
 Page 80, line 7 from bottom, *add a comma before and after pometaria.*
 Page 81, last line, *for nidi read nidus.*
 Page 94, in the sub-head, *for GALL-INHABITING read ROOT-INHABITING.*
 Page 99, line 7, in note, *for nerves read tracheæ.*
 Page 108, line 17, *for two read too.*
 Page 117, line 15, *for V read IV.*
 Page 118, line 17 from bottom, *for hight read height.*
 Page 147, line 20, *for 1873 read 1866.*
 Page 162, line 20, *for larva read lava.*

REPORT VIII.

- Page III, line 13 from bottom, *add 1 before the 3.*
 Page 7, line 26, *for copper read soda.*
 Page 22, in notes, *transpose the * and †.*
 Page 34, line 6, *for tulietes read tuélites.*
 Page 34, line 11, *for three-hundredths read two-hundredths.*
 Page 37, under Fig. 23, *for exerted read exserted.*
 Page 38, line 6, *for glass read grass.*
 Page 52, line 1 in note, *for Doliconyx read Dolichonyx, and for orizivora read oryzivora.*
 Page 53, line 32, *for veridascens read viridascens.*
 Page 98, second line, in explanation of Fig., *for e read c and for c read e.*
 Page 100, line 2, *after they add are still imperceptible; in the third stage (after second molt) they.*
 Page 100, line 4, *for third read fourth, and for second read third.*
 Page 100, line 7, *for fourth read fifth, and for third read fourth.*
 Page 100, line 8, *for fourth read fifth and for fifth read sixth.*
 Page 114, line 7 from bottom, *for distingulsh read distinguish.*

- Page 115, line 5, *after* histories *add* a comma.
Page 115, line 5 from bottom, for *Pesotettix* read *Pezotettix*.
Page 121, line 18, *after* limbs *add* and.
Page 149, under Fig. 46, for larva read pupa.
Page 150, line 10, for *gran* read *granu*.
Page 154, line 4 from bottom, for sheli read shell.

REPORT IX.

- Page 6, line 26, insert after "moth" (*Euphanessa mendica*, Walk.).
Page 15, line 3, for entite read entire.
Page 29, in explanation of cut, for Abbott's read Abbot's.
Page 50, explanation of cut, for *e* read *c*.
Page 50, line 3 from bottom, for Hubner read Hübner.
Page 54, last line, in place of the comma, write is.
Page 55, line 1, for the other read the second.
Page 55, line 9 from bottom, for m. m read mm.
Page 55, line 7 from bottom, strike out the on.
Page 56, line 1, for m. m read mm.
Page 56, line 2, for the last and read anal.
Page 56, line 32, commence a new ¶ with "Chrysalis" and italicize it.
Page 57, in the heading for *Spretus* read *spretus*.
Page 58, line 14, strike out have.
Page 87, strike out the *g* in line 17 and also in figure.
Page 89, line 13, strike out the *i* after embryon.
Page 90, last line, for ambion read amnion.
Page 98, line 11 from bottom, for *Compoplex* read *Campoplex*.
Page 98, line 6, add a comma before De Geer.
Page 98, note *, for *Bastardii* read *Bastardi*.
Page 98, under Fig. 28, for BASTARDII read BASTARDI.

NOTES AND ADDITIONS.

Under this head it is not my purpose to publish the many additional notes of observations which have been made by myself and others on the various insects treated of in the reports ; but rather to indicate a few of the more important facts, especially such as are unpublished and bear on life-histories left incomplete. As, in preparing the reports, the older and better known generic nomenclature was almost uniformly employed, it is thought advisable to indicate in this bulletin the more recent nomenclature, and this is accordingly done either in these "Notes" or in connection with the reproduced "Descriptions of New Species" which follow :

HYMENOPTERA.

STIZUS GRANDIS Say (Rep. I, p. 27, Fig. 12)—This has been shown by Mr. W. H. Patton (Bull. U. S. Geol.-Geog. Survey, vol. V, p. 342) to be only a variety of *speciosus* Drury, which is the type of the genus *Sphecius* Dahlb.

CRYPTUS EXTREMATIS Cress. (Rep. IV, p. 111)—The questions in regard to the character of *C. samiae* Pack. have been settled by Dr. Hagen from an examination of the types (Bull. Buff. Soc. Nat. Sci., II, 206 ; 1875) confirming the conclusion which I came to. In Bulletin No. 3 of the Commission (p. 47) I have suggested that *extrematis* should sink as a synonym of *samiæ*, because two species (one of which is *nuncius*) were combined under it in the original description.

MICROGASTER MILITARIS Walsh (Rep. I, p. 89 and subsequently)—This is an *Apanteles* (See my Notes on N. A. Microgasters, etc. (Extr. from Trans. St. Louis Ac. Sc. IV), p. 19.)

ANTIGASTER MIRABILIS Walsh (Rep. VI, p. 162)—Mr. L. O. Howard has shown (*Can. Ent.* October, 1880. p. 209, and February, 1881, p. 31) that the habit of rolling back is not uncommon in the *Eupelmides*, and that *Antigaster* cannot well be separated from *Eupelmus* as at present understood.

COLEOPTERA.

CARABID LARVÆ (Rep. IX, p. 97)—The second larva mentioned on this page was subsequently reared by me to the perfect state and proved to be *Amara obesa* Say. It will be found figured and described in the First Report of the Commission (p. 290).

LEBIA GRANDIS, Hentz (Rep. III, p. 100)—This belongs to Chaudoir's genus *Loxopeza*.

LEBIA ATRIVENTRIS Say (Rep. VIII, p. 3)—Belongs to Chaudoir's genus *Loxopeza*.

HIPPODAMIA MACULATA, DeGeer (Rep. I, p. 112 and subsequently)—Now referred to Mulsant's genus *Megilla*. It does not appear that this species occurs also in Europe as stated in the text.

COCCINELLA MUNDA Say (Rep. II, p. 25)—This is now considered synonymous with *Cycloneda sanguinea* Linn.

COCCINELLA PICTA Randall (Rep. V, p. 101)—Now known as *Harmonia picta*.

MYSIA 15-PUNCTATA, Oliv. (Rep. IV, p. 18)—This has been referred to the genus *Anatis*, Mulsant.

LACHNOSTERA QUERCINA, Knoch. (Rep. I, p. 156)—This is synonymous with *fusca* Fröhlich, which has priority. The fungus affecting it (p. 158 and Rep. 6, p. 125) is *Cor-dyceps ravenelii* Berkeley. (See *American Entomologist*, III, p. 139.)

CHAULIOGNATHUS PENNSYLVANICUS, De Geer (Rep. I, p. 57 and subsequently)—This is now known as *Ch. americanus* Forst., the latter name having priority. For an account of the eggs and young larvæ, see Second Report of the Commission, p. 261.

SAPERDA BIVITTATA Say (Rep. I, p. 42)—This is now admitted to be a synonym of *candida* Fabr. For a correct description of the eggs and mode of oviposition, see an article by me in *New York Weekly Tribune*, Feb. 20, 1878.

BRUCHUS PISI Linn. (Rep. III, p. 44)—This name of the 12th edition of Linnaeus's "Systema Naturæ" gives way in modern catalogues to *pisorum* L. of the 10th edition.

FIDIA VITICIDA Walsh (Rep. I, p. 32)—This species is not mentioned by Crotch in his "Materials for the Study of the Phytophaga of the U. S." (Proc. Ac. Nat. Sc. Phil., 1873), but his *Fidia murina* (l. c. p. 33) is undoubtedly synonymous with Walsh's *viticida*, the latter name having priority by several years. In Crotch's "Check list" this species is also omitted, but the *Fidia vitis* Walsh in the "Omissions" to that list (p. 127) is probably meant for *viticida*.

HALTICA CHALYBEA, Illiger (Rep. III, p. 79)—This belongs to the genus *Graptodera* Chev.

HALTICA CUCUMERIS Harris (Rep. I, p. 101)—This is now referred to the genus *Epitrix*, Foudras.

PHYSONOTA QUINQUEPUNCTATA Walsh & Riley (Rep. II, p. 59)—This is synonymous with *Ph. unipunctata* (Say), there being no question as to the specific identity of the two, both having been bred by Mr. F. H. Chittenden, of Ithaca, N. Y., from larvæ on a wild sun-flower (*Helianthus*).

CASSIDA NIGRIPES Oliv. (Rep. II, p. 63)—The eggs of this species are much like those of *aurichalcea* (Rep. II, Fig. 31) in size, form and color, though the spine-like appendages break off more easily. They may, however, be distinguished by being larger (1.6^{mm} long without projections), having, in fact, nearly double the bulk, and by the flat posteriorly projecting piece which bears the spine-like appendages being generally greatly developed so as sometimes to extend beyond the apex fully one-third the length of the whole egg. Sometimes this piece divides distinctly into three spines, but in other cases it is quite blunt.

CASSIDA BIVITTATA Say (Rep. II, p. 61)—The eggs of this species are pale and ovoid, just 1^{mm} long, but invariably covered with a yellowish secretion which dries and spreads out each side, and this by a black excrementitious material which gives the egg from above the appearance of an ovoid bit of excrement flattened on the adhering side. The eggs are laid singly or in twos or threes.

CASSIDA AURICHALCEA Fabr. (Rep. II, p. 62)—This is now referred to the genus *Coptocycla* Chev.

CASSIDA PALLIDA Herbst (Rep. II p. 62)—This is now recognized as a synonym of *Coptocycla aurichalcea* (Fabr.).

COPTOCYCLA GUTTATA, Oliv. (Rep. II, p. 63)—The eggs of this species, which I have often since observed, are deposited singly or in twos, threes or fours. They are rather more than 1^{mm} long, of the same general form and character as those of *Cassida bivittata*, but more narrow and elongate. The color is pale yellowish and translucent. The egg is always covered with a viscid fluid which dries to form a transparent covering verging to fulvous or gamboge in color. This covering almost always spreads out on each side of the egg in ray-like ridges, those on each side parallel and slightly oblique, and whenever the egg is single these ridges are remarkably regular and have a neat appearance. There is occasionally on the top of this a varying amount of ex-

crement. The structure of the covering is similar to that found in the egg of *Cassida texana* Cr. (which feeds on *Solanum eleagnifolium*), where, however, the ribs are finer and transverse, and there is no excrementitious covering. The newly hatched larva of *guttata*, like that of the other species is whitish, strongly recalling in general appearance an ordinary mite, the head not being concealed as it subsequently is, the hairs at the tip of the legs being frequently clavate or knobbed, and resembling those on the young of many Coccids. The marginal spines and the anal fork are quite well developed but simpler than in the subsequent larval stages. This newly hatched larva is quite nimble and crawls easily over glass.

DELOYALA CLAVATA, Oliv. (Rep. II, p. 56)—Now referred to the genus *Coptocycla*.

BLISTER-BEETLES (Rep. I, p. 96 ff.)—The larvæ feed on locust eggs. For account of their larval economy see my paper "On the larval Characters and Habits of the Blister-beetles," etc., Trans. Ac. Sc. St. Louis III, p. 544 ff.: also Reports of the Commission I, p. 292 ff.; II, 262 ff. Remarks on synonymy are also there given, but the following may be repeated.

LYTTA CINEREA Fabr. (Rep. I, p. 97)—This is now known as *Macrobasis unicolor* (Kirby).

LYTTA MURINA Lec. (Rep. I, p. 98)—This is a color variety of *Macrobasis unicolor*.

LYTTA MARGINATA Fabr. (Rep. I, p. 98)—This is believed by Horn to be a color-variety of *Epicauta cinerea* (Forst.).

LYTTA ATRATA Fabr. (Rep. I, p. 98)—This is the *Epicauta pensylvanica* (De Geer) of Crotch's List.

ANTHONOMUS PRUNICIDA, Walsh. (Rep. III, p. 39) — Upon this species, which is a synonym of *scutellaris* Lec., Dr. Leconte has since founded the genus *Coccotorus* (Proc. Am. Philos. Soc. vol. XV, 1876, p. 193).

CONOTRACHELUS NENUPHAR, Hbst. (Rep. III, p. 127, note) — The phytophagic variety of this species from Walnut and Butternut has since been characterized by Dr. Leconte as a distinct species, *C. juglandis* (Proc. Am. Philos. Soc. vol. XV, p. 226).

CÆLIODES INÆQUALIS, Say (Rep. I, p. 128) — Dr. Leconte has since founded upon this species the genus *Craponius* (Proc. Am. Philos. Soc. vol. XV, 1876, p. 268). The egg of this snout-beetle is quite large, bright yellow in color and deposited in a cavity half as large as the beetle, though the puncture leading to it is small. The lateral angularities or tubercles of the joints, as described by Walsh, are quite characteristic, and the dorsal view in my figure, given to show them, conveys a somewhat false impression of the larva, which is more or less curved, and has the general characteristics of Curculionid larvæ. The figure is rather more attenuated than it should be. That the beetle hibernates I have since proved beyond question.

BARIDIUS TRINOTATUS Say (Rep. I, p. 93) — Dr. Leconte (Proc. Am. Philos. Soc. XV, 1876, p. 287) has since established for this and two allied species the genus *Trichobaris*.

SPHENOPHORUS ZEÆ Walsh (Rep. III, p. 59) — This has been previously described by Mr. Uhler as *S. sculptilis* (Proc. Ac. Phil. VII, 1855, p. 416).

SPHENOPHORUS PULCHELLUS Schœnherr (Rep. III, p. 60) — As intimated in the footnote on the same page, this species is synonymous with Say's *S. 13-punctatus*, for which species and for *Sphenophorus pustulosus* Gyllh. Dr. Leconte has established the genus *Rhodobænus* (Proc. Am. Philos. Soc. vol. XV, 1876, p. 332). I have reared both, and also intermediate forms, from Helianthus in Texas, and Ambrosia in Missouri.

SCOLYTUS CARYÆ Riley (Rep. V, p. 107) — Dr. Leconte (Proc. Am. Phil. Soc. XV, 1876, p. 371) has since decided that 4-spinosus Say is the ♂ of this species, and Say's name consequently obtains.

LEPIDOPTERA.

PAPILIO PHILENOR Drury (Rep. II, p. 116) — Referred by Scudder to Hübner's genus *Laërtias*. For further notes and description of the egg and young larva, see *Canadian Entomologist*, January, 1881, p. 9, and *American Naturalist*, April, 1881, p. 327.

DANAIS ARCHIPPUS, Fabr. (Rep. III, p. 143)—For further facts respecting the swarming and migrations of this butterfly, see the *American Entomologist* (III, p. 101), and for a fuller and more accurate account of the mode of pupation, see my paper on the "Philosophy of the Pupation of Butterflies and particularly of the Nymphalidæ" (Proc. Am. Ass. Adv. Sc. vol. XXVIII, 1880).

ÆGERIA ACERNI, Clem. (Rep. VI, p. 110)—Mr. D. S. Kellicott has an interesting article in the *Canadian Entomologist* for January, 1881, on the Ægerians inhabiting the vicinity of Buffalo, N. Y., in which he states that the chrysalis of this species in his locality does not agree with my description as "unarmed," if that description refers to the dorso-abdominal teeth. A reëxamination of my specimens shows that my statement applies to the absence of these teeth. It is, however, possible that there is some variation in this regard and that the eastern specimens from the Hard maple differ from the western ones from the Soft maple in having the teeth as indicated by Mr. Kellicott.

ARCTIA ISABELLA, Smith (Rep. IV, p. 143)—Referred to *Pyrrharctia* Packard. For further account of larval variation and parasites, see *American Entomologist*, III, p. 134 (June, 1880).

HYPHANTRIA TEXTOR Harr. (Rep. III, 130)—There is no doubt in my mind, from frequent breeding of specimens, that this is synonymous with *cunea* Drury and *punctata* Fitch, which are but varieties, Drury's name having priority.

CALLIMORPHA FULVICOSTA, Clem. (Rep. III, 132)—Grote and Robinson give the synonymy of this species in their "List of Lepidoptera of N. A.," etc., *lecontei* Boisd. having priority. The late Jacob Boll bred all the forms from larvæ feeding on the same species of plant.

SAMIA COLUMBIA Smith (Rep. IV, p. 107)—Mr. Herman Strecker has given a beautiful figure of the male of this species in his "Lepidoptera Rhopaloceres and Heteroceres, etc.," 1875 (Pl. XII, Fig. 3), and Mr. F. B. Caulfield has described and figured the larva (*Canadian Entomologist*, X, p. 41, 1878) showing that it is structurally identical with that of *cecropia* and differs only in the intenser green of the body, in the lateral tubercles and bases of the others being white instead of pale blue and in the upper thoracic tubercles being of a deeper coral-red. It accords more with the *cecropia* larva in the fourth stage. It is placed as a good species in Grote's "List of N. A. Platypterices," etc. (Am. Phil. Soc., 1874), but I am still of opinion that it should not be considered a distinct species but simply a well-marked local color-variety worthy of name. There is great variation in color, whether of the larva, cocoon or imago, in *cecropia*.

CALLOSAMIA ANGULIFERA, Walker (Rep. IV, p. 122, note)—This is still considered a good species by systematists. Mr. Jno. Akhurst, of Brooklyn, N. Y., informs me that he finds it rather constant from larvæ which seem to differ in no respect from those of *promethea*, but which feed on the Tulip tree (*Liriodendron tulipifera*), and make the cocoon near the ground without pedicel. I learn from Dr. Packard that Mr. Uhler has bred both it and *promethia* from the same lot of larvæ.

CLISIOCAMPA SYLVATICA Harr. (Rep. III, 121)—This is now referred to *disstria* Hübn., which has priority.

AGROTIS INERMIS Harr. (Rep. I, p. 72)—This is now recognized to be identical with the European *A. saucia* Treitschke.

NOCTUA CLANDESTINA Harr. (Rep. I, p. 79)—An *Agrotis*.

AGROTIS TELIFERA Harr. (Rep. I, p. 80)—This is now recognized as the European *A. ypsilon* Hüfn. = *A. suffusa* (S. V.) = *A. ortonii* Pack.

AGROTIS SUBGOTHICA Harr. (Rep. I, p. 81)—The moth represented under this name at Fig. 29, *a*, has since been described by Grote as *A. herilis*, and that at Fig. 29, *b*, has since been described by Lintner as *A. tricola*. (Notes on some N. Y. Noctuidæ, Ent. Cont. III in Rep. N. Y. St. Mus. Nat. Hist., 1872, p. 159.)

AGROTIS JACULIFERA Guen. (Rep. I, p. 82)—This is the true *subgothica* of Haw. (See Grote, List of Noctuidæ of N. A., Bulletin Buffalo Soc. Nat. Sc. II, 1874, and Lintner l. c.)

AGROTIS DEVASTATOR, Brace (Rep. I, p. 83)—Grote refers it to *Hadena*.

CELÆNA RENIGERA Stephens (Rep. I, p. 86)—Referred by Grote to *Hadena*. Specimens in the Fitch collection marked with names (evidently from Walker) *infecta*, *egens*, *defectua*, *subcadens* ? and *murcimaculata* seem to be all synonyms and mere variations.

PRODENIA AUTUMNALIS Riley (Rep. III, p. 116 and subsequently)—As stated in the 8th Report (p. 48) this in the more typical form is recognized as *Laphygma frugiperda*, Sm. & Abb. The variety *obscura*, as Prof. Zeller, who has seen it, informs me is so near the European *exigua* Hübn. that it is not easily distinguished.

PRODENIA COMMELINÆ, Sm. & Abb. (Rep. I, p. 88, and III, p. 113)—Dr. Leon F. Harvey (Bull. Buff. Soc. Nat. Sci., vol. II, pp. 274, 275; 1875) has since proposed specific names for two of the forms hitherto considered to be but varieties of *commelinæ*. The moth represented at Fig. 48, c, of the Third Report, is named by him *flavimedia*, that at Fig. 48, b, *lineatella*, the true *commelinæ*, being a larger species. From larvæ with the series of black triangles bordered exteriorly by a yellow line (such as are represented on Plate I, Fig. 12 of Rep. I, and at Fig. 48 a of Rep. III) I have bred the *flavimedia*. But larvæ found on cotton in the Southern States, and differing in having black triangles on the second joint only, and also varying greatly in coloration, have produced the same moth. Abbot's figure of the larva of *commelinæ* shows the full series of black triangles, but without any yellow exterior line.

GORTYNA NITELA Guen. (Rep. I, p. 92)—I have proved by breeding that *G. nebris* Gn. is but a large, southern form of this species. In the Southern States it is most common in stems of *Ambrosia trifida*, often producing a swelling or pseudo-gall. Both forms are indiscriminately bred with intermediate variations. See an article by Miss E. A. Smith (7th Report on the insects of Illinois, Cyrus Thomas, pp. 112-114) for additional food-plants and the habit of the younger larvæ to infest wheat-stalks, corn, etc. See also Am. Ent. I, p. 252; my "Potato Pests" (Orange, Judd & Co., 1877, p. 91) and *Prairie Farmer*, August 11, 1877. The insect normally pupates in the stem and when infesting thin stalks like those of most cereals and blue-grass (in which it is also found) often of necessity leaves one stalk for another.

ANOMIS XYLINA, Say (Rep. II, p. 37; VI, 17)—This has since been referred by Grote to Hübner's *Aletia argillacea*, which has been generally adopted. See Bulletin 3 of the Commission on the Cotton Worm. While it will doubtless be found convenient in future to separate it from the other species of the genus *Anomis*, and Hübner's generic name may therefore obtain, I must confess, after a careful examination of Hübner's figure of *argillacea*, to grave doubts as to the correctness of Grote's reference thereto of our Cotton-worm Moth (*xylina*, Say). Hübner's figure lacks several of the most constant characteristics of *xylina*. It is fulvo-testaceous shaded with brown, with the under side bright yellow. It lacks the three white specks on primaries and has a dark (orbicular?) spot in place of the outer one. It has a large white circular spot with black annulus in place of the dusky elongate discal spot with its double pupil. The wavy lines are almost black and differ in form; the fringes are unicolorous, and the abdomen is narrower. The figure more nearly represents in fact a species which I have received from Bahia, Brazil, and which differs from *xylina*, though the larva (also quite different) feeds on cotton.

We are all inclined to follow determinations of those who make a specialty of any group, but after due allowance for faulty coloring in Hübner's figure, I am constrained to believe that in this instance Mr. Grote has been in fault.

CANKER-WORMS (Rep. VIII, p. 12)—For additional remarks as to the generic characters of the two Canker-worms, see my paper "On the differences between *Anisopteryx pometaria* Harr. and *Anisopteryx æscularia* W. V., with remarks on the genus *Paleacrita*. (Trans. Ac. Sc. St. Louis, Vol. III, p. 573 ff.)

GALLEREA CEREANA, L. (Rep. I, p. 166)—This is the *mellonella* L. of the 10th edition Syst. Naturæ.

PEMPELIA GROSSULARIÆ Packard (Rep. I, p. 140)—The European *Zophodia convolutella* Hübn. (*Phycis grossulariella* Treitschke), which has precisely similar habits, closely resembles this species. In 1871 I compared it with this last in Mr. Stainton's collection and with specimens received from Prof. Zeller and could detect no essential differences. The European specimens are slightly larger, with broader wings and usually clearer, paler gray coloring. Colorational markings are, however, very variable in specimens from both sides of the Atlantic.

P. grossulariæ Packard was subsequently described by Grote as *Dakruma turbatella* (Bull. U. S. Geol.-Geog. Survey, IV, No. 3, p. 702; 1878). *Dakruma* seems to differ from *Zophodia* in nothing but the absence of the basal portion of the subcostal vein and possibly, although this character is not mentioned by Grote, in the recurved palpi. According to the synoptical table given by Heinemann, *grossulariæ* would fall in the genus *Stenoptycha*, distinguished from *Zophodia* by the recurved palpi. We may well question the generic value of this character, for different authors describe it quite differently: thus, Heinemann describes the palpi of *Stenoptycha* and *Homeosoma* as recurved, whereas Grote describes them as porrect in these two genera, if we accept his statement that *Honora* Grote is to be considered a section of *Stenoptycha*: there appears also to be a difference in position in specimens of the same species, according as the palpi are heavily scaled or have lost the scales. From the known individual variation in the venation of these and other moths, especially in the hind wings, we cannot attach any specific, much less any generic, value to the slight difference in the subcostal vein of *Dakruma* noted above. Moreover, authentic specimens of *grossulariæ* do not appear to possess this character of *Dakruma*. I am, therefore, of opinion that a study of sufficient material from both continents will prove the two specifically identical, or at the most that our American insect is a variety, and that *Dakruma* will not obtain. Packard is of this opinion, as in the later editions of his *Guide* the species is called *Myelois convolutella*.

PENTHINA VITIVORANA Packard (Rep. I, p. 133)—This is identical with a European insect having the same habits. It was first described over a century since by Schiffermiller & Denis as *Tortrix botrana*, and has been referred to various genera since, and finally to *Eudemis* Hübn., so that the insect should be known as *Eudemis botrana* (Schiff.). *Conchylis ambiguella* (Hübn.) has very similar habits in Europe. See Nördlinger's "Die Kleinen Feinde der Landwirthschaft," p 424 ff. It is the *Lobesia botrana* of the later editions of Packard's *Guide*.

EURYPTYCHIA SALIGNEANA Clem. (Rep. II, 134).—This according to Prof. Fernald, who has seen the type, is the same as Clemens's *Hedya scudderiana* (Proc. Acad. Sci. Phila., 1860, p. 358), the description of which is very brief and presumably taken from a female. The genus *Euryptychia* (Proc. Ent. Soc. Phila. V, 140) is founded on the male, which has a broad fold extending to the middle of costa on the primaries and covering up a pencil of yellowish hairs. Zeller subsequently redescribed it as *Pædisca affusana* (Beiträge, etc., pt. III, p. 101 [307]). From a comparison of female specimens I am led to believe that this is the same species that is commonly known in Europe as *Spilonota roborana* Schiff., though in Staudinger and Wocke's Catalogue *cynosbana* Fabr., described in 1875, is given the priority and *aquana* Hübn. is placed as a synonym. The obliquity of the edge of the basal dark patch and the details of the ocellated spot upon which species have been separated, I find to be variable.

The insect in Europe is known to feed on the leaf-buds of the rose. I have abundant proof that in this country it is not a gall-maker, but, as was inferred in the Report, an inquiline. I have found its larva feeding upon the flowers as well as amid the terminal leaves of the Golden-rod, and have also found it in other galls. When feeding in the more exposed positions it generally has a carneous or rosy tint.

ANCHYLOPERA FRAGARIÆ W. & R. (Rep. I, 142)—This has been referred to *Phoxopteris comptana* Fröhl., and while the two very closely resemble each other Prof. Fernald informs me that he yet believes *fragariæ* to be distinct.

ÆTA COMPTA, Clem. (Rep. I, p. 151)—Notwithstanding Mr. Grote doubts the identity of this insect with Cramer's *Phalæna punctella*, there is no question in my mind about it, and I entirely agree with Zeller, who makes also the *Tinea pustulella* Fabr. a synonym (Beitr. z. Kenntn. N. A. Nachfalter II, p. 28). It was first described in this country in 1856 by Fitch as *Deiopeia aurea* (3rd Rep. Ins. N. Y., p. 168.) See also "Zygænidæ and Bombycidæ of N. A." by R. H. Stretch, 1872, pp. 159 and 241.

The egg of this insect is one of the most singular Lepidopterous eggs with which I am familiar. I have found it numerous in the South in midsummer. It is 0.9^{mm} long, soft and plastic so as to be variable in form; but when laid (as it often is) on the web which the young larvæ make, where it takes on the more natural form, it is ovoid, somewhat compressed, with frequently a median ridge and one end narrowed and produced into a short neck. The color is cream-yellow and the delicate shell is corrugulate. It is laid singly and generally slightly attached by the broad side to the side of the mid-rib of the tenderest leaves, and its contact (by virtue, doubtless, of some poisonous liquid with which it is laid) causes a well defined swelling of the leaf-vein.

The species is placed among the *Zygænidæ* in Grote and Robinson's *List*, and has evidently more affinities therewith than with the *Teneidæ*.

PRONUBA YUCCASELLA Riley (Rep. V, p. 150 and subsequently)—For further facts regarding this species, see my papers in Trans. St. Louis Ac. Sc. III, p. 568; *American Entomologist* III, pp. 141, 182, 293, and also a paper read before the American Association for the Advancement of Science at Boston, Aug., 1880, and to be published in the Proceedings of the Association for that year.

PTEROPHORUS PERISCOLIDACTYLUS (Rep. III, p. 65)—This belongs to the genus *Oxyptilus*, Zeller.

HETEROPTERA.

ARMA SPINOSA Dallas (Rep. II, p. 113 and subsequently)—Now referred to Stål's genus *Podisus*.

EUSCHISTUS PUNCTIPES, Say (Rep. IV, p. 19 and subsequently)—This is now known as *Euschistus variolarius* Beauv., this last having priority over Say's name.

COREUS TRISTIS, De Geer (Rep. I, p. 113 and subsequently)—Belongs to Amyot & Serville's genus *Anasa*.

MICROPUS LEUCOPTERUS, Say (Rep. II, p. 15 and subsequently)—Now referred to Burmeister's genus *Blissus*.

ANTHOCORIS INSIDIOSUS, Say (Rep. II, p. 27 and subsequently)—Belongs to Fieber's genus *Triphleps*.

REDUVIUS RAPTATORIUS Say (Rep. I, p. 114)—Belongs to *Sinea*, Amyot & Serv., and is synonymous with *diadema* Fabr.

HARPACTOR CINCTUS Fabr. (Rep. I, p. 114 and subsequently)—Belongs to Stål's genus *Milyas*.

HOMOPTERA.

CICADA SEPTEMDECIM (Rep. I, p. 18)—This orthography, used in the Reports, is grammatically correct, but I find that Linnæus himself wrote *septendecim* (*Systema Naturæ*, Tom I, Pars II, 12th Ed. Stockholm 1767). Fitch used both forms of spelling, but Westwood, Harris and most other authors follow Linnæus, and *septendecim* is, therefore, preferable. As to whether the 17 and 13-year broods should be considered specifically distinct, I am still of the opinion expressed in the First Report that the insects should not be looked upon as distinct species, but that *tredecim* Riley should rather be considered a race, or as Walsh (in a letter to Charles Darwin, which has kindly been shown me by Mr. G. H. Darwin) puts it, an incipient species, to which, for convenience, it is desirable to give a distinctive name. That it may be looked upon as a good species by excellent authority, will be seen by Walsh's discussion of the subject (*American Entomologist* II, p. 335) which I here quote:

What candid entomologist, who has worked much upon any particular order, will not allow that there are certain genera where it is often or almost or quite impossible

to distinguish species by the mere comparison of cabinet specimens of the imago? Lœw and Osten Sacken have said this of the genus *Cecidomyia* in Diptera; Osten Sacken of two other Dipterous genera, *Sciara* and *Ceratopogon*; Norton of the genus *Nematus* in Hymenoptera; and Dr. Le Conte lately assured me that, although when he was a young man he thought himself able to discriminate, in the closet, between the different species of *Brachinus* in Coleoptera, he now considered it quite impracticable to do so with any degree of certainty. And yet who doubts the fact of the existence, in North America, of very numerous distinct species of *Cecidomyia*, of *Sciara*, of *Ceratopogon*, of *Nematus*, and of *Brachinus*.

Upon the same principle I strongly incline to believe that the 17-year form of the Periodical Cicada (*C. septemdecim*, Linn.) is a distinct species from the 13-year form (*C. tredecim*, Riley) although it has been impossible for me, on the closest examination of very numerous specimens, to detect any specific difference between these two forms.* It is very true that the 13-year form is confined to the more southerly regions of the United States, while the 17-year form is generally, but not universally, peculiar to the Northern States; whence it has been, with some show of plausibility, inferred that the 13-year form is nothing but the 17-year form accelerated in its metamorphosis by the influence of a hot southern climate. But as these two forms interlock and overlap each other in various localities, and as it frequently happens that particular broods of the two forms come out in the same year, we should certainly expect that, if the two forms belonged to the same species, they would occasionally intercross, whence would arise an intermediate variety having a periodic time of 14, 15 or 16 years. As this does not appear to have taken place, but, on the contrary, there is a pretty sharp dividing line between the habits of the two forms, without any intermediate grades of any consequence, I infer that the internal organization of the two forms must be distinct, although externally, when placed side by side, they are exactly alike. Otherwise, what possible reason could there be for one and the same species to lie underground in the larva state for nearly 17 years in one county, and in the next adjoining county to lie underground in the larva state for scarcely 13 years? I presume that even the most bigoted believer in the old theory of species would allow that, if it can once be proved to his satisfaction that two apparently identical forms are always structurally distinct, whether in their external or in their internal organization, they must necessarily be distinct species.

On the other hand, I firmly believe that many perfectly distinct forms, which at one time passed current, or which even now pass current, as true species, are in reality mere dimorphous forms of one and the same species. We find a good example of this in the dimorphous ♀ *Cynips*, *q. aciculata*, O. S., which has already been treated of at great length. We find another good example of the same thing in *Cicada Cassinii* ♂ ♀, Fisher, which is sufficiently distinct from the Periodical Cicada to have been classified as a distinct species, and yet never occurs except in the same year and in the same locality as this last, and what is more extraordinary still, is found not only along with the 17-year form (*C. septemdecim*), but also along with the 13-year form (*C. tredecim*).

Now, if *Cassinii* were a distinct species, and not, as I believe it to be, a mere dimorphous form of *C. septemdecim* and *C. tredecim*, the chances are more than a million millions to one against its always coinciding with the two other forms, not only as to the particular locality but as to the particular year of its appearance.

I do not know that any one has heretofore attempted to set at rest, by actual proof, the very general skepticism as to this insect remaining so long underground, on the part of those persons who have given little attention to the subject. I have been able to trace the development from year to year of my *tredecim* brood XVIII in the vicinity of Saint Louis by digging up the larvæ each year from 1868 to 1876, and noting the annual growth. They could always be found within from two to five feet of the surface upon the roots of trees, and had by the 8th year attained the first pupa stage, and I have no doubt but that, at this writing, the true pupæ are nearing the surface of the ground to appear in myriads in the perfect state in May and June of this year.

The fungus affecting this Cicada has since been described by Mr. C. H. Peck as *Massospora cicadina* (31st Rep. N. Y. State Mus. Nat. Hist., pp. 44, 1879).

ERIOSOMA PYRI, Fitch (Rep. I, p. 118) — After comparing specimens in Europe with our American insect, I have no doubt of the specific identity of the two, or of the root-inhabiting and twig-inhabiting forms. The insect should be known, therefore, as *Schizoneura lanigera* (Hausm.). See my remarks in *American Entomologist*, II, 359;

*For an excellent statement of the facts bearing upon this curious question, see a paper by Mr. Riley, the State Entomologist of Missouri, in No. 4 of the *American Entomologist*, and a still more complete one in his First Annual Report.

Rep. 3, p 95, and "Notes on Aphididæ of the U. S." (Hayden's Bull. U. S. Geol. & Geogr. Surv. of Terr., Vol. V, p. 3).

ASPIDIOTUS HARRISII Walsh (Rep. I, p. 7) — This belongs to Costa's genus *Diaspis*, and is apparently the species named *ostreæformis* by Curtis (Gardener's Chronicle, 1843, p. 805).

DIPTERA.

TRUPANEA APIVORA Fitch (Rep. I, p. 168; II, 122) — This has been renamed *Pro-machus Fitchii* by Osten Sacken (Cat. of the described Diptera of N. A. 2nd Ed., 1878, p. 234), the species proving different from *Bastardii* Lœw, and Fitch's name being pre-occupied.

BEE-FLY LARVA (Rep. IX, p. 96) — The undetermined larva here illustrated (Fig. 24) has since proved to be that of a *Systæchus*, a genus of Bombyliid flies. For further details and determinations see the Second Report of the Commission (pp. 262-9).

SARCOPHAGA CARNARIA, L. (Rep. IX, p. 95) — The variety *sarracenix* of this species there mentioned is now considered a good species, for reasons stated in Bulletin 3 of the Commission (pp. 39, 40, note).

EXORISTA LEUCANÆ, Kirkpatrick (Rep. II, p. 50 and subsequently) — Referred to the genus *Nemoræa* Desv. by Osten Sacken (Catalogue, etc., 1878, p. 150). The variety *cecropix* of this (Rep. IV, p. 108) is quoted by him as a distinct species under *Exorista*, probably a mistake caused by my employing the wrong figure in the *American Entomologist*, Vol. II, p. 101, where that of *E. flavicauda* is used for *leucanix*.

LYDELLA DORYPHORÆ Riley (Rep. I, p. 111) — Now included in the genus *Exorista*.

ORTHOPTERA.

ÆCANTHUS NIVEUS, De Geer (Rep. I, p. 138, and V, p. 120) — This species is common in all parts of the country, and I have proved, by breeding, that its eggs are those described and figured as such in the 5th Report. I agree with Scudder in considering *fasciatus* De Geer but a dark and rather well marked variety of it. Its chirp is intermittent, resembling a shrill *te-reat te-reat te-reat* with a slight pause between each. The eggs and punctures figured on page 119 of the 5th Report (Fig. 47) as probably those of *Orocharis saltator* are, as I have since proved by breeding and by watching the process of oviposition, those of a large species of *Æcanthus*, hitherto, I believe, very generally confounded with *niveus*, and which is described below as *Æ. latipennis* N. Sp. While *niveus* punctures all kinds of soft stems and pithy twigs, *latipennis* seems to prefer the more slender parts of the Grape-vine. The female, when she has sufficiently proceeded in the act of ovipositing, is so intent that she can very well be watched at night by the aid of a "bull's-eye."

The jaws are first used to slightly tear the outer bark. With the antennæ stretched straight forward and the abdomen bent up so as to bring the ovipositor at right angles with the cane, she then commences drilling, working the abdomen convulsively up and down about twice each second. The eggs, as described in the Report, are laid lengthwise in the pith, but always in two sets, one each side of the hole. The number varies according to the size of the cane, and the distance between the holes is also variable but usually less than in my figure. The hole is usually filled up with a white mucous secretion, though there is very little of it about the eggs. This secretion also doubtless serves to facilitate the drilling. The same female will lay over 200 eggs, and will sometimes puncture the same cane at intervals of $\frac{1}{8}$ inch for $1\frac{1}{2}$ feet or more.

The shrill of *latipennis* is continuous and recalls the trilling of a high-pitched dog-whistle in the distance. The key varies, however, and is sometimes much less high and more musical than at others. The commingled shrill of this species recalls also the distant croaking of frogs in spring. The broad wings are thoroughly elevated during the act or even bent forward, and the vibration is so rapid that there appears

to be no motion. The species, in addition to these differences in stridulation and habits, may be distinguished from *niveus* by the following characters:

ÆCANTHUS LATIPENNIS N. Sp.—White, the elytra of the ♀ sometimes grayish and the posterior femora in one specimen discolored. Antennæ immaculate, with the basal joints and the front of head usually roseate. Tip of ovipositor black. Pronotum as in *niveus*. Hind wings ♀ as long as the elytra or sometimes a trifle longer; of ♂ somewhat shorter than elytra. Elytra of ♀ irregularly reticulate between the parallel oblique veins, especially toward the base. Elytra of ♂ when unfolded $\frac{4}{5}$ as wide as long, the dorsal surface 14^{mm} to 16.5^{mm} long by 7^{mm} to 8^{mm} wide; the rasp 1.5^{mm} long and the teeth distinctly seen with a lens of low power. Ovipositor 6^{mm} long; subgenital plate broadly excavated. Claspers of ♂ with their tips broad, but slightly broader at base than at tip, not deeply separated.

Described from 15 ♂ ♀ specimens from Missouri, 1 ♂ from Alabama, and 1 ♂ from South Texas.

The form of the subgenital plate, the immaculate antennæ with their roseate base, and the larger size serve to distinguish the species as well in the pupa as in the imago state.

Æ. latipennis is a larger insect than *niveus* usually is. The ovipositor measures 6^{mm} in length, whereas in *niveus* it rarely exceeds 5^{mm} and in only one specimen, a sanguineous variety captured July 10, 1874, does it equal 6^{mm}. The male elytra of *niveus* in only one specimen, captured September 19, 1877, reach 13^{mm} in length by 6^{mm} in width on the upper face, and the size is generally much less. In *niveus* the unfolded male elytra are less than $\frac{3}{4}$, and usually only $\frac{1}{2}$, as wide as long, and the rasp is only 1^{mm} long, and the teeth are not so easily seen. The elytra of *niveus* female sometimes show an irregularity in the reticulation between the parallel oblique veins but never so great an irregularity as in *latipennis*, there being fewer cells. In only one specimen of *latipennis*, a male taken on cotton at Columbus, Tex., are there any black marks on the lower surface of the basal joints of the antennæ, representing the lines or dots which are always present in *niveus*. But the two species are most sharply separated by the form of the subgenital plate of the female, which in *niveus* narrows rapidly towards the tip which has a minute angular notch, and by the form of the male claspers, which in *niveus* have their tips very slender and parallel, being deeply parted, and then retreating rapidly from one another on each side.

Besides *niveus* there are recognized from North America three other species of *Æcanthus*, one of which, *californica* Sauss.,* recorded only from California, is described as having the posterior wings abortive.† The other two species, *nigricornis* Walk. from Illinois (description quoted in the *American Entomologist*, Vol. II, p. 207; 1870) and *varicornis* Walk. from Mexico, both described only in the female sex and differing from *niveus* in nothing but the slightly longer hind wings and the slightly greater size of the insect, and in *varicornis* having a slightly longer prothorax, have been retained as distinct species by Saussure. But *niveus*, as may be seen in a series of specimens, varies in these characters indefinitely, just as other species of crickets are admitted to vary; so we may consider Walker's species to be but varieties of *niveus*. They cannot be referred to *latipennis*, for in this species the wings rarely, and then but slightly, exceed the elytra.

One other North American species, *bipunctatus* DeG., has been referred to *Æcanthus*. It belongs, however, to the genus *Xabea* and should be known as *Xabea bipunctata* (DeG.).

As the female of *Xabea*† has not hitherto been described and Saussure did not recognize the genus as distinct from *Æcanthus*, it may be well to give here the characters drawn from both sexes to show how very clearly the two genera differ. The type of the genus is from Sumatra, and Walker, being unacquainted with our species, and Saussure, having only imperfect specimens, both failed to recognize the existence of the genus in North America.

* Études sur les Orthoptères, (in Mission Scientifique au Mexique, etc. Recherches Zoologiques 6^{me} partie.) 3^{me} livraison; p. 462; 1874.

†By "abortive" is evidently meant, from the description following the diagnosis, simply shorter than abdomen. In this respect and in the male (which alone is described) being shorter than *niveus*, *californicus*, which I know only from the description, may most easily be distinguished.

‡ Walker, Cat. Derm. Salt. Brit. Mus., Pt. I, p. 109.

XABEA Walk.—First joint of antennæ armed with a stout, blunt tooth in front. Female elytra irregularly reticulated, the oblique longitudinal veins not being conspicuous; male elytra with the mediastinal vein strongly arcuated; no humeral angle. Wings twice as long as the elytra. Cerci only half as long as the abdomen, sinuous. Outer valves of the ovipositor ending in a single outwardly directed tooth which is preceded on the outside by a longitudinal series of three teeth; the inner valves compressed, ending in three teeth of which the middle one is much the longest. Posterior tibiæ with neither spurs nor serrations and having only 4 apical spurs, 2 within and 2 without; the first joint of posterior tarsi unarmed, the tarsi clearly but 3-jointed, the second joint short as in the other legs; tarsal claws with the inner tooth acute.

OROCHARIS SALTATOR Uhler (Rep. V, p. 119).—The eggs figured and described on page 119 as probably those of this insect are, as above stated, those of *Æcanthus latipennis*. I have, however, frequently obtained the eggs of the Orocharis since. In December, 1877, I watched a female ovipositing in the end of a dead and rather soft twig of the Soft-maple at Kirkwood, Mo. The twig had been pruned and the bark was somewhat gnawed by the cricket and the eggs thrust in irregularly from the end and from the sides. Both wood and pith were crammed with eggs, but all longitudinally inserted. The favorite nidus of the species is, however, the soft and somewhat corky, rough bark of the trunk and older branches of the American elm, the eggs being thrust in singly or in small batches, either longitudinally with, or very slightly obliquing from, the axis of trunk or branch. The female is very intent in the act, working her abdomen deliberately from side to side during the perforation. The ovipositor is held more obliquely than in *Æcanthus*.

The egg is amber-colored and very slender and elongate, the tip rather pointed and very faintly opaque with the surface but slightly granulate. It has scarcely any curve and varies from 3.5^{mm} to 4^{mm} in length and from 0.4^{mm} to 0.5^{mm} in diameter at middle.

The stridulation of this cricket is a rather soft and musical piping of not quite half a second's duration, with from 4 to 6 trills, but so rapid that they are lost in the distance. The key is very high, but varies in different individuals and according to moisture and temperature. It most resembles the vibrating touch of the finger on the rim of an ordinary tumbler when three-fourths filled with water—repeated at intervals of from 2 to 4 per second, and it may be very well likened to the piping of a young chick and of some tree frogs. As the species is very common in the Southwest its chirp is everywhere heard and is so distinctive that when once studied it is never lost amid the louder racket of the katydids and other night choristers. It is frequently heard during the day time in cloudy or damp weather, and I have heard it at Saint Louis the first days of November after a slight frost. The elytra in stridulating are raised less than in *Æcanthus* and are depressed at intervals.

The courting of the sexes is amusing. They face each other and play with their antennæ for the best part of an hour or more than an hour. The female is, otherwise, pretty quiet, but the male continually mouths the twig or the bark upon which the courting is being done, and plays his palpi at a great rate, very stealthily approaching nearer to his mate meanwhile. At last the antennal fencing ceases and those of the female bend back and then the male approaches until their heads touch. He then deliberately turns round, elevates the elytra and slips his abdomen under the female, who virtually mounts and assists him, his elytra overshadowing her head.

The eggs of this insect, as also those of *Æcanthus latipennis*, are devoured by a parasitic larva of similar form and size, and which I have not yet reared to the perfect state.

ORCHELIMUM GLABERIMUM, Burm. (Rep. V, p. 123).—The egg-punctures illustrated at Fig. 56 are, as there correctly supposed, those of this species, as I have since proved by watching the act of oviposition and by rearing from the eggs. The insect is very fond of using the tops of corn-stalks for the same purpose.

NEUROPTERA.

CORYDALUS CORNUTUS, L. (Rep. V, p. 141; IX, p. 125)—For additional facts relating to the early larval stages, see my notes on the "Larval Characteristics of Corydalus and Chauliodes and on the development of Corydalus cornutus (Proc. Am. Ass. Adv. Sc., 1878).

MITES.

TROMBIDIUM SERICEUM Say (Rep. VII, p. 175 and subsequently)—For the natural history of this species and the specific identity with it of the larval form known as *Astoma gryllaria* LeBaron, and for further facts respecting the other mites mentioned in the Report, see my remarks in the Transactions of the Academy of Science of Saint Louis, (Vol. III, p. cclxvii, October, 1877) in the *American Naturalist* for March, 1878, and in the First Report of the Commission (p. 306 ff.).

DESCRIPTIONS OF NEW SPECIES AND VARIETIES.

Some systematists have questioned whether descriptions of species in Agricultural Reports should be recognized. While my own views on this subject are pretty freely expressed on page 56 of my Third Missouri Report and elsewhere, the publication of this Bulletin affords a good opportunity to bring the descriptions that are scattered through the nine volumes together, with such notes on synonymy as present knowledge suggests, and such corrections as are given in the Errata. In the earlier reports the measurements were expressed in inches and hundredths of an inch, while in the later volumes the metric system was adopted as most convenient and accurate, and the measurements which follow have all been reduced to this standard. All changes of this character or other changes from the original are included in brackets, while the additional notes are in Long Primer type.

HYMENOPTERA.

PORIZON CONOTRACHELI, N. SP.—*Head* pitchy-black, opaque, the ocelli triangularly placed and close together; eyes oval, polished, and black; face covered with a silvery-white pubescence; labrum rufous, with yellowish hairs; mandibles and palpi, pale yellowish-brown; antennæ inserted in depressions between the eyes, reaching to metathorax when turned back, filiform, 24-jointed; black with basal joints 6-1 becoming more and more rufous, the bulbus always distinctly rufous; bulbus rather longer and twice as thick as joint 3; joint 2 about one-third as long. *Thorax* pitchy-black, opaque, the sides slightly pubescent with whitish hairs, the mesothorax rounded and bulging anteriorly, the scutellum slightly excavated and sharply defined by a carina each side; metathorax with the elevated lines well defined and running parallel and close together from scutellum to about one-fourth their length, then suddenly diverging and each forking about the middle. *Abdomen* glabrous, polished, very slender at base, gradually broader and much compressed from the sides at the apex which is truncated; peduncle uniform in diameter and as long as joints 2 and 3 together; joints 2-5 subequal in length; color rufous with the peduncle wholly, dorsum of joint 2, a lateral shade on joint 3, and more or less of the two apical joints superiorly, especially at their anterior edges, black; venter more yellowish: ovipositor about as long as abdomen, porrect when in use, curved upwards when at rest, rufous, with the sheaths longer and black. *Legs*, including trochanters and coxæ uniformly pale yellowish-brown with the tips of tarsi dusky. *Wings*, subhyaline and iridescent, with veins and stigma dark brown, the stigma quite large, and the two discoidal cells subequal and, as usual in this genus, joining end to end, but with the upper veins which separate them from the radial cell, slightly elbowed instead of being straight, thus giving the radial cell a quadrangular rather than a triangular appearance. ♂ differs from ♀ only in his somewhat smaller size and unarmed abdomen. Expanse ♀ 0.32 inch [= 8^{mm}], length of body, exclusive of ovipositor, 0.22 [= 5.5^{mm}]; expanse ♂ 0.28 [= 7^{mm}], length 0.18 [= 4.5^{mm}].

Described from 3 ♀ ♀, 1 ♂ bred May 26th-28th, 1870, from cocoons received from Dr.

I. P. Trimble, of New Jersey, and 1 ♀ subsequently received from the same gentleman—all obtained from larvæ of *Conotrachelus nenuphar*.

As I am informed by Mr. E. T. Cresson, of Philadelphia, who pays especial attention to the classification of the *Ichneumonidæ*, it might more properly be referred to Holmgren's genus *Thersilochus*, which differs from *Porizon* in the greater distance between the antennæ at base, and in the venation of the wing.—[Third Rept., p. 28, Fig. 9.

LIMNERIA LOPHYRI, N. SP.—♀, length 0.30—0.35 inch [7.5—8.7^{mm}]. Head and thorax black with silvery white pile. Antennæ piceous, more than half as long as body; but slightly paler toward tip; bulbus either yellowish or rufous. Ocelli either rufous or black. Mandibles, palpi, front and middle coxæ trochanters and tibiæ, pale yellow. Tegulæ almost white. Abdomen, with faint pile, rufous, the petiole and sides of next joint usually blackish. Hind legs rufous, the base of tibiæ and of tarsi paler.

♂ somewhat smaller, and with more black on the abdomen.

Four ♂'s, 12 ♀'s bred from larvæ of *Lophyrus Abbotii*.—[Ninth Rept., p. 32.

HEMITELES (?) CRESSONII, [N. Sp.]—♂—Length 0.25 [6^{mm}]. Black, opaque, head transversely-subquadrate; face clothed with pale glittering pubescence; spot on mandibles, palpi, scape of antennæ in front and the tegulæ, white; eyes large, ovate; antennæ longer than head and thorax, slender, black; thorax closely and minutely punctured; mesothorax with a deeply impressed line on each side anteriorly; scutellum convex, closely punctured, deeply excavated at base; metathorax coarsely sculptured, truncate and excavated behind, the elevated lines sharply defined, forming an irregularly shaped central area, and a triangular one on each side of it, the outer posterior angle of which is prominent and subacute; wings hyaline, iridescent, nervures blackish, stigma large, areolet incomplete, the outer nervure wanting; legs pale honey-yellow, coxæ paler, tips of posterior femora, and their tibiæ and tarsi entirely blackish; abdomen elongate ovate, flattened, petiolated, the first segment flat, gradually dilated posteriorly, somewhat shining, and indistinctly longitudinally aciculate; the two following segments opaque, indistinctly sculptured; remaining segments smooth and shining.—[First Rept., p. 177. Figured at Pl. II, Fig. 7.

HEMITELES (?) THYRIDOPTERIGIS, N. SP.—♀ Length 0.36 [inch = 9^{mm}]; expanse 0.50 [inch = 12.5^{mm}]. Ferruginous, opaque. Head transverse, rather broader than thorax, the front much depressed; face prominent centrally beneath antennæ, closely punctured, thinly clothed with pale pubescence; clypeus and cheeks shining; tips of mandibles black; antennæ, long, slender, filiform, ferruginous, blackish at tips; thorax rugose; scutellum prominent, with sharp lateral margins; metathorax prominent, quadrate, abrupt laterally and posteriorly, finely reticulated and pubescent, the upper posterior angles produced on each side into a long, divergent, flattened, subacute spine; disk with two longitudinal carinæ, from which diverges a central transverse carina; tegulæ piceous; wings hyaline, subiridescent; a narrow, dark fuliginous band crosses the anterior pair a little before the middle, and a broad band of same color between middle and apex, this band having a median transverse hyaline streak; areolet wanting, second recurrent nervure straight, slightly oblique; apex of posterior wing fuscous; legs long and slender, ferruginous, more or less varied with fuscous; posterior coxæ, tips of their femora, and their tibiæ and tarsi, fuscous; base of four posterior tibiæ more or less whitish, forming a rather broad annulus on posterior pair; abdomen petiolated, subconvex, densely and finely sculptured, blackish, basal segment tinged with reddish, the second and third segments distinctly margined at tip with whitish; apical segments smooth and shining, thinly pubescent; ovipositor half as long as abdomen, sheaths blackish.

♂.—Not at all like the ♀. Length 0.33 [inch = 8^{mm}], expanse 0.44 [inch = 11^{mm}]. Long, slender, black, polished, without distinct punctures, thinly clothed with white pubescence; palpi white; antennæ long, slender; scape reddish; mesothorax gibbous, with two deeply impressed longitudinal lines; metathorax with well-defined elevated

lines, forming several irregular areas; sides rugulose, apex without spines or tubercles; tegulae white; wings whitish-hyaline, subiridescent, the nervures and stigma white, subhyaline, neuration as in ♀; legs long, slender, pale honey-yellow; coxae, posterior trochanters, apex of their femora, and their tibiae and tarsi, blackish; base of posterior tibiae with a white annulus; abdomen long, slender, flattened, petiolated, smooth and polished, the apical margin of second segment being narrowly whitish.

Described from four ♀ and one ♂ specimens bred from the same [*Thyridopteryx*] cocoon.—[First Rept. p. 150. Figured at Pl. II, Figs. 11, 12.]

The species is quite common in Washington, D. C., and is often attacked by a secondary Chalcid parasite.

MICROGASTER LIMENITIDIS, N. SP.—♂ ♀. Length 0.09 inch [= 2¼^{mm}]. Color pitchy-black. Antennae black, about as long as body; palpi whitish. *Thorax* minutely punctured. *Abdomen* with the two or three basal joints emarginate and rugose, the terminal joints smooth and polished. *Legs* dusky; front and middle femora yellowish, hind femora black; front and middle tibiae yellowish, hind tibiae with terminal half dusky, but the spur pale; front and middle tarsi yellowish tipped with dusky, hind tarsi dusky above, paler below. *Wings* hyaline, iridescent, the nervures and stigma black or dark-brown, the radial nervule, the cubital nervules and the exterior nervule of the discoidal cell, sub-obsolete.

Described from 6 ♀, 1 ♂, bred from larvae of *Limenitis disippus*.—[Third Rept., pp. 158, 159.]

The specimens referred to in connection with this description as bred from *Gelechia gallasolidaginis* prove to belong to a distinct species. Both species belong to the genus *Apanteles* Först. as at present accepted. See my "Notes on N. A. Microgasters" (Trans. Ac. Sc. St. Louis, IV, *Author's separata*, p. 13.)

MICROGASTER GELECHIAE.—Length 0.20 [= 5^{mm}] ♂ ♀.—Black, clothed with a short, thin, glittering, whitish pubescence, most dense on the face, which latter is closely punctured; occiput and cheeks shining; mandibles rufopiceous; palpi whitish; eyes pubescent; antennae as long as the body in ♂, shorter in ♀, 18-jointed; thorax shining, feebly punctured, mesothorax closely and more strongly punctured, with a deeply impressed longitudinal line on each side over base of wings; scutellum smooth and polished, the lateral groove broad, deep, arched and crenulated; metathorax opaque, densely rugose, with a sharp, central, longitudinal carina, and a smooth, flat, transverse carina at base; tegulae testaceous, wings hyaline, iridescent, apex smoky, nervures blackish, areolet complete, subtriangular, radial nervure indistinct; legs pale honey-yellow, coxae blackish, pale at tips, middle pair in ♀ concolorous with legs; abdomen with the two basal segments densely rugose and opaque, the remainder smooth and shining; venter more or less varied with pale testaceous.—[First Rept., p. 178.]

This is a true *Microgaster*.

PERILITUS INDAGATOR, N. Sp.—Imago—♀, *Head* almost glabrous, transverse, deep honey-yellow, the trophi pale, except the tips of jaws, which are dusky; ocelli touching each other, black; eyes black, very large, occupying nearly the whole side of face, and with a few very short hairs; antennae with about 24 joints, pale fuscous; reaching, when turned back, to about the middle of abdomen. *Thorax* honey-yellow beneath and very slightly pubescent; very finely punctured and slightly pubescent above; prothorax honey-yellow and prominently convex; mesothorax with lateral and posterior sutures black; metathorax black. *Abdomen* with the pedicel black and slightly punctured; depressed, narrow at base, widening behind, slightly pubescent above; the other joints glabrous, polished, deep honey-yellow, the second joint largest and as long as all the subsequent ones together; ovipositor extending about the length of the abdomen beyond its tip, rufous with the sheaths black. *Legs*

pale honey-yellow, the tarsi, especially at tips, slightly dusky, the hind femora and tibiae a little dusky towards tips, and a narrow rufous ring at base of former. *Wings* hyaline, iridescent; veins brown; stigma honey-yellow, with an opaque brown cloud; two cubital cells, the outer small, sub-quadrate; the radial large; one discoidal, long and narrow. Length, exclusive of ovipositor, 0.18 inch [= 4.5^{mm}].

Described from 1 ♀ bred from *Acrobasis juglandis*, LeB.—[Fourth Rept., p. 43.

SPATHIUS TRIFASCIATUS, N. Sp.—♀. Average length, 0.18 inch [= 4.5^{mm}]. Color, light-brown. *Head* pubescent, palpi long and pale; eyes black; ocelli black, contiguous; antennae smooth, pale, and reaching to second abdominal joint. *Thorax* with sutures dark-brown; legs more or less dusky, the tarsi (except at tip) an annulus at base of tibiae, and the trochanters, pale; wings fuliginous, with a white fascia at base, at tip and across outer middle of front wing, including the inner half of stigma, the outer half of which is dark-brown; middle fascia most clearly defined. *Abdomen* slightly pubescent at sides and tip; first joint pale, petiolate, and with short and longitudinal aciculations above; second joint pale above, the others more or less brown; ovipositor pale, dusky at tip, and long as abdomen.

One bred specimen.

♂—Differs in being much darker colored, the head, thorax and femora being brown, and the metathorax and base of first abdominal joint black.

One bred specimen.—[Fifth Rept., p. 106.

BRACON CHARUS, N. sp.—♀ Length of body 0.35 inch [= 8.7^{mm}]; of ovipositor 0.40 inch [= 10^{mm}]; expanse of wing 0.65 inch [= 16^{mm}]. Colors black and deep rufous. Head, thorax, legs and antennae polished black, the legs and sides of head and thorax with a fine grayish pubescence; trophi also black. Abdomen uniformly deep rufous. Terebra of ovipositor pale yellow, the sheaths black and very faintly pubescent. Wings deep fuliginous with a faint zig-zag, clear line across the middle from the stigma.

Described from 7 ♀'s, all bred from *Chrysobothris femorata*.—[Seventh Rept., p. 75. Fig. 13.

BRACON SCOLYTIVORUS, Cress.—♀—Black, shining, metathorax and base of abdomen pubescent; face, anterior orbits, lower half of cheeks, clypeus, mandibles, except tips, palpi, tegulae, legs, including coxae, and abdomen, honey-yellow, the latter darker; posterior coxae sometimes dusky; antennae at base beneath, dull testaceous; wings fuliginous, apical half paler, iridescent; abdomen shining, first segment whitish laterally, the base and disc sometimes dusky; base of second segment with a large subtriangular flattened space inclosed by a deep groove, the posterior side of which is generally blackish; ovipositor longer than abdomen; sheaths black; length, .15—.17 inch [= 3³/₄—4¹/₄^{mm}].

♂—More pubescent; posterior coxae blackish, also the femora above, especially the posterior pair; posterior tibiae dusky; abdomen black, polished; apex of first, basal half of second, and sides of apical segments more or less honey-yellow; sides of basal segment whitish; wings paler; abdomen narrower and rather more convex; length, .16 inch [= 4^{mm}].

Three ♂, three ♀ specimens.—[Mr. E. T. Cresson, in Fifth Rept., p. 106.

SIGALPHUS CURCULIONIS, Fitch—*Imago*.—*Head* black, sub-polished, and sparsely covered on the face with short whitish hairs; ocelli touching each other; labrum and jaws brown; palpi pale yellow; antennae (Fig. 7, c) 27-jointed, filiform, reaching, when turned back, to middle joint of abdomen and beyond, the bulbous and small second joint rufous and glabrous, the rest black or dark brown, though 3-10 in many specimens are more or less tinged with rufous; 3-14 very gradually diminishing in size; 14-27 subequal. *Thorax* black, polished, the metathorax distinctly and broadly punctate, and the rest more or less distinctly punctate or rugose, with the sides sparsely pubescent. *Abdomen* pitchy-black, flattened, the dorsum convex, the venter concave, and the sides narrow-edged and slightly carinated; the three joints distinctly separated and of about equal length; the first joint having two dorsal longitudinal carinae down the

middle; all densely marked with very fine longitudinally impressed lines, and sparsely pubescent; (Dr. Fitch in his description published in the *Country Gentleman*, under date of September, 1859, states that these lines leave "a smooth stripe along the middle of its second segment and a large smooth space on the base of the third;" which is true of a few specimens, but not of the majority, in which the impressed lines generally cover the whole abdomen.) Ovipositor longer than abdomen, but when stretched in a line with it, projecting backwards about the same length beyond; rufous, with the sheaths black. Legs pale rufous, with the upper part of hind tibiae and tarsi, and sometimes the hind femora, dusky. Wings subhyaline and iridescent, the veins pale rufous, and the stigma black. Length ♀, 0.15–0.16 inch [=3.7–4^{mm}], expanse 0.30 [=7.5^{mm}]; ♂ differs only in his somewhat smaller size and in lacking the ovipositor. In many specimens the mesothorax and the eyes are more or less distinctly rufous.

Described from 50 ♀♀, 10 ♂♂, bred June 23d–July 29th, 1870, from larvæ of *Conotrachelus nenuphar*, and 2 ♀♀ obtained from Dr. Fitch.

Larva (Fig. 8, *a*)—White, with translucent yellowish mottlings.

Pupa (Fig. 8, *c* ♀)—0.17 inch [=4¹/₄^{mm}] long; whitish, the members all distinct, the antennæ touching hind tarsi, the ovipositor curved round behind, reaching and touching with its tip the third abdominal joint, which afterwards forms the apical joint of imago; five ventral joints, which in the imago become much absorbed and hidden, being strongly developed.

Cocoon (Fig. *c*, *b*)—Composed of one layer of closely woven yellowish silk.

VARIETY RUFUS—Head, thorax, and most of the first abdominal joint entirely rufous, with the middle and hind tibiae dusky, and the ovipositor three times as long as abdomen and projecting more than twice the length of the same beyond its tip.

Described from three ♀♀ bred promiscuously with the others. This variety is slightly larger and differs so remarkably from the normal form that, were it not for the absolute correspondence in all the sculpturing of the thorax and body, and in the venation of the wings, it might be considered distinct. The greater length of the ovipositor is very characteristic, and accompanies the other variation in all three of the specimens.—[Third Rept., p. 27. Fig. 7.]

EURYTOMA BOLTERI, N. Sp.—♀ Length 0.18 inch [=4.5^{mm}]. Antennæ black, not much longer than the face, perceptibly thicker towards the end, and apparently 10-jointed, though the three terminal joints are almost always confluent. Dimensions and appearance of joints, represented in the annexed Figure 97, *a*. Head and thorax rough-punctured and finely bearded with short, stiff gray hairs. Abdomen about as long as thorax, scarcely so broad, viewed from above, but wider viewed laterally; highly polished, smooth and black, the three terminal segments with minute stiff gray hairs along the sutures; visibly divided into seven segments, the four anterior ones of about equal length, the two following shorter, and the terminal one produced into a point. Legs fulvous with the *coxae*, [trochanters], thighs and more or less of the shanks blackish-brown. Wings perfectly transparent, glossy, colorless, and with the nerves very faint.

♂ Measures but 0.14 inch [=3.5^{mm}], and differs in the antennæ, being twice as long as the face, in their narrowing towards the tip and in being furnished with whorls of long hairs. The number of joints are not readily made out, and I have consequently presented at Figure 97, *b*, a magnified figure. His body is but half as wide and half as long as the thorax viewed from above, and not quite as broad as the thorax, viewed laterally; it also lacks the produced point of the ♀. His wings are also cut off more squarely and more distinctly nerved.—[First Rept., p. 187. Pl. II, Fig. 9.]

For further descriptive details see Walsh's posthumous paper on the *Eurytomides* (*Am. Ent. II*, p. 298–9), where the insect is looked upon as a variety of *Eurytoma diastrophii*.

[TRICHOGRAMMA MINUTA, N. Sp.] * * * It comes nearest the genus *Trichogramma*, Westw., and may be provisionally called *Trichogramma* (?) *minuta*. It differs

from that genus and from all other Chalcididan genera with which I am acquainted, in the antennæ being but 5-jointed (scape, plus 4 joints), the scape stout and as long, or longer, than joints 2, 3, and 4 together; joints 3 and 4 small and together as long as joint 2; 5 very stout, fusiform, and as long as 2, 3, and 4 together. The legs have the trochanters stout and long, the tibiæ not quite so long nor so stout as the femora, and with a long tooth; the tarsi are 3-jointed, with the joints of equal length and with the claws and pulvilli sub-obsolete. The abdomen is apparently 6-jointed, the basal joint wide, the 2nd narrower, 2-5 increasing in width till 5 is as wide as 1. The ovipositor of ♀ extends a little beyond the apex, and starts from the anterior edge of the 5th joint.—[Third Rept., p. 158. Fig. 72.]

The species was provisionally referred to *Trichogramma*, and I subsequently proposed for it the generic name *Pentarthron* (Record of Am. Ent. 1871, p. 8). *Pentarthrum* has, however, been used by Wollaston in beetles, and until allied genera are better characterized than at present, the old generic name may be retained.

COLEOPTERA.

BRUCHUS FABÆ N. Sp. (Fig. 19.)—General color tawny-gray with more or less dull yellowish. *Body* black tinged with brown and with dull yellowish pubescence, the pygidium and sides of abdomen almost always brownish. *Head* dull yellowish-gray with the jaws dark brown and palpi black; antennæ not deeply serrate in ♀, more so in ♂; dark brown or black with usually 5, sometimes only 4, sometimes 4 and part of 5 basal joints, and with the terminal joint, more or less distinctly rufous, or testaceous, the color being so slight in some specimens as scarcely to contrast at all with the darker joints. *Thorax* narrowed before, immaculate, but with the pubescence almost always exhibiting a single pale medio-dorsal line, sometimes three dorsal lines, more rarely a transverse line in addition, and still more rarely (two specimens) forming a large dark, almost black patch each side, leaving a median stripe and the extreme borders pale and thus approaching closely to *erythrocerus* Dej.; base with the edges almost angulated; central lobe almost truncate and with a short longitudinal deeply impressed median line; no lateral notch; scutel concolorous and quadrate with the hind legs more or less notched. *Elytra* with the interstitial lines having a slight appearance of alternating transversely with dull yellowish and dusky; so slight however that in most of the specimens it can hardly be traced: the dark shadings form a spot on each shoulder and three transverse bands tolerably distinct in some, almost obsolete in others, the intermediate row being the most persistent and conspicuous: between these dark transverse rows the interstices are alternately more or less pale, especially on the middle of the 3rd interstitial lines. *Legs* covered with grayish pubescence, and with the tibiæ and tarsi, especially of first and second pair, reddish-brown; the hind thighs usually somewhat darker, becoming black below and inside, and with a tolerably long black spine followed by two very minute ones. Length 0.09—0.14 inch [= 2¼ — 3.5^{mm}]. Described from 40 specimens all bred from different kinds of beans. Hundreds of others examined.

This insect has been for several years ticketed in some of the Eastern collections by the name of *B. fabæ*, or else, what is worse, the corruption of it, *fabi*. The former name has been disseminated by my friend F. G. Sanborn of Boston, Massachusetts, who says that he received the weevil thus named, together with beans attacked by it, in the year 1862 from Rhode Island. The name was credited to Fabricius, but I can find no notice in any of the works I possess of any European *Bruchus fabæ*, and several of my Eastern correspondents who have access to large libraries have been unable to find any description or allusion to a species by that name. Dr. LeConte has given it the MS name of *varicornis* but as his description will not appear perhaps for years to come and as no comprehensive description has yet been published, I have deemed it advis-

able to dispel in a measure the confusion that surrounds the nomenclature of the species. There is need of a description of so injurious an insect, and as *fabæ* is not pre-occupied I adopt the name because it is entirely appropriate and because it is more easily rendered into terse popular language than *varicornis*.

It resembles most closely of any other species which I have seen, the *B. erythrocerus*, Dej., which, however, is smaller, and differs in having a narrower thorax which has light sides and a dark, broad dorsal stripe divided down the middle by a pale narrow line: *erythrocerus* is further distinguished by the antennæ being entirely testaceous and the hind thighs more swollen.

From *obsoletus* Say, *fabæ* differs materially: *obsoletus* is a smaller species, dark gray, with the antennæ all dark, the pygidium not rufous, the thorax with a perceptibly darker dorsal shade so that the sides appear more cinereous, a white scutel, and each interstitial line of the elytra with a slight appearance of alternating whitish and dusky along its whole length; for though there is nothing in Say's language to indicate whether it is the interstitial lines that alternate transversely, whitish and dusky, or each line that so alternates longitudinally, I find from an examination of a specimen in the Walsh collection, that the latter is the case, and so much so that the insect almost appears speckled. The two species differ both in size and color, though, as Say's description is short and imperfect it is not surprising that *fabæ* should have been referred to it.

From the European bean-feeding *Br. flavimanus* (which is apparently either a clerical error for, or a synonym of *Br. rufimanus*, Schœnh.) as described by Curtis, it differs notably; as it does likewise from their *Br. serratus*, Ill., which also attacks beans.

Dr. LeConte, according to Mr. Rathvon, was inclined to consider this insect the *obsoletus* of Say, from the fact that in specimens which the latter gentleman sent him, the antennæ were not varied as in his MS. *varicornis*, but uniformly black. A few specimens which Mr. Rathvon sent me nearly two years ago, taken from the same lot as were those which he forwarded to Dr. LeConte, were singularly enough, all decapitated but two; and these two showed the varied antennæ. These specimens had all been kept in alcohol, and I am greatly inclined to believe that the uniformly dark appearance of the antennæ that was noticed by LeConte was the effect of the alcohol on those which naturally had the rufous joints but faintly indicated. At all events, though Mr. Rathvon tells me that he found a small proportion of beetles with dark antennæ, after examining, at my suggestion, over two hundred specimens that had thus been kept in alcohol; yet from over one hundred specimens which he had the kindness to send me, I only find (after thoroughly drying them) three with the terminal joint really as dark as the subterminal, and not a single one in which the rufous basal joints cannot be more or less distinctly traced.—[Third Rept., p. 55-56. Fig. 19.

Since the above was written, Dr. Horn has given us a revision of the *Bruchidæ* of the United States (Trans. Am. Ent. Soc., Vol. IV, 1873), in which he makes *fabæ* a synonym of *obsoletus* Say, expresses regret that another synonym must be added and states that the *obsoletus* which I referred to is the *transversus* Say (= *hibisci* Oliv.). This criticism is not deserved, and while the decision of one who has done such excellent work in Coleoptera as Dr. Horn has will be generally accepted as final, yet no one can compare his redescription of *obsoletus* with Say's description and not feel that the two apply to different insects. *Fabæ* is usually one-third larger, tawny-gray above with vari-colored antennæ, concolorous scutel, emarginate behind, and rufous legs and abdomen; *obsoletus*, on the contrary, according to Say, is blackish-cinereous, the thorax cinereous each side, with a whitish scutel and with the abdomen and legs not differing in color from the rest of the body. *Fabæ* breeds in beans; *obsoletus* in the seeds

of *Astragalus*. Indeed one would be far more justified in considering *B. alboscuteclatus* Horn a synonym of *obsoletus* Say than in considering *fabæ* a synonym of it, and when the *Bruchus* from *Astragalus* in the Eastern States is bred, I fully expect Dr. Horn to change his mind. Nor is the assumption justifiable that the *obsoletus* referred to by me, and destroyed in the Walsh collection, is *hibisci* Oliv. It was far more like *alboscuteclatus* as far as I remember, and there is not a character about this species which does not accord with Say's description of *obsoletus* except that the scutel is described as *rounded*, while that of *obsoletus* is described by Say as quadrate. I am of opinion that too much stress has been laid on this difference by Dr. Horn, as, when the pubescence is separated behind, the scutel appears quadrate, whereas in *fabæ* it appears bifid. The scutel of *alboscuteclatus* when denuded is quadrate, but it is doubtless the clothed appearance which Say described. Say, as appears from his text, had abundant material, and it is assuming too much to suppose that he could overlook the striking differences in size and coloration of *fabæ*, as above indicated.

The specific name *fabæ* was used by Brullé for *Bruchus pisorum* Linn.

MADARUS VITIS, N. Sp.—Length, exclusive of rostrum 0.10 [inch=2.5mm]. Color uniformly rufous, without maculations, the eyes alone being darker. Highly polished; rostrum arcuated, stout and about as long as thorax; thorax and body with extremely minute and distant punctures, anterior margin of thorax abruptly narrowed, especially laterally, into a collar; elytra slightly undulate, with 4 distinct elevations, one on the extreme outer margin close to the thorax, and one on the middle of each, near the extremity.—[First Rept., p. 132. Fig. 74.]

For further details as to the synonymy of this insect, see *American Entomologist* I, p. 105. Dr. LeConte's description of *Baridius sesostris* was published about three months earlier than my own and he subsequently (Proc. Amer. Phil. Soc., Vol. XV, 1876, p. 299) erected the genus *Ampelogypter* for this and two other species, so that *Madarus vitis*=*Ampelogypter sesostris* Lec.

ANALCIS FRAGARIÆ, N. Sp.—*Imago*, (Fig. 14, b, c)—Color deep chestnut-brown, sub-polished, the elytra somewhat lighter. Head and rostrum dark, finely and densely punctate and with short fulvous hairs, longest at tip of rostrum; antennæ rather lighter towards base, 10-jointed, the scape much thickened at apex, join 2 longest and robust, 3 moderately long, 4-7 short, 8-10 connate and forming a stout club. Thorax dark, cylindrical, slightly swollen across the middle and uniformly covered with large thimble-like punctures, and with a few short coarse fulvous hairs, unusually arranged in three more or less distinct longitudinal lines; pectoral groove ending between front legs. Abdomen with small remote punctures and hairs which are denser towards apex. Legs of equal stoutness, and with shallow dilated punctures and uniform very short hairs. Elytra more yellowish-brown, dilated at the lower sides anteriorly, and with about 9 deeply-punctured striæ, the striæ themselves sometimes obsolete; more or less covered with coarse and short pale yellow hairs which form by their greater density, three more or less conspicuous transverse bands, the first of which is at base; between the second and third band, in the middle of the elytron, is a smooth dark-brown or black spot, with a less distinct spot of the same color below the third, and a still less distinct one above the second band. Length 0.16 inch [=4mm].

Described from four specimens bred from strawberry-boring larvæ. The black spots

on the elytra are quite distinct and conspicuous on two specimens, less so on one, and entirely obsolete on the other.

Larva, (Fig. 14 *a*)—White with back arched Lamellicorn-fashion. Head gamboges yellow, glabrous, with some faint transverse striations above mouth; mandibles rufous-tipped with black; labrum emarginate, and with palpi, pale. A faint narrow dorsal vascular line. Legs replaced by fleshy tubercles. Length 0.20 inch [=5^{mm}] when stretched out.—[Third Rept., p. 44. Fig. 14.

Say's generic name *Tyloderma* having priority over Schönherr's *Anal-cis*, the name of this insect becomes *Tyloderma fragariæ*.

LEPIDOPTERA.

ÆGERIA RUBI, N. Sp.—*Imago*.—Expanse, ♂, 1.00 [inch=25^{mm}]; ♀, 1.25 inch [=31^{mm}]. Front wings transparent, with a broad costal border extending half the width of wing at base, a narrow discal spot, and more or less of the tip dull-ferruginous; the inner border, the inner longitudinal vein, the intermediate space toward posterior angle, and sometimes its whole length, of the same color; veins brownish within and black without the discal spot. Hind wings perfectly transparent, or rarely with a few sparse ferruginous scales; the transverse discal vein pale, the others pale at base, but black toward extremities; costa narrowly golden-yellow, becoming darker toward apex. Fringes dark-brown, those of hind wings appearing darkest by virtue of a dark wing border. Under surface somewhat paler. Abdomen stout, with a very slight anal tuft in ♀; a stouter one in ♂. Antennæ blue-black, not enlarging toward tip, quite pectinate in ♂. Palpi, a narrow ring around neck, the sides of the collar, a broad band curving across tegulæ and around the base of wings, a faint line across middle of thorax, two faint longitudinal lines between it and collar, legs, except outer base (sometimes whole length) of femora and tibiæ, hind third of abdominal joints, and a dorsal and lateral series of abdominal tufts or patches (the dorsal ones, especially on 3d and 7th joints, most persistent and conspicuous)—all golden-yellow: the rest of body black. The orbits are of a somewhat paler-yellow, and the face either gray or bluish.

♂ differs from ♀ in the darker color of primaries, the narrower fringe of secondaries, the narrower ferruginous spot at apex of primaries, the more tufted abdomen, the broader and darker anal tuft, and the pectinate antennæ.

Described from 6 ♂'s, 6 ♀'s, bred from *Rubus*. Approaches nearest to *Trochilium marginatum* Harr., and *T. tibiale* Harr.,* from which it differs in the thoracic marks and the abdominal tufts.

Larva—Length 0.90-1.10 inch [=22.5-27.5^{mm}]; diameter 0.18 [=4.5^{mm}]. Color pale-yellow. Head dark-brown, with a few whitish hairs; mandibles black, the other trophi paler. Cervical shield horny, pale-brown. Each joint with 8 pale, shiny piliferous spots, transversely arranged on 2, 3 and 12; the dorsal 4 quadrangularly arranged and the lateral 2 interrupted by stigmata on all the others. Thoracic legs slightly tinged with brown; prolegs, with the hooklets dark. Several specimens examined.—[Sixth Rept., p. 113. Fig. 30.

ACRONYCTA POPULI, N. Sp.—*Larva*—Length 1.50 [inch, =37^{mm}]. Color yellowish-green, covered with long soft bright yellow hairs which spring immediately from the body, part on the back, and curl round on each side. On top of joints 4, 6, 7, 8 and 11, a long straight double tuft of black hairs, those on 7 and 8 the smallest. Head polished black with a few white bristles. Joint 1 with a black spot above, divided longitudinally by a pale yellow line, giving it the appearance of a pair of triangles. Joint 2 with two less distinct black spots. Thoracic legs black; prolegs black with brownish extremities. Venter greenish-brown. Described from many specimens. When young of a much lighter color, or almost white, with the black tufts short but

* Silliman's Am. Journ. of Sc., XXXVI, p. 309 (1839).

more conspicuous, with a distinct black dorsal line, two lateral purplish-brown bands, and with hairs white, sparse and straight.

Individuals vary much: some have a black dorsal line, some have but three distinct black tufts; some have a sixth tuft of black hairs on joint 9, and others have a few black hairs on all but the thoracic joints. Just before spinning up, many of the hairs are frequently lost, and the body acquires a dull livid hue.

Moth.—♀, front wings, white, finely powdered with dark atoms which give them a very pale gray appearance; marked with black spots as follows: a complete series of small spots on posterior border extending on the fringes, one between each nerve; near the anal angle between nerves 1 and 2 a large and conspicuous spot bearing a partial resemblance to a Greek *psi*, placed sidewise, and from this spot a somewhat zigzag line running parallel with posterior border, but somewhat more arcuated towards costa, least distinct between nerves 3 and 4, and forming a large distinct dart-like spot between nerves 5 and 6; space between this line and posterior border, slightly darker than the rest of the wing-surface on account of the dark atoms being more thickly sprinkled over it; four costal marks, one subobsolete in a transverse line with the reniform spot, one conspicuous about the middle, and in a line with reniform spot and anal angle, one about the same size as the last and looking like a blurred X about one-third the length of wing from base, and one subobsolete, near the base; orbicular spot flattened and well defined by a black annulation; reniform spot indicated by a blurred black mark running on the cross-vein and sometimes somewhat crescent-formed; a V-shaped spot pointing towards base half-way between costa and interior margin, in a transverse line with the large costal spot which looks like a blurred X; a blurred mark in middle at base, and lastly a narrow spot on the inferior margin, half-way between base and anal angle. Hind wings same color as front wings; somewhat more glossy, with the lunule, a band on posterior border one-fourth the width of wing, and sometimes a narrow coincident inner line, somewhat darker than the rest; the posterior border also with a series of spots one between each nerve. Under surface of front wings pearly-white with an arcuated brown band, most distinct towards costa, across the posterior one-third, all inside of this band of a faint yellowish-brown; lunule and fringe spots distinct, and with a faint trace of the *psi*-spot; hind wings uniform pearly-white with a distinct and well defined dark wavy line running parallel with posterior margin across the posterior one-third of wing, and with the lunule and fringe spots distinct. Antennæ simple and bristle-formed, gray above, brown beneath. Head thorax and body, both above and below, silvery-gray. Legs with the tarsi alternately dusky and gray. ♂ differs from ♀ by his somewhat stouter antennæ; much narrower body, and narrower wings and fringes, the front wings having the apex more acuminate, and the hind wings scarcely showing the darker hind border.

Described from 2 ♀, 2 ♂ all bred. In the ornamentation of the front wings this species bears some resemblance to the European species *tridens* and *psi*, but otherwise differs remarkably, and especially in its larval characters. It bears a still closer resemblance both in the larva and imago state to the pale variety of a common species known in England as the "Miller" (*A. leporina*), but judging from the figures and description in "Newman's Natural History of British Moths," it may be easily distinguished from *leporina* by the well defined orbicular spot, by the greater proximity of the two large costal spots, by lacking a round spot behind the disk, and by the more prolonged apex. It differs also in the larva state from *leporina* which feeds on the Birch. It likewise closely resembles *interrupta*, though the larvæ are remarkably different; and it also resembles *lepusculina*, the larva of which is unknown; but the specific differences will be readily perceived upon comparing Guenée's descriptions. How near it approaches to *Acronycta occidentalis*, Grote, it is impossible to tell, as the author's description is exceedingly brief, considering the number of closely allied forms; but as that species has a bright testaceous tinge on the reniform spot, it evidently differs from mine. Harris's *Apatela* [*Acronycta*] *Americana*, though very differ-

ent in the imago, yet closely resembles *populi* in the larva state. I have on two occasions found the larva of *Americana* feeding on the Soft Maple, and it may be distinguished from *populi*, by its greater size; by the paler color of the body; by the hairs being paler, more numerous, shorter and pointing in all directions, especially anteriorly and posteriorly of each segment; by having on each of joints 4 and 6 two distinct long black pencils, one originating each side of dorsum, and on joints 11 one thicker one originating from the top of dorsum; by a substigmatal row of small black spots (three to each segment, the middle one lower than the others) and by a trapezoidal velvety black patch starting from anterior portion of joint 11 and widening to anus.—[Second Rept., pp. 120, 121. Figs. 87, 88.]

Grote refers it, in his *List*, to *lepusculina* Gn.; having, I believe, seen the type. Guenée must have had a uniformly colored and pale specimen as my typical specimens have a distinct orbicular mark, deeper subterminal markings and the terminal space contrasting by its darker gray with the rest of primaries—all unmentioned in Guenée's description.

XYLINA CINEREA, N. Sp.—*Larva*—Length when full grown 1.20—1.30 inches [=30—32^{mm}], color shiny silvery-green on the back, darker below. A medio-dorsal cream-colored stripe; a subdorsal one represented by 3 or 4 irregularly shaped spots on each joint. A broad deep cream-colored stigmatal line, with a few green dents in it, extending to anal prolegs. Four slightly elevated cream-colored spots, encircled by a ring of rather darker green than the body, in the dorsal space, and in the subdorsal space there are four or more similar but smaller spots. Venter glaucous-gray. Head as large as joint 1, free, glassy-green with white mottlings at sides and top, and pearly-white lips. Thoracic legs whitish. Prolegs concolorous with venter. When young the body is darker and the markings paler. Described from two living specimens.

Imago (Fig. 57, b)—*Front wings*, with the ground-color pale cinereous shaded and marked either with light brown, having a faint purplish tint, or with darker brown, having a similar reflection, or with a colder grayish-brown with the faintest moss-green reflection: in the first two cases the dark color either blends and suffuses with the ground-color so as to give the wing a nearly uniform and smooth appearance, or else contrasts sufficiently to bring out all the marks distinct; in the latter case (two specimens) the markings are very distinct and the ground color is whiter and more irrorate. In the well-marked specimens the usual lines are readily distinguished, the basal half line, transverse anterior and transverse posterior being quite wavy, pale, and bordered each side with a dark shade, the median shade dark and well defined and the subterminal line, though sometimes pale near costa, forming a series of dark angular spots: in the more uniform specimens these lines are barely distinguishable and perhaps the most constant is the sub-terminal which most often takes the form of a series of dark angular spots: the ordinary spots have a pale inner and a more or less distinct dark outer annulation; the orbicular is larger than the reniform and is sufficiently double to take on the form of an 8, the upper part of which is always largest and with the interior space paler than the general surface, while that of the lower part is either concolorous or darker; the form is, however, quite irregular and differs sometimes in the two wings of the same species: the reniform spot is generally well defined, and is either darker, or has a tinge of reddish-brown, interiorly: at the base of the wing is a more or less distinct pale space occupying the upper half, and bordered below by a brown line which is straight about half its length and then extends upwards and outwards towards transverse anterior. A tolerably distinct terminal line, with the fringes dark. In taking a general view of the varying specimens this pale basal space, the pale upper part of the orbicular and the dark subterminal line, seem to be the most constant characters of the species. *Hind wings* gray-brown in-

clining to cinnamon-brown, with the posterior border but slightly darker and the fringe paler. Under surface quite uniform, that of front wings being nacreous gray with a faint discal spot and with a narrow costal and broad terminal border of pale fulvous, dusted with purple-gray; the hind wings of this last color with the lunule and line distinct. *Head* nearly entire, though the quadrifid arrangement of the hairs is traceable; palpi hairy throughout. *Thorax* quite square, of same color as primaries and with the collar bordered behind with brown and sometimes the edges of the tegulae similarly bordered. *Abdomen* of same color as hind wings with lateral tufts, and cut off squarely at apex. Expanse 1.32—1.82 inches [= 34—45^{mm}].

Described from 3 specimens fed on grape-vine, 2 on peaches and 1 on *Cercis canadensis*. Other captured specimens examined.

This species is the analogue of, and very closely resembles the European *Xylina conformis*, which is known under various synonyms. A specimen sent to Mr. P. C. Zeller of Stettin, Prussia, was, however, pronounced distinct. The well-marked irrorate form still more closely resembles Guenée's *cinerosa* found in Switzerland, and which he himself thinks may prove to be a variety of *conformis*. The more I study the species of the NOCTUIDÆ as they occur in nature, the more I am struck with their great variability, and there can be no doubt that many of the so-called species will turn out to be but varieties when we better understand them. In this large family none but the more strikingly marked species should ever be described without an accompanying description of their preparatory states and of their principal variations. I am unacquainted with any of Walker's species except *subcostalis*, which is very different, and if this should prove to be a synonym of any of them the fault must be laid to the difficulty under which the naturalist in the Western States labors for want of proper libraries to refer to. It differs essentially from Grote's *Bethunei* and *capax* as described and illustrated in Volume I of the Transactions of the American Entomological Society. I am informed by Mr. [J.] A. Lintner of Albany, N. Y., that Dr. A. Speyer of Rhoden, Fürstenthum Waldeck, Prussia, who gives much attention to the Noctuidæ, has it marked *Celæna oblonga* in his MS., but the insect evidently does not belong to that genus, and as the German pronunciation of *Xylina* much resembles the English pronunciation of *Celæna*, the reference to the latter is doubtless due to a verbal misunderstanding.—[Third Rept., pp. 135, 136. Fig. 57.]

Now referred, in Grote's *List of Noctuidæ of N. A.*, to Hübner's genus *Lithophane*.

AMPHIPYRA CONSPERSA, N. Sp.—*Larva*.—Found full grown July 2, 1867, on Hazel. No pyramidal hump, and of a uniform emerald-green, the dorsal palpitations visible and the stigmata pale, with a black annulation, but with no other markings either on the head, body, or legs.

Imago—Like *pyramidoides* in every particular except that the brown of front wings is almost uniformly spattered over, more or less suffusely, with pale-grayish spots, so that no regular marks appear. The costal marks are, however, tolerably distinct as in *pyramidoides*, and by careful examination and comparison traces of the more conspicuous marks of that species may be discerned.

Described from one ♀ bred July 31.—[Third Rept., p. 75.]

As remarked at the time, the specimen from which the description was made was a bred one and perfect. Grote, in his *List of Noctuidæ*, considers it simply an aberration of *pyramidoides*, but this can hardly be the case, as the larva also shows differences.

AGROTIS SCANDENS, N. Sp.—*Larva*.—Average length when full grown 1.40 [inch, = 35^{mm}]. Ground-color very light yellowish gray, variegated with glaucous in the shape of different sized patches, which are distinctly seen under the lens to be separated by fine lines of the light ground-color. A well-defined dorsal and less distinct

subdorsal and stigmatal line, caused by these patches becoming larger and darker; another and still less distinct line of the same kind under stigmata. The dorsal line frequently with a very fine white line along its middle, especially at sutures of segments. Piliferous spots in the normal position; those above black, those at the sides lighter. Stigmata black. Head and cervical shield tawny, the latter with a small black spot each side, the former with two in front, and two eye-spots each side. Caudal plate tawny, speckled with black. Venter and legs glaucous. Bristles fine and small. Filled with food it wears a much greener appearance than otherwise, while when young it is of a more uniform dirty whitish-yellow, the lines less distinct but the piliferous spots proportionately larger. Head quite variable in depth of shade.

Perfect Insect.—Average length 0.70 [inch, = 17.5^{mm}]; alar expanse 1.50 [inch, = 37^{mm}]. General color of fore wings very light pearly bluish-gray, with a perceptible deepening posteriorly. Quite variable, sometimes of a more decided blue, at others inclining to buff as in *Leucania unipuncta*, Haw. Markings, when distinct, as in Plate 1, Figures 5 and 6. With the exception of the reniform spot and subterminal line, however, they are usually distinct only on costa, being either indistinct or entirely obsolete on the rest of the wing. The subterminal line is light, with a more or less dark diffuse shade each side, which, in some instances, forms into sagittate spots. A black stain at the lower part of reniform spot forms a most distinctive character. Hind wings very pale and lacking the bluish cast of fore wings; lunule distinct, and a dark shade, enclosing a lighter mark, as in *Heliothis*, along posterior margin. Eyes dark; head and thorax same as fore wings; abdomen same as hind wings. The whole under surface the same as hind wings above, the lunules and arcuated bands faintly traced, the fore wings having a darker shade in the middle.

Described from 30 bred specimens.—[First Rept., pp. 78-79. Pl. 1, Figs. 5, 6.]

AGROTIS COCHRANII, Riley.—*Imago*.—Fore wings of a light warm cinereous, shaded with vandyke brown and umber, the terminal space, except at apex, being darker and smoky. Basal, middle and limbal areas of almost equal width, the middle exceeding somewhat the others. A geminate dark basal half-line, usually quite distinct. Transverse anterior geminate, dark, somewhat irregularly undulate, and slightly obliquing outwards from costa to interior margin. Transverse posterior geminate, the inner line being dark, distinct and regularly undulate between the nerves, while the outer line is plain and much paler; it is arcuated superiorly and inversely obliques for two-thirds its width. Orbicular and reniform spots of normal shape, having a fine, dark annulation, which is however obsolete in both, anteriorly; the orbicular is concolorous with the wing, whilst the reniform has a dark inner shade with a central light one, and forms with the transverse posterior a somewhat oval spot which is also dark. Median shade dark and distinct interiorly, shading off and becoming indistinct in center of wing, and quite dark between the two spots, giving them a fair relief. Subterminal line single, light, acutely and irregularly dentate, with an inner dark shade, but warmer than that of terminal space. Terminal line very fine, almost black, slightly undulate. Fringes of same color as wing, with a light central line, having an outer dark coincident shade. A dark costal spot in basal area; at termini of the usual lines, and two light ones in subterminal space. In some specimens one or two fine dark sagittate marks are discernable, and also a fine black claviform mark. Hind wings: whitish, with a darker shade along posterior margin. Under surface of fore wings somewhat lighter than the upper surface and pearlaceous interiorly, with a smoky arcuated band—more definite near the costa than elsewhere—and a tolerably distinct lunule. Under surface of hind wings concolorous; slightly irrorate with brown anteriorly and posteriorly, and with an indistinct lunule and band. Antennæ, prothorax, thorax, tegulæ and body of same color as primaries, the prothorax having a darker central line, and in common with the tegulæ a carneous margin. Under surface lighter; legs with the tarsi spotted.

This moth, in its general appearance, bears a great resemblance to *Hadena chenopodii*, but the two are found to differ essentially when compared. From specimens of *H.*

chenopodii, kindly furnished me by Mr. Walsh, and named by Grote, I am enabled to give the essential differences, which are: 1st. In *A. Cochranii*, as already stated, the middle area exceeds somewhat in width either of the other two, while in *H. chenopodii* it is but half as wide as either. 2d. In the *Agrotis* the space between the spots and between the reniform and transverse posterior is dark, relieving the spots and giving them a light appearance, whilst in the *Hadena* this space is of the same color as the wing, and the reniform spot is dark. The claviform spot in the *Hadena* is also quite prominent, and one of its distinctive features, while in the *Agrotis* it is just about obsolete.

There are specimens that seem to be intermediate between these two, but all those bred by me, both male and female, were quite constant in their markings, and their intermediates will doubtless prove to be distinct species or mere varieties.

Larva—Length 1.07 inches [= 26.8mm]. Slightly shagreened. General color, dingy ash-gray, with lighter or darker shadings. Dorsum light, inclining to flesh color, with a darker dingy line along its middle. The sides, particularly along the sub-dorsal line are of a darker shade. On each segment there are eight small, black, shiny, slightly elevated points, having the appearance of black sealing-wax, from each of which originates a small black bristle. The stigmata are of the same black color, and one of the black spots is placed quite close to them anteriorly. Head shiny and of the same dingy color as the body, with two darker marks, thick and almost joining at the upper surface, becoming thinner below and diverging toward the palpi. The upper surface of first segment is also shiny like the head. Ventral region of the same dingy color, but lighter, having a greenish tinge anteriorly and inclining to yellow under the anal segment. Legs of same color. It has a few short bristles on the anterior and posterior segments.

Chrysalis.—Length 0.70 of an inch [= 17.5mm]. Light yellowish brown with a dusky line along top of abdomen. Joints, especially of the three segments immediately behind the wing-sheaths, dark brown. The brown part of these three segments, minutely punctured on the back. Eyes dark brown, and just above them, a smaller brownish spot. Two quite minute bristles at extremity.

Described from numerous bred specimens. — [First Rept., pp. 75–76. Fig. 26.]

There is little question but that this is the moth briefly characterized by Harris (Ins. Inj. to Veg., p. 444) as *Agrotis messoria*, an examination of the types confirming this view. *A. repentis* G. & R. and *A. lycarum* are also conceded by Grote to be synonyms.

PLUSIA BRASSICÆ, N. Sp. — *Larva*—Pale yellowish translucent green, the dorsum made lighter and less translucent by longitudinal opaque lines of a whitish-green; these consist each side, of a rather dark vesicular dorsal line, and of two very fine light lines, with an intermediate broad one. Tapers gradually from segments 1–10, descending abruptly from 11 to extremity. Piliferous spots white, giving rise to hairs, sometimes black, sometimes light colored; and laterally a few scattering white specks in addition to these spots. A rather indistinct narrow, pale stigmatal line, with a darker shade above it. Head and legs translucent yellowish-green, the head having five minute black eyelets each side, which are not readily noticed with the naked eye. Some specimens are of a beautiful emerald-green, and lack entirely the pale longitudinal lines. Described from numerous specimens.

Chrysalis—Of the normal *Plusia*-form, and varying from yellowish-green to brown.

Moth—*Front wings* dark gray inclining to brown, the basal half line, transverse anterior, transverse posterior, and subterminal lines pale yellow inclining to fulvous, irregularly undulate, and relieved more or less by deep brown margins; the undulations of the subterminal line more acuminate than in the others, and forming some dark sagittate points; the basal half-line, the transverse anterior near costa, and the transverse posterior its whole length, being sometimes obscurely double: four distinct equidistant costal spots on the terminal half of wing, the third from apex formed by

the termination of the transverse posterior; posterior border undulate with a dark brown line which is sometimes marked with pale crescents; a series of similar crescents (often mere dots) just inside the terminal space; the small sub-cellular silver spot oval, sometimes uniformly silvery-white but more often with a fulvous centre, sometimes free from, but more often attached to the larger one which has the shape of a constricted U, very generally with a fulvous mark inside, which extends basally to the transverse anterior at costa. Fringes dentate, of the color of the wing, and with a single undulating line parallel to that on the terminal border. *Hind wings* fuliginous, inclining to yellowish towards base, and with but a slight pearly lustre; fringes very pale with a darker inner line. Under surfaces pale fuliginous with a pearly lustre, the front wings with a distinct fulvous mark under the sub-cellular spots, speckled more or less with the same color around the borders of the wing, the fringes being dentate with light and dark; the hind wings speckled with fulvous on their basal half, and with the fringes as above. *Thorax* variegated with the same color as front wings, the tufts being fulvous inclining to pink. *Abdomen* ♀ gray, with a few pale hairs near the base, and scarcely extending beyond the margin of the hind wings; ♂ longer, covered with pale silky hairs, a distinct dorsal brown tuft on each of the three basal segments, and two large lateral either fawn-colored or golden-yellow brushes on the fifth segment, meeting on the back and partly covering two smaller brushes on the sixth, which are tipped with black; terminal segment flattened and with two lateral more dusky and smaller tufts: underside of thorax and abdomen gray, mixed with flesh-color. Alar expanse 1.55 inches. Described from numerous bred specimens. In a suite of specimens bred from the same brood of larvæ a considerable difference in the general depth of color is found, some being fully as dark again as others.

Closely resembles *Plusia ni*, Engr., which occurs in Italy, Sicily, France, and the northern parts of America. Mr. P. Zeller of Stettin, Prussia, to whom I sent specimens, considers it distinct however from the European *ni*, and I have consequently given it a name in accordance with its habits. — [Second Rept., pp. 111-112. Fig. 81.]

Notwithstanding its close resemblance to *ni*, the best authorities agree with Zeller in considering it distinct, as it certainly is. Strangely enough this same *brassicæ*, or what is extremely close to it, occurs also in South Europe and is figured in Stainton's *Entomologist's Annual* for 1870 as *P. ni*, one specimen having been found on the south coast of England, which specimen Zeller, as he wrote me, believes to have come from America. Staudinger would probably characterize *brassicæ* as a "species Darwiniana," and there are doubtless individuals of both the species which approach each other so closely as to be undistinguishable. There is such variation in the silver spot in either that it cannot be depended on alone, but Speyer (*Europäisch-Americanische Verwandtschaften; Stettiner Ent. Zeit.*, June, 1875, p. 165) has presented other differences that are constant in detail, the most noticeable of which are the darker and more irrorate coloring and the interrupted and wavy terminal line of *brassicæ*, against the paler, smoother, more metallic coloring and the perfectly straight and unbroken terminal line of *ni*.

The larva is the most common cabbage pest in the Southern States, and is infested with an undetermined parasite. Mr. E. A. Popenoe has found it feeding on the leaves of *Crepis*, and what appears to be the same has been found by my assistants on Clover, Dandelion, *Senecio scandens*, and *Chenopodium*.

APLODES RUBIVORA, N. Sp. — *Larva* — Average length 0.80 inch [= 20^{mm}]. Color light yellowish-gray, darker just behind each joint, and very minutely shagreened all over. On each segment a prominent pointed straight projection each side of dorsum, and several minor warts and prickles below. Two very slightly raised, longitudinal lighter lines along dorsum, between the prominent prickles. Ten legs.

Perfect insect — Alar expanse 0.50 inch [= 12.5^{mm}]; length of body 0.25 inch [= 6^{mm}]. Color verdigris-green, the scales being sparse so that the wings appear sub-hyaline. Fore-wings with two transverse lighter lines dividing the wing into three parts, proportionate in width as 3, 4, 2 counting from base, and parallel with posterior margin; also a faint line between these two, running to about $\frac{1}{3}$ of wing from costa. Hind wings with two similar transverse lines, dividing the wing in like proportion, the outer line not parallel with margin, but wavy and produced posteriorly near its middle. Costa pale; fringes obsolete. Head, thorax and abdomen green above, but, together with antennæ and palpi, white beneath.

Described from one ♀ specimen. — [First Rept., pp. 139-140. Pl. II, Fig. 25.

Dr. Packard, in his *Monograph of the Geometrid Moths, etc.* (U. S. Geol. Surv. of Terr., Vol. X, 1876, p. 382), refers it to the genus *Synchlora* Gor., and adds the conventional ending to the specific name, so that the species becomes *Synchlora rubivoraria*. *Synchlora albolineata* Pack. and *Eunemoria gracilaria* Pack. are given as synonyms.

PHYCITA [ACROBASIS] NEBULO, Walsh — *Imago*. — I reproduce here the description of the moth in Mr. Walsh's original words: "Expansion of wings 7-10. Length of body 3-10. General color light cinereous, varied with dusky. A row of about seven sub-semilunar or linear dark spots on outer margin of fore wing. Then one-fourth of the distance to the body a waving light cinereous band parallel to the exterior margin, marked on each side with dusky black. Nearly at the centre a much abbreviated black band. Beyond the centre on the costal margin a subtriangular dusky black spot, the apex of which connects with the apex of a much larger subobsolete triangular brick-red spot which extends to the interior margin, and is bounded on the outside by a wavy light cinereous band, which is again bounded by a wavy dusky black band proceeding from the apex of the costal triangle. Base of wing dusky black, inclosing a small round light cinereous spot. Hind wings and all beneath light cinereous shaded with dusky, the fore wings darker. Tarsi dusky with a narrow light cinereous fascia at the apex of each joint. Hind tibia fasciate with dusky at the apex, sometimes obscurely bifasciate. Intermediate tibia fasciate with dusky at the centre, the fascia generally extending to the base, but becoming lighter. Anterior tibia dusky, with a narrow apical light cinereous fascia. Palpi, both labial and maxillary, dusky."

When compared with other closely allied and resembling species, this little moth may be characterized in the following manner: The ground color of the front wing is decidedly bright and pale; the discal spots are almost always confluent, thus forming an abbreviated transverse bar; the dark markings are well defined and the triangular dark costal spots starting from the inner third of the wing is distinctly relieved, while the "brick-red" (nearer a cinnamon-brown) triangular spot which opposes it is large, so that the space it occupies on the inner margin is nearly as wide (generally within one-third) as that between it and the transverse posterior line. The lower half of the basal space is often of a distinct cinnamon-brown, and an oblique dusky band, which Mr. Walsh has not mentioned, is often quite distinct, running from near the apex to the brown triangle, where it connects with the inner margin. The species recalls, in facies, the European *Myelois suavella*. In a suite of specimens bred from Apple, Quince, Plum and Cherry, there is sufficient variation to prevent a too rigidly drawn description, but the above characters obtain in all of them, and such variation as occurs runs in the direction of the variety presently to be described.

Larva — [Length 0.5 inch] Brown or greenish in color. Cylindrical. Tapering grad-

ually from first to last joint. Head and cervical shield darker than the rest of body, slightly shagreened, sparsely covered with long hairs, the shield quite large, convex, and occupying the whole surface between stigmata—there being in front of the latter a sub-cervical dark horny plate. Joints 2 and 3 wrinkled as at Fig. 18, *c* the former with two rather conspicuous dark dorsal piliferous spots. The other joints with a few fine hairs, the stigmata plainly visible, and the anal covering but slightly horny. Legs and prolegs of moderate size and of same color as body.

Described from numerous specimens.

Chrysalis—Mahogany-brown, with no striking character. Abdomen, especially above, with very minute punctures.

VARIETY NEBULELLA (Fig. 20, *e*).—I have bred a single specimen from wild 'Crab (*Cratægus*) which differs in some essential features from the normal form, but which nevertheless can only be considered a variety of it, as I observed no larval differences. It differs in the more uniform and subdued tone of the front wings, the markings being more suffused and indistinct; but principally in the relative narrowness of the space outside the transverse posterior line the greater consequent width of the middle area, and smallness of the triangular brown spot—the space it occupies on the inner margin being scarcely one-half as wide as that between it and the transverse posterior line. The discal spots are also separated.

Described from one good specimen. An interesting fact connected with this variety is, that precisely the same form occurs in Europe, as I found a single specimen in the cabinet of M. J. Lichtenstein of Montpellier, France, which he had captured in that vicinity, and which he allowed me to bring home for comparison. It seems to be rare, even there, and whether indigenous or imported from this country, is a question yet to be solved.—[Fourth Rept., p. 41-42, Figs. 18, 19, 20.

ACROBASIS JUGLANDIS, LeBaron.—(Fig. 20, *d*)—I have bred this species from Hickory, but as Dr. LeBaron has also bred it abundantly from Walnut, and has signified his intention of describing it in his second annual Report, I adopt his proposed name, and shall content myself with pointing out the manner in which it may generally be distinguished from *nebulo*. Firstly, by the paler basal area of the front wings, which is sometimes almost white, especially near the costa, and by the head and shoulders and sometimes the ♂ antennal horn partaking of this paler color. Secondly, by the darker median space, the dark triangular costal spot not being well relieved posteriorly, but extending so as sometimes to darken the whole space. Thirdly, by the discal spots always being well separated.

Such are its specific characters as taken from 3 hickory-bred and 6 walnut-bred specimens; but of the former there is 1 which when placed alongside of some of the more abnormal specimens of *nebulo*, can scarcely be distinguished from them, and, if chosen without knowledge of its larva, would certainly be placed with them; while of the latter there are two which nearly as closely resemble the variety *nebulella*. In general characters, in the size of the brown triangular spot, and the manner in which the inner margin is divided, *juglandis* is intermediate between *nebulo* and *nebulella*. In one of the hickory-bred specimens, the general color is quite warm, and the basal area carneous rather than white.—[Fourth Rept., p. 43. Fig. 20, *a*, *b*, *d*.

Dr. LeBaron published his description of it about the same time, under the name *Phycita juglandis*, in his Second Report on the Insects of Illinois, p. 123.

PEMPELIA HAMMONDI, N. Sp. *Imago* (Fig. 21, *d*).—Average expanse 0.48 inch [=12^{mm}]. Front wings glossy purplish-brown with two silvery gray transverse bands dividing the wing on costa in about three equal parts, the basal band sharply defined outwardly and always extending to inner margin, the posterior band never extending more than half way across the wing, and generally not more than one-third, illy defined. In some specimens the basal transverse band is quite narrow, with the basal space a shade paler than the median: in others the band forms a double line. In some

specimens also, a narrow pale transverse line outside the second band, and a pale terminal shade, are visible. Hind wings uniformly paler gray. Under surface glossy gray, with no marks, the front wings a shade darker than the hind. ♂ differs from ♀ in the basal portion of the antennæ being curved, and the curve filled with a tuft of scales.

Described from numerous bred specimens. The species has the general facies of the European *Cryptoblabes bistriga*, which is a larger insect.

Larva.—Length 0.45—0.50 inch [=11—12.5^{mm}]. General color olive, or pale green, or brown, with a broad dark stripe along each side of back. Tapers slightly both ways, joints 4-12 inclusive, divided into two transverse folds. Freckled with numerous pale specks and with piliferous spots, the specks often taking the form of two pale broken lines along the upper edge of dark stripe. The piliferous spots are pale with a central black dot, and are best seen in the dark specimens. On joints 4-12 inclusive they are placed 4 in a square on the middle of the back, and four more each side, the two upper lateral ones being on the anterior fold, the stigmata appearing as minute rufous specks between them. Both these spots are often double. The third lateral spot is on the posterior fold and the fourth is subventral and anterior. The hairs proceeding from these spots are long and setaceous. Head horizontal, freckled, pale behind, tinged with green in front and with a few long hairs. Joint 1 also freckled and with a large black piliferous tubercle with a pale basal annulation and in range with middle of dark stripe. Joint 2 with similar black tubercles with a white centre and replacing the uppermost lateral pale spot. There are but two of the small pale dorsal piliferous spots on this joint (between the tubercles) as well as on joint 3. Beneath immaculate, except that the thoracic legs have sometimes a few dusky dots.

In the very dark specimens the head, cervical shield and anal plate remain pale. The cervical shield is then well defined with four small piliferous specks at anterior edge, and the large shiny tubercle forms the extreme anterior angle.

Described from numerous specimens.

Pupa.—0.24 inch [=6^{mm}] long; rather stout and short, with two minute diverging spines and a few stiff bristles at tip.

In many specimens the subdorsal dark stripe is obsolete or sub-obsolete, but even then the four black tubercles on joints 1 and 2 characterize the larva sufficiently.—[Fourth Rept., p. 46. Fig. 21.]

TORTRIX RILEYANA, Grote—*Larva*—Length, Hickory feeding, 0.60-0.80 inch [=15—20^{mm}]; Snowberry feeding, 0.40-0.50 inch [=10—12.5^{mm}]. Largest on segment 2, tapering thence gradually to anus. Ground color dull yellow. Covered with large, distinct, black, sealing-wax-like, slightly elevated spots, each giving rise to several fine bristles. These spots are thus arranged on each segment: 2 each side of dorsum the posterior ones widest apart; 1 at sides in the middle of the segment, containing the stigmata in its lower hind margin; 1 smaller and narrower just below this, on a somewhat elevated longitudinal ridge, and 1 round one below this ridge on the posterior part of the segment. Segments 2 and 3 have but one spot each side of dorsum. Two distinct wrinkles on all the segments, more on 2 and 3. Head, cervical shield, and caudal plate black. Venter dirty yellow with black marks; legs ditto.

Chrysalis—Honey-yellow, robust in the middle, and with two transverse rows of minute teeth across the back of each segment.

Perfect Insect—From Hickory—Average expanse 1 inch, length of body, 0.35 [=8.8^{mm}]. Deep ochreous. Fore wings evenly washed with purplish, leaving the fringes and costal edge dark ochreous. The markings take the shape of dark velvety brown rounded maculations, generally of small size and faintly shaded with ochreous on the edges. Three of these subterminally at the base of the wing, subequal, situated interspaceally between the nervures. At a little within the middle of the costa are two fused maculations, the most prominent. Before and beyond these, some faint costal marks. At the extremity of the discal cell, above median nervure, is the first of a

series of maculations, normally four in number but not constant, usually uneven in size. A subterminal series of spots is inaugurated on costa by a large, compound shaded maculation. Below this, over the median nervules, sweeps an outwardly rounded series of small approximate dots. Two dots on costa, within and at the apex, and a faint terminal series of minute streaks is shortly discontinued. Hind wings of a lustrous bright deep ochreous; pale along the costal margin and darker shaded along internal margin. Beneath, as are the hind wings above; both wings immaculate, fore wings the darker. Body and appendages concolorous, bright deep ochreous. Antennæ simple. Numerous bred specimens.

From Snowberry—var. symphoricarpi—Much paler, the fore wings not being as dark as the hind wings of the above. The upper surface of fore wings not washed with purplish but merely of a darker ochreous than the hind wing. The maculations entirely similar but ferruginous, paler and the slighter costal marks obsolete. Legs at base and under thoracic surface almost whitish. Average expanse, 0.62 [=15.5^{mm}]; length of body, 0.30 [=7.5^{mm}]. Described from numerous specimens. Under surfaces exactly alike in both varieties.—[First Rept., p. 154. Fig. 85, and Pl. 2, Figs. 3, 4.

TORTRIX CINDERELLA, N. Sp.—*Imago*.—Alar expanse exactly 1-2 inch [=12.5^{mm}]. Front wings deep glossy ash-gray, immaculate. Under a lens they have an irrorate appearance, while in certain lights some of the scales appear to form a series of darker transverse sinuous lines. Also scattered over the wing may be noticed a dozen or more reddish scales, which are not sufficient, however, to destroy the uniform immaculate appearance. Head, mouth-parts, antennæ, legs, and abdomen of same color. Hind wings paler and semi-transparent. Fringes of all wings concolorous. Under surface of wings pale nacreous, inclining to pale fulvous around the margins.

Described from two bred specimens.

Larva (Fig. 22, a).—Length 0.50 inch [=12.5^{mm}]. Form of that of *Acrobasis nebulo*, wrinkled very much in the same manner. Color yellowish-green, the piliferous spots of the same color, but readily distinguished by their polished surface; they are placed in a transverse row on thoracic joints, and on joints 4—12 there are four trapezoidally on dorsum, two laterally on the first fold and one subventral. Stigmata between the two lateral spots, and yellowish. Head and cervical shield gamboge-yellow; only a shade darker than body; labrum and two basal joints of antennæ paler or white, the terminal joint brown; ocelli on a somewhat crescent-shaped black spot (the most conspicuous character) a second dusky spot at base of head laterally. Legs immaculate.

Described from many specimens.

Pupa (Fig. 22, b).—Length 0.25—0.30 inch [=6—7.5^{mm}]. Brown, characterized by a peculiar rounded projection from front of head; by a little pointed prominence at base of each antennæ, and each side of penultimate abdominal joint; and by terminating in a broad suppressed piece which produces two decurved hooks. Posterior rim of abdominal joints rasped dorsally, and a slight rasped dorsal ridge near the anterior edge of larger joints. Legs reaching only to end of wing-sheaths. The head-prominence varies in size and slightly in form.—[Fourth Rept., p. 47.

From specimens reared from cranberry-feeding larvæ received from Mr. Jno. H. Brakeley, of Bordentown, N. J., I am satisfied that this is the same species briefly characterized by Packard in the 1st edition of his *Guide* (p. 334) as *Tortrix oxycoccana*, and that *T. malivorana* LeBaron (my Rep. IV, p. 47) is but a dimorphic orange form, subsequently described by Packard as *T. vacciniivorana* (Hayden's Report of the U. S. Geol. and Geogr. Survey of the Territories 1878, p. 522). The orange and ash-gray specimens are thus bred both from Apple and Cranberry. I have reared both forms from Cranberry and from Apple, and they are undistinguishable in the larva and pupa states. The gray form is often

more or less suffused with orange scales and the orange form less frequently with gray scales. This is the most remarkable case of dimorphism with which I am familiar in the family, and points strongly to the important bearing of biological facts on a true classification. The dimorphic coloring is not sexual, but occurs in both sexes. The eggs of this species are very flat, circular and translucent, with a diameter of 0.7^{mm} , and are laid singly on the underside of the leaf near the mid rib. The species belongs to the genus *Teras*, and as Packard's specific name *oxycoccana* has priority, the insect should be known as *Teras oxycoccana*, Pack. The insect, according to Mr. Brakeley, who gives an account of it in the Report of the Seventh Annual Convention of the New Jersey Cranberry Association (1879, p. 7), commonly affects, also, the high-bush whortleberry. The gray form of the moth is most frequent in autumn.

GELECHIA GALLÆSOLIDAGINIS, N. Sp.—*Larva*.—Length 0.60 [inch, = 15^{mm}]. Cylindrical. Color dark dull-brown, without shine. Largest on middle segments; tapering from 4th to head, and from 9th to extremity. Each segment impressed transversely in the middle, thus forming two folds, the thoracic segment having other such folds. Six small piliferous spots, two each side of dorsum and one above stigmata, which, together with the stigmata, are shiny and of a lighter brown than the body. Head and cervical shield light shiny-brown.

Chrysalis.—Length 0.50 [inch, = 12.5^{mm}]. Mahogany-brown. Form normal. Blunt at extremity.

Perfect moth.—Average length 0.38 [= 9.5^{mm}]. Alar expanse ♀ 0.95 [inch, = 24^{mm}], ♂ 0.75 [inch, = 18.8^{mm}]. Fore wings deep purplish-brown, more or less sprinkled with carneous. A light carneous band starts from the costa near the base, and curves towards the middle of the inner margin, which it occupies to a little beyond the beginning of the cilia, where it curves upwards towards the tip, reaching only half way up the wing. Here it is approached from above by a somewhat diffuse spot of the same color, which starts from the costa just behind the apex, and runs down to the middle of the wing.

In the plainly marked individuals there is an extra line running from the middle of the inner margin, outwardly obliquing to the middle of the wing, and then back to the inner margin a little beyond where the cilia commences, but in the great majority of specimens this mark is indistinct. Cilia light carneous. Hind wings slate-gray, with the cilia lighter. Antennæ finely annulated with the same two dark and light colors. Head, thorax and palpi light, with a sprinkling of the dark brown. Body dark, with light annulations. The species varies in the distinctness of its markings, and the light parts of the front wing appear finely sprinkled with brown under the lens. Male generally smaller than female, with the antennæ proportionately a little longer.

Described from numerous bred specimens.

It seems to resemble *G. longifasciella* of Clemens, in coloration and pattern; but unfortunately our late lamented microlepidopterist, failed almost always to give the measurement of the species he described, and it is impossible to tell how much mine resembles that species. Yet, as *longifasciella* was described from two mutilated specimens, received from A. S. Packard, jr., and as that gentleman has seen my insect and declared it an undescribed species, there can be little doubt of the fact.—[First Rept., p. 175. Pl. II, Figs. 1, 2, 5.]

PTEROPHORUS CARDUI, N. Sp.—*Larva*.—Average length 0.60. Largest in the middle of body, tapering thence each way. Color light straw-yellow—greener when young. Somewhat darker, partly translucent, dorsal, subdorsal and stigmatal lines. Two lateral rows of black spots, the lower spots rather smaller and placed behind the

upper ones. A third row above these, and others along the back, but so small that they are generally imperceptible with the naked eye, except on the thoracic segments, being especially distinct on segment 2. Head small, black, sometimes inclining to brown. Cervical shield black, divided longitudinally in the middle by a lighter line. Caudal plate also black. Segment 11, besides the spots above mentioned, has two transverse black marks, the posterior one the largest. Thoracic legs black, the others of the same color as the body.

Described from 12 specimens.

Pupa.—Average length 0.45. Of form of Plate 2, Fig. 14. Soft, dull yellow, with a lateral dusky line each side of dorsum, and another, less distinct, each side of venter. Also dusky about the head and wing-sheaths.

Perfect insect.—Length 0.45; alar expanse 0.80. Front wings bifid, the cleft reaching not much more than $\frac{1}{4}$ of wing; tawny yellow, with a distinct dark brown triangular spot running from costa to the base of cleft—sometimes a little below it—its posterior margin with a slight concave curve. Three dusky, diffuse longitudinal spots, one placed on the basal third of the wing at costa and frequently reaching along the costa to the triangular spot; one near the interior margin, a little nearer to the base of wing than the last, and one on the outer third of the interior margin. Two light-colored transverse lines across the end of wing, one very near and parallel with posterior margin, the other bordering the triangular spot behind, and curving across the lower lobe towards posterior angle. The space between these two light lines usually darker than the ground-color. Fringes dark with a light margin. Hind wings trifid, the upper cleft reaching a little beyond the middle, the lower one to the base of wing. Color ashy-brown, the lower lobe produced into a dark angular spot about their middle posteriorly. Antennæ, palpi, head, thorax, and body, tawny yellow; legs of the same color with the exception of the tarsi, which are almost white, with alternate dark brown spots, the spines being black, with dusky tips.—[First Rept., pp. 180–181. Fig. 98, and Pl. II, Figs. 13, 14.]

Zeller has since (1872) referred it to the genus *Platyptilia* (Beitr. zur Kenntn. N. A. Nachtfalter, 2nd part, p. 118), and indicates the difference between it and a very closely allied European species, *P. Zetterstedtii*. He very properly, because of the incongruous compound, drops the conventional ending *dactylus* which I used in the original description.

HETEROPTERA.

NYSIUS DESTRUCTOR, N. Sp.—General color grayish-brown; of shape of *N. thymi* Wolff. *Head* either minutely or more coarsely punctate, and more or less distinctly pubescent; the surface usually brown, with a distinct black, longitudinal line each side, broadening on the crown, but generally leaving the orbit of the eyes pale; these lines sometimes more diffuse and occupying the whole surface, except a median brown spot at base of crown, and a narrow, paler spot on the clypeus; ocelli piceous; eyes opaque, either black or slate-color; face sometimes uniformly pubescent and appearing dark grayish-brown; but more generally black each side of rostrum, with a distinct yellowish-brown spot on the cheeks below the eyes; rostrum piceous, paler at base and reaching to hind coxæ; antennæ either pale yellowish-brown or darker brown, the torulus and first joint darkest. *Thorax*, pronotum narrowing anteriorly, the sides slightly sinuate, irregularly and more coarsely punctate than the head, more or less pubescent, dingy yellow or brown, with a transverse black band near the anterior edge, obscuring the incision and leaving the edge pale, especially in the middle, where there is often a conspicuous pale spot; also five more or less distinct longitudinal dark lines, the central one most persistent and leading on the posterior margin to a pale, shiny, impunctate spot; the callus at hind angles, and sometimes an intermediate spot between it and the median one, and the entire posterior margin, also pale and impunctate; scutellum dark, coarsely punctate, sometimes with a smooth median lon-

gitudinal ridge ending in a pale spot, and with the lateral margins pale; prosternum dark, more or less pubescent, the anterior and posterior margins, and a band outside of coxæ, more or less broadly pale; mesosternum and metasternum also dark, with the pale spots outside of coxæ. *Legs* pale yellow, inclining more or less to brown; coxæ dark at base, pale at tip; trochanters pale; front and middle femora spotted more or less confluent on the outside with brown; hind femora, ♂ dark brown, except at tips and base; ♀ spotted only; tibiæ ringed with brown at base; tarsi marked more or less with brown, especially at tip. *Hemelytra* either colorless, transparent and prismatic, or distinctly tinged with dingy yellow; shallowly punctate and very finely pubescent, the veins of corium and clavus dingy yellow, with brown streaks, the more constant of these streaks being two on posterior margin of corium, and one at the tip of clavus. *Abdomen*, ♂ tergum piceous, with the sutures and sides of some of the joints rarely paler; venter piceous, minutely and regularly covered with gray pubescence; ♀ sutures and spots on tergum more often pale; venter dingy yellow, except at base; ♀ paler than ♂, and generally larger. Average length 0.13 inch [= 3.4^{mm}].

Larva.—Dingy yellow, with more or less distinct longitudinal dark lines, especially on head.

Pupa.—Same color, with more distinct red and brown longitudinal lines, and two little tooth-like, pale yellow processes at inner base of hemelytra pads, indicating the wings; the abdomen paler than the rest of the body.

Described from numerous specimens. I have some, especially males, in which the black so predominates that the paler parts of the head and thorax are scarcely traceable, while in others again the pale parts predominate almost to the exclusion of the black. Indeed, so variable is the species that it is difficult to see wherein some of the specimens differ from the European *thymi*, or from *N. angustatus* Uhler, and it is barely possible that future comparison will show specific identity between some or all of the three. But as long as authors fail to give the variation a species is liable to, or the number of specimens a description is drawn up from, it will remain impossible to decide such questions satisfactorily, and I name *destructor* at the suggestion of our Hemipterist, Mr. P. R. Uhler, of Baltimore, who has examined specimens which I sent him.—[Fifth Rept., p. 113. Fig. 41.

MYTILASPIS POMICORTICIS, N. SP.—*Eggs*—from 30 to 100 under each scale; length scarcely 0.01 inch, irregularly ovoid, nearly thrice as long as wide, snow-white, except just prior to hatching, when they become yellowish. *Larva*—Length of body 0.01 inch, ovoid, thrice as long as wide, pale yellow, with a darker yellow spot near each end; a few short hairs seen around border; two fine anal setæ about half as long as body springing from two lobes between which two spinous hairs are always seen; antennæ quite variable, the joints irregular and not easily resolved, sometimes appearing only 6-jointed, but more generally 7-jointed, with a few hairs, two or three at tip the longest and most persistent; legs with a one-jointed tarsus, a feeble claw, and, among other hairs, four more or less distinctly knobbed ones near tip, the two uppermost longest.

♂—Length of body, 0.022 inch [= .5.5^{mm}]; color, translucent carneous-gray; a dorsal transverse band on each abdominal joint, and portions of the mesothorax and metathorax darker, or purple-gray; the members somewhat lighter. *Head*, sub-triangular; rostrum rudimentary; ocular tubercles, one each side of it, plainly visible, the eyes on the upper surface prominent, dark, and with few facets; antennæ as long as body, 10-jointed, joints 1 and 2 bulbous and sometimes indistinctly separated; 3—9 about four times as long as wide, slightly constricted; 10 half as long and fusiform; all but basal two with a whorl of about eight hairs, slightly clavate and as long as width of joint. *Thorax* very large, oval; prothoracic portion narrowing in front, composed of two transverse folds, the anterior one having a transverse row of four dusky dots; the mesothoracic portion large and elevated, showing three lateral swellings; a well-defined medio-dorsal plate, rounded in front, shallowly-notched behind, with a medio-

longitudinal suture, and a transverse one dividing it in two, the anterior half pale, the posterior darker; the metathoracic portion showing a sub-triangular scutel, and separated from mesothorax by the transverse band (*apodema* of Targioni). *Wings* about as long as body, arising from base of mesothorax, spatulate, closing flat on back in repose, and appearing whitish, finely and uniformly covered with short, stiff hairs; supported by a bifurcate vein, the bifurcation arising from basal fourth, and each fork running near and almost parallel with the wing-margins; balancers dark, with the hook quite long. *Legs* with the middle pair longest, and—from large size of coxæ—further from front than from hind pair; the coxæ and femora large and swollen, the latter with a more or less distinct lobe near the base below; the tarsi one-jointed, with a constriction occasionally indicated, and terminating in a single flexible claw, surrounded by four clubbed hairs; the tibiæ and tarsi are quite bristly, but on the femora there are usually but two bristles, one about the middle above, and one on the basal lobe below; the coxæ also have one above. *Abdomen*, seen from above, nearly as long as thorax; appearing shorter from below; 8 joints only discerned; the last joint abruptly narrowed into a large tubercle bearing four bristles on the under side, and sending forth the genital armor in the form of an awl-shaped style as long as the abdomen.

♂ *Scale*—Larval part golden yellow; the anal shield yellowish-brown, sometimes quite pale, inclining to white, flattened, straight, rather more than twice the length of larval scale, increasing in width from tip to end, where it is slightly truncate; attached by a white film; average length, 0.035 inch.

♀—Average length, 0.05 inch; color, pale yellow; jug-shaped and flattened when young, more globular when mature, and twice as long as wide; the cephalo-thoracic portion rounded and entire, but narrower than the abdominal, at the juncture with which it forms a more or less conspicuous lateral projection; on its inferior side is a tubercle, having two longitudinal ridges, and giving rise to a corneous, filiform proboscis, longer than the body, and composed of four separate parts; posterior abdominal joints deeply lobed laterally, with two or three blunt, fleshy hairs to each lobe; anal plate gamboge-yellow, corneous, with an irregular border, presenting two larger, slightly tri-lobed, median projections, and one or more smaller ones each side, furnished with spinous hairs, two especially between the tri-lobed projections aforementioned; five more or less complete sets of secretors visible from below, arranged around anus in form of an arc, the median set with normally 10, the upper laterals 20, and the lower laterals 14; besides these, some six or more blunt tubes, and a series of shorter pointed ones, may be noticed along the border, and doubtless serve as secretors. (See Fig. 32 *b*.)

♀ *Scale*—Larval scale golden-yellow; median scale somewhat darker; anal shield varying from pale brown to deep purplish-gray, and generally of a color with the bark it is upon. The whole scale is often incanous, but the hoary film easily rubs off; it averages 0.12 inch in length, but is quite variable in form and size, being either straight or curved, narrow and strongly arched, or broad and flatter, but always rounded at the end; the white inferior laminae at sides sometimes show distinctly from above, and give the appearance of a pale border.

The lice, whether ♂ or ♀, vary in appearance according to position and state of maturity. In making the foregoing descriptions and figures, I have taken what appeared the most natural positions, after examination of many specimens. The ♂ abdomen shrinks very much in drying, and the more detailed ♀ characters are variable. While the normal number of secretors in the middle set is never more than 10, I have sometimes found but 8 or 9; that of the upper laterals never surpasses 20, but may be as low as 15; while that of the lower laterals is more uniformly 14, though I have sometimes found 16, and at others 12. Opposite sets do not always contain the same number.—[Fifth Rept., pp. 95–96. Figs. 31, 32.]

This is the species previously known as *Aspidiotus conchiformis*, or popularly as the Oyster-shell Bark-louse, and the reasons for separating it are given in the report.

ERIOSOMA ULMI, N. Sp.—Color dark blue. Length to tip of closed wings, exclusive of antennæ, 0.12 [inch, = 3^{mm}]. Wings hyaline, three times as long as wide, and more pointed at the ends than in *E. pyri*. Costal and subcostal veins, and that bounding the stigma behind, robust and black. Discoidal veins together with the 3d forked and stigmal veins, all slender and black, the forked vein being as distinct to its base as are the others, with the fork but $\frac{1}{3}$ as long as the vein itself and curved in an opposite direction to the stigmal vein. Antennæ 6-jointed and of the same color as the body; joints 1, 2, 4, 5 and 6 of about equal length, joint 3 thrice as long as either. Legs of the same color as body.

The young lice are narrower and usually lighter colored than the mature individuals, varying from flesh or pink to various shades of blue and purple.—[First Rept., p. 124.]

Professor Thomas (Trans. Ill. St. Hort. Soc., 1876, p. 191) has called it *Erisoma Rileyi** because of *ulmi* being preoccupied by an European species. It belongs to *Schizoneura*. For subsequent remarks see "Notes on the Aphididæ of the United States, etc., by C. V. Riley & J. Monell," (Bull. Hayden's U. S. Geol. & Geogr. Survey, Vol. V, No. 1, p. 3.)

DIPTERA.

ASILUS MISSOURIENSIS N. Sp.—Alar expanse 1.85 [inches, = 47^{mm}]; length of body 1.30 inches [= 33^{mm}]. Wings transparent, with a smoky yellow tinge, more distinct around the veins, which are brown. Head pale yellow, sometimes brownish; moustache straw-yellow with a few stiff black hairs below; beard pale straw-yellow; crown very deeply excavated; base of the same pale yellow with short, stiff, yellowish hairs, and a crown of black ones near the border; eyes large, prominent, finely reticulated and almost black; antennæ, first joint black tipped with brown, cylindrical and hairy; second joint black, short, thick and rounded at tip, with a few stiff hairs; third joint as long as first, tapering each way, smooth, black and terminating in a long, brown bristle; proboscis black and nearly as long as face; neck with pale and black hairs. Thorax leaden-black, slightly opalescent with reddish brown at sides, more or less pubescent with pale yellow, especially laterally and posteriorly and in three narrow longitudinal dorsal lines which gradually approach towards metathorax; bearded at sides and behind with a few decurved black bristles, those behind interspersed with a few smaller pale hairs; scutel of the same color, with upward-curving, black bristles; halteres brown. Abdomen, ♂, general color dull leaden-yellow, with darker transverse bands at intersections; the light color produced by a yellowish pubescence and numerous short close-lying yellow hairs, the dark bands produced by the absence of this covering at the borders of each segment; basal segment broad, bilobed, and with lateral black bristles; segments 6, 7, 8 and anal valves with a decided pink tint, especially 7; 8 but one-third as long as 7 above. ♀, broader, flatter, more polished and brassy, with no transverse darker bands, segments 7 and 8 polished black, the latter narrow and longer than any of the others; anus with a few black bristles. Legs, dull purple-brown, with black bristles; thighs very stout, the hind pair rather darker than the others, the two front pair of trochanters with long, yellowish hairs; pulvilli, generally fulvous.

Described from two ♂, and two ♀, all captured while sucking honey-bees. I have not access to Loew's descriptions, and cannot therefore compare it with already described species; but specimens have been sent to Dr. Wm. LeBaron, of Geneva, Illinois, and to Baron Osten Sacken, of New York, and both these gentlemen are unacquainted with it, and believe it to be new. In the well marked ♂ specimens, the body bears a general resemblance to that of *Trupanea* [*Promachus*] *vertebrata*, Say.—[Second Rept., pp. 122-123. Fig. 89.]

* By typographical error *Rilepi*.

Baron Osten Sacken has since placed this as a synonym of *Proctacanthus Milbertii* Macq. in the second edition of his Catalogue of the described Diptera of North America (1878), p. 81.

LYDELLA DORYPHORÆ, New Species.—Length 0.25 [=6mm]. Alar expanse 0.48 [=12mm]. Antennæ black. Palpi fulvous. Face silvery white. Front silvery, tinted with pale golden-brown, with a broad middle stripe black. Thorax cinereous with imperfect black stripes. Abdomen black and silvery-ash, changing into each other when viewed from different angles. When viewed from above: first segment deep black with a posterior border of silver-ash very narrow in the middle, much widened laterally, but abbreviated at the sides of the abdomen. The other segments with the basal half silvery-ash, terminal half black. Legs black. Fourth longitudinal vein of the wings straight after the angle. Posterior transverse vein arcuate.

Described from numerous bred specimens.—[First Rept., pp. 111-112. Fig. 48.]

This species is referred by Ostén Sacken to the genus *Exorista* of Schiner, *Lydella* not being received as a distinct genus. The name *Lydella* is used also for a genus of Acarina.

EXORISTA FLAVICAUDA, N. Sp.—Length 0.35 to 0.50 inch [=8.5-12.5mm]. Head broader than thorax; face, silvery-white, the cheeks inclining to yellow, with lateral black hairs extending to near the base of antennæ, and one stiffer and longer bristle at top of cheeks; front, dusky, ferruginous, with two rows of black converging bristles; divided by a broad depressed stripe of a brighter ferruginous color and without bristles; occiput bright ferruginous; labium ferruginous with hairs of same color; maxipalps rufous; eyes dark mahogany-brown, and perfectly smooth; antennæ, two basal joints rufous, with black hairs, third joint flattened, dusky, and thrice as long as second; seta, black; entire hinder part of head covered with dense white hairs. Thorax, more decidedly blue than in *leucaniæ*, broader (instead of narrower) in front than behind; the vittæ less distinct; scutellum of same color as thorax. Abdomen, stout and more cylindrical than in *leucaniæ*; first joint dark bluish-gray; second, light bluish-gray, becoming darker along the middle, at sides and at lower border; third joint, like second above, but golden-gray at sides (no rufous); last joint entirely yellow or pale orange, with no other color and but few black bristles around anus. Wings more dusky than in *leucaniæ*; alulae, opaque bluish-white. Legs, black; pulvilli pale yellow.

Described from one captured, 4 bred ♀. Space between eyes at occiput fully one-third the width of head.—[Second Rept., pp. 51-52. Fig. 18.]

TACHINA [EXORISTA] PHYCITÆ, LeBaron—*Imago*.—Length, 0.20 inch [=5mm]. Antennæ black, third joint twice as long as the second; face silvery, without bristles at the sides; sides of the front silvery at the lower part, pale golden above; the middle black vitta occupying a little more than half of the width of the inter-ocular space; frontal bristles continued down the face to opposite the end of the second joint of antennæ; palpi blackish-brown; eyes hairy. Thorax black, with the ordinary cinereous stripes scarcely perceptible. Abdomen black, varied with cinereous at the base of the segments; a large fulvous spot on the side of the abdomen occupying nearly the whole of the side of the second segment, half or more of the third, and sometimes a small spot on the first; bristles on the middle as well as at the hind-margin of the second and third segments. Venation of the wings of the usual type; first posterior cell almost closed, before the end of wing; fourth long vein slightly curved after the angle; fifth long vein prolonged to the margin; hind cross vein moderately sinuous. Tarsal claws and pulvilli unusually long.

Female? A single specimen, a very little larger than the others, was obtained from the same lot of leaf-crumplers, which possibly may be the ♀ of the same species. It differs as follows: Front broader; antennæ dark brown; the cinereous markings of the body more distinct; the tip of abdomen fulvous, but without the fulvous spot at the sides; and with the tarsal claws of ordinary length.

This species appears to belong to the subgenus *Exorista* of Meigen, closely allied to *Tachina* proper, and differing from it chiefly in having the eyes hairy, and in the presence of bristles on the middle, as well as at the hind margin of the second and third abdominal segments, whereas *Tachina* has only the latter.—[Fourth Rept., p. 40-41.]

This species was simultaneously published by Dr. LeBaron in his 2d Rept. Ins. Ill., p. 123. It is retained in *Exorista* by Osten Sacken.

ANTHOMYIA ZEAE ♀, N. Sp. (Pl. 2, Fig. 24). Length 0.20 [inch, = 5^{mm}]; alar expanse 0.38 [inch, = 9.5^{mm}]. Antennæ black; style microscopically pubescent; front, fulvous, with a distinct, rather narrow, brownish, cinereous margin; face and orbits brownish-white; palpi and proboscis black; ocellar area somewhat heart-shaped; thorax and abdomen pale yellow-brownish cinereous, with minute black points at the insertion of the bristles; thorax with an indistinct middle stripe of brown; legs black, tinted with cinereous; poisers pale ochre-yellow; scales small, the upper valve larger than the lower.—[First Rept., p. 155. Figs. 86, 87, and Pl. II, Fig. 24.]

ANTHOMYIA RADICUM (Linn.) var. CALOPTENI—Egg—Oval, smooth, white, 0.04 inch long.

Larva—Skin unarmed, 0.24 inch [= 6^{mm}] long when extended, of the normal form, the mandibular hooks black, quite conspicuous, and diverging at base. Prothoracic spiracles elongate. Anal spiracles minute, yellowish-brown, with the 8 fleshy surrounding tubercles, small.

Pupa—Pale-brown, rounded at each end, with the prothoracic spiracles and lips anteriorly, and the anal spiracles and lower tubercles posteriorly, showing as minute points.

Imago—♀. Average expanse 0.48 inch [= 12^{mm}]. General color ash-gray with a ferruginous hue, especially above, and a more or less intense metallic reflection. Face with white reflections below; eyes smooth, brown, encircled by the ground color, and this behind and on forehead bordered by a brown line; 2 similar lines at back of head from upper corners of eyes and approaching to neck; forehead dusky-brown, becoming bright yellowish-red toward base of antennæ, and the brown forking at right angles around occiput. Trophi and antennæ black, the style simple and somewhat longer than the whole antennæ. Thorax with three dusky longitudinal lines, obsolete behind; legs black, with cinereous hue beneath; wings faintly smoky, with brown-black veins, the discal cross-vein straight and transverse, the outer one bent and more oblique; balancers crumpled, yellowish. Abdomen with faint dusty medio-dorsal spots, broad at base, tapering and obsolescing toward end of each joint.

In the ♂, aside from the larger eyes, stronger bristles, and narrower, less tapering abdomen with its additional joint—all characteristic of the sex—the face is whiter, and the medio-dorsal dark mark of abdomen continuous.

Described from 25 specimens of both sexes, reared from locust-egg-feeding larvæ.

Specimens bred from cabbage and radish roots, and others in my cabinet taken from the burrows (made in Osage Orange in Missouri) of *Crabro stirpicola* Pack.; do not differ specifically.—[Ninth Rept., p. 95.]

For further details see First Rept. of the Commission (pp. 285-9), where the species is shown to be the *Anthomyia angustifrons* of Meigen.

ORTHOPTERA.

CALOPTENUS ATLANS N. sp.—Length to tip of abdomen 0.70-0.85 inch [= 17.5-21^{mm}]; to tip of closed wings 0.92-1.05 inches [= 23-26^{mm}]. At once distinguished from *femur-rubrum* by the notched character of the anal abdominal joint in the male and by the shorter, less tapering cerci; also by the greater relative length of wings which extend, on an average, nearly one-third their length beyond the tip of the abdomen in the dried specimens; also by the larger and more distinct spot on the wings—in all which characters it much more closely resembles *spretus* than *femur-rubrum*.

From *spretus*, again, it is at once distinguished by the smaller size, the more distinct separation of the dark mark running from the eyes on the prothorax and of the pale line from base of wings to hind thigh; also by the anal joint in the ♂, tapering more suddenly and by the two lobes forming the notch being less marked. From both species it is distinguished not only by its smaller size but by the deeper, more livid color of the dark parts, and the paler yellow of the light parts—the colors thus more strongly contrasting.

6 ♂'s, ♀ 7's from New Hampshire. Just as the typical *femur-rubrum* is at once distinguished from the typical *spretus* by the characters indicated; so *Atlantis*, though structurally nearer to *spretus*, is distinguished from it at a glance by its much smaller size and darker, more marbled coloring. The contrast is all the greater in the living specimens, and I have seen no specimens of *spretus* that at all approach it in these respects.

Whether this is the *femur-rubrum* as defined by DeGeer or by Harris, it is almost impossible to decide, though Harris's figure of *femur-rubrum* better represents it than the true *femur-rubrum*, as subsequently defined by Thomas, and as found in Illinois and Missouri.—[Seventh Rept., pp. 169–170.]

For further details and structural differences between it and *C. spretus* see First Report of the Commission.

LIST OF DESCRIPTIONS OF ADOLESCENT STATES.

In making out the following list of descriptions of adolescent states, etc., that appeared in the Reports, the nomenclature there used is retained. Unless otherwise stated the insects, in the particular states indicated, were at the time unknown or undescribed, the descriptions first appearing in the Reports. Those published in connection with the preceding descriptions of new species are omitted here:

HYMENOPTERA.

- Nematus ventricosus*; larva: IX, 21. (Previously described by several writers.)
Pristiphora grossulariæ; larva: IX, 26. (Description quoted from Walsh.)
Emphytus maculatus; larva and pupa: IX, 23-29. (Previously described by me in the *Prairie Farmer*, May 25, 1867.)
Lophyrus abbotii; larva: IX, 32.
Lophyrus lecontei; larva: IX, 33. (This and *abbotii* both partially described by me in the *Prairie Farmer*, November 10, 1866; May 25, 1867; May 2, 1868, and in the *Prairie Farmer Annual*, 1869.)
Tiphia inornata; larva: VI, 126.

COLEOPTERA.

- Harpalus* (probably *herbivagus* Say); larva: IX, 97.
Harpalid; larva: I, 59.
Mysia 15-punctata; larva: IV, 18.
Chilocorus bivulnerus; larva and pupa: I, 16.
Hippodamia convergens; larva and pupa: I, 112. (Previously mentioned in the *Am. Ent.* I, 46, and elsewhere.)
Coccinella picta; larva: V, 101.
Passalus cornutus; larva and pupa: IV, 140-141. (Previously mentioned by Burmeister and by Walsh.); egg: V, 55.
LACHNOSTERNA QUERCINA; egg: V, 55.
Pelidnota punctata; larva and pupa: III, 78-79. (First described by me in *Am. Ent.* II, 295.)
Telephorus bilineatus; larva: IV, 30. (First described by Packard.)
Chauliognathus pensylvanicus; larva: I, 57. (Quoted from the *Am. Ent.* I, 35.)
Chrysobothris femorata; eggs: VII, 73; larva, I, 46. (Previously described by Fitch and others); eggs, larva, and pupa: VII, 73.
Sinoxylon basilare; larva and pupa: IV, 54.
Corynetes rufipes; larva and pupa: VI, 101, 102.
Prionus laticollis; larva: I, 126; larva and pupa: II, 87; egg: V, 56. (*laticollis*.)
Saperda bivittata; pupa: I, 43. (Previously described by Harris.)
Lema trilineata; larva and pupa: I, 99. (From the *Prairie Farmer*; and the *Am. Ent.* I, 26. Previously described by Harris and others.)
Doryphora juncta; larva: I, 106. (First described in the *Am. Ent.* I, 43.)
Doryphora 10-lineata; eggs and larva: I, 105. (From the *Am. Ent.* I, 43. Previously described by me in *Prairie Farmer* Aug. 8, 1863.)

- Colaspis flavida*; larva: III, 84, and IV, 34.
Coscinoptera dominicana; eggs and larva: VI, 128, 130.
Haltica chalybea; larva and pupa: III, 81. (Quoted from *Am. Ent.* II, 327. The larva first described by Packard, *Guide*, p. 507.)
Blepharida rhois; egg, larva and pupa: VI, 121.
Cassida bivittata; larva and pupa: II, 61. (First described by me in the *Prairie Farmer Annual* for 1868, p. 53.)
Cassida aurichalcea; egg: II, 60; larva and pupa: II, 62. (Previously described by Harris.)
Cassida pallida; larva: II, 62.
Cassida guttata; larva and pupa: II, 63.
Cassida nigripes; larva and pupa: II, 63, 64.
Bruchus pisi; egg: III, 47.
Tenebrionid?; larva: VI, 113. (Previously described as the larva of *Eupsalis* by Harris.)
Eupsalis minuta; larva and pupa: VI, 115, 116. (The pupa first described by Harris.)
Conotrachelus cratægi; larva and pupa: III, 39.
Baridius trinotatus; larva and pupa: I, 95. (From the *Am. Ent.* I, 22.)
Anthonomus quadrigibbus; egg: III, 31; larva and pupa: III, 35.

LEPIDOPTERA.

- Papilio philenor*; larva and pupa: II, 117. (Previously described by Smith and Abbot, and by Boisduval and Le Conte; also by Harris in *Ent. Corr.*)
Pieris protodice; larva and pupa: II, 104. (Published simultaneously in the *Am. Ent.* II, 77.)
Pieris rapæ; larva and pupa: II, 108. (Previously described by various authors.)
Danaïs archippus; egg: III, 144.
Limenitis disippus; egg and larva: III, 154. (The mature larva previously described by various authors.)
Apatura lycaon; egg, larva and pupa: VI, 146, 147. (The larva and pupa badly described by Boisd. & Lec.)
Apatura herse; egg, larva and pupa: VI, 148. (The larva and pupa badly described by Boisd. & Lec.)
Paphia glycerium; larva and pupa: II, 127. (First published by me in *Am. Ent.* II, 123); egg and larval changes: V, 146.
Megathymus yuccæ; egg, larva and larval changes: VIII, 174, 181. (First published by me in *Trans. St. Louis Ac.*); IX, 129.
Chærocampa pampinatrix; egg, larva and pupa: II, 71, 72. (Previously*described, except egg, by various authors.)
Philampelus achemon; young and full grown larvæ and pupa: II, 74, 75. (Previously described by various authors.)
Philampelus satellitia; eggs, young and full grown larvæ, and pupa: II, 76-78. (Previously described, except egg, by various authors.)
Sphinx 5-maculata; larva pupa: I, 95. (From the *Am. Ent.* I, 23; previously described by several authors.)
Thyreus Abbotii; larva and pupa: II, 78, 79. (Previously described by various authors.)
Deilephila lineata; two forms of larva: III, 141, 142. (Previously described, but not in connection. Quoted from the *Am. Ent.*, II, 258.)
Ægeria acerni; larva and pupa: VI, 110.
Ægeria rubi; larva: VI, 113.
Psychomorpha epimenis; larva and pupa: III, 64, 65; VI, 88. (First described as the possible larva and pupa of *End. unio*, *Am. Ent.* II, 152 and in 1st Rept., p. 84.)
Eudryas grata; eggs, larva and pupa: II, 83; VI, 89, 90. (The larva previously described by Harris and others.)

- Eudryas unio*; larva and pupa: VI, 92. (First described by Lintner.)
- Alypia octomaculata*; larva: I, 136, (previously mentioned by Fitch); II, 80, published simultaneously in the *Am. Ent.*, II, 151, (previously described in *Harris' Corr.*); VI, 94.
- Procris americana*; larva and pupa: II, 86. (First described by Harris.)
- Callimorpha fulvicosta*; larva: III, 134.
- Spilosoma virginica*; larva and pupa: III, 69. (Previously described by various authors.)
- Hyphantria textor*; larva: III, 132. (First described by Harris.)
- Ecpantheria scribonia*; larva: IV, 143. (Previously described by other authors.)
- Bombyx mori*; egg and larva: IV, 86. (Previously well known.)
- Attacus cecropia*; larval changes: IV, 106. (Quoted from the *Am. Ent.*, II, 100.)
- Attacus cynthia*; larval changes: IV, 117. (Previously described by other authors.)
- Attacus promethea*; larval changes: IV, 121. (Partially given by other authors previously.)
- Attacus luna*; larval changes: IV, 124. (Previously given by Lintner.)
- Attacus polyphemus*; larval changes: IV, 126.
- Attacus yama-mai*; larval changes: IV, 132. (Previously described by other authors.)
- Attacus pernyi*; egg, larva, and cocoon: IV, 137. (Previously described by other authors.)
- Hemileuca maia*; egg and larval changes: V, 128, 129. (Previously described by Lintner.)
- Hyperchiria io*; larval changes: V, 135. (Previously given by Lintner.)
- Anisota rubicunda*; eggs and larval changes: V, 138.
- Acronycta oblonga*; larva and pupa: III, 71. (The larva first figured by Smith & Abb.)
- Acronycta xylinoides*; larva: V, 126.
- Amphipyra pyramidoides*; larva and pupa: III, 73, 74.
- Leucania unipuncta*; larva and pupa: II, 49; VIII, 33, and larva: II, 55 (previously described by various authors); egg: VIII, 34; egg and larval changes: VIII, 184, 185.
- Gortyna nitela*; larva: I, 92. (From the *Am. Ent.*, II, 22. Briefly described by Harris, *Treatise*, p. 440; but first identified by me in the *Prairie Farmer*.)
- Agrotis inermis*; larva and pupa: I, 74.
- Agrotis cochrani*; larva and pupa: I, 76. (First described by me in the *Prairie Farmer*, June 22, 1867.)
- Agrotis clandestina*; larva and pupa: I, 79. (Previously mentioned by Harris.)
- Agrotis telifera*; larva and pupa: I, 81. (Described by me in the *Prairie Farmer*, June 22, 1867; and previously described in Europe, where the species also occurs and is known as *A. ypsilon*.)
- Agrotis subgothica*; larva: I, 82.
- Agrotis jaculifera*; larva and pupa: I, 83.
- Agrotis devastator*; larva and pupa: I, 84.
- Hadena subjuncta*; larva and pupa: I, 85.
- Celæna renigera*; larva and pupa: I, 86.
- Prodenia commelinæ*; larva: I, 88; III, 114 (from *Am. Ent.*, II, 363). [See Notes.]
- Anisopteryx vernata*; larva and pupa: II, 95-97 (previously described by other authors); eggs, larva and pupa: VII, 82 (and 86-87, adapted from Mann); *Paleacrita vernata*, VIII, 13-17 (from the *Trans. St. Louis Acad.*)
- Anisopteryx pomataria*; eggs: II, 94-95 (the two species confounded); eggs, larva and pupa: VII, 84 (and 86-87, adapted from Mann); VIII, 13-17 (from the *Trans. St. Louis Acad.*)
- Eufitchia ribearia*; egg, larva and pupa: IX, 3, 4. (The larva first described by Fitch.)
- Phacellura nitidalis*; larva: II, 67.
- Asopia costalis*; larva and pupa: VI, 106. (The larva mentioned by Harris, but first described by Walsh in the *Prac. Ent.*, and first bred and determined by me, *Prairie Farmer*, April 20, 1867.)

- Phycita nebulo*; *larva* and *pupa*: IV, 41. (The larva first described by LeBaron.)
Pempelia grossulariæ; *larva* and *pupa*: I, 141. (Larva previously described by Fitch and by Packard.)
Tortrix rileyana; *larva* and *pupa*: I, 154.
Anchylopera fragariæ; *larva*: I, 143. (First described in the *Am. Ent.*, I, 90.)
Penthina vitivorana; *larva* and *pupa*: I, 135. (The larva first described, but not identified, by Rathvon.)
Carpocapsa pomonella; *larva* and *pupa*: I, 63. (Previously described by various authors.)
Walshia amorphella; *larva* and *pupa*: II, 133.
Bucculatrix pomifoliella; *larva* and *pupa*: IV, 51. (Larva previously described by Clemens.)
Æta compta; *larva* and *pupa*: I, 152.
Pterophorus periscelidactylus; *larva* and *pupa*: I, 137; III, 66. (Previously described by Fitch.)
Pterophorus carduidactylus; *larva* and *pupa*: I, 180.
Pronuba yuccasella; *larva*: V, 155; *pupa*, VI, 131 (from Trans. St. Louis Acad.); *egg*, VI, 133 (from *Am. Nat.*).
Orgyia leucostigma; *eggs*, *larva* and *pupa*: I, 144-146. (Previously described by others.)
Thyridopteryx ephemeræformis; *eggs*, *larva* and *pupa*: I, 148, 149. (Previously described by others.)
Hæmatopis grataria; *eggs*, *larva* and *pupa*: I, 179.
Galleria cereana; *larva* and *pupa*: I, 166. (Previously described by other authors.)

HEMIPTERA.

- Strachia histrionica*; *eggs*, *larva* and *pupa*: IV, 37.
Micropus leucopterus; *egg*, *larval stages* and *pupa*: VII, 21.
Cicada septemdecim; *egg* and *young larva*: I, 25. (The eggs previously described by several writers.)
Pæcilopectera pruinosa; *eggs*: V, 122.
Ceresa bubalus; *eggs*: V, 121.
Mytilaspis pinifoliæ; *eggs* and *larva*: V, 98. (First mentioned by LeBaron.)
Phylloxera rileyi; *larva* and *pupa*: VI, 64, 86; VII, 120.
Phylloxera vastatrix; *various forms*: VI, 66 (previously described elsewhere and by others); *impregnated egg*: VIII, 159. (Previously described by me in the Trans. St. Louis Acad. for Oct. 18, 1875, and independently by Balbiani in the Comptes rendus de l'Ac. d. Sc. Paris for Oct. 4, 1875.)
Eriosoma pyri; *larva*: I, 120. (From the *Am. Ent.*, I, 82; previously described by several authors.)

DIPTERA.

- Tabanus atratus*; *larva* and *pupa*: II, 130, 131. (Previously described, but not specifically identified, by Walsh.)
Erax bastardi; *larva* and *pupa*: II, 124.
Bombyliid; *larva*: IX, 96.
Pipiza radicum; *larva* and *pupa*: I, 122. (Quoted from the *Am. Ent.*, I, 84.)
Anthomyia zææ; *larva* and *puparium*: I, 155.
Meromyza americana; *larva* and *pupa*: I, 160.
Æstrus ovis; *larva* and *puparium*: I, 162. (Previously described by other authors.)

ORTHOPTERA.

- Mantis carolina*; *eggs* and *larva*: I, 170-171. (Previously described by several authors.)
Æcanthus niveus; *eggs*: V, 120. (Previously described as eggs of *Ceresa bubulus* by Fitch.)

Orchelimum glaberimum; *eggs*: V, 123.

Phaneroptera curvicauda; *eggs*: V, 124, and VI, 165; *larva* and *pupa*: VI, 166.

Microcentrus retinervis; *eggs*: V, 123; VI, 155 (previously described as eggs of *Platyphyllum* by Harris); *larva* and *pupa*: VI, 161.

Phylloptera oblongifolia; *eggs*: V, 123. (See *Microcentrus*.)

Platyphyllum concavum; *eggs*: V, 124; VI, 167.

Caloptenus spretus; *eggs* and *egg-mass*: IX, 88, 89; *larva* and *pupa*: VII, 129.

NEUROPTERA.

Corydalis cornutus; *larva* and *pupa*: V, 143, 144 (Previously described by Haldeman);
eggs and *egg-mass*, and *young larva*: IX, 127.

LIST OF DESCRIPTIONS, MOSTLY AM- PLIFIED, OF SPECIES NOT NEW.

The following list includes the species, already known, of which a complete redescription of the adult is given in the Reports, either because the original description was in a foreign language, or not easily accessible, or of one sex only, or for other reasons.

HYMENOPTERA.

- Tiphia inornata* Say : VI, 126.
Cryptus extrematis Cress. : IV, 111.
Pezomachus minimus Walsh : II, 52. (From Walsh.)
Ophion purgatus Say : II, 53.
Mesochorus vitreus Walsh : II, 52. (From Walsh.)
Pimpla annulipes Brullé : V, 49.
Macrocentrus delicatus Cress. : V, 50.
Microgaster militaris Walsh : II, 52. (From Walsh.)
Chalcis mariæ Riley : IV, 110. (From the *Am. Ent.*, II, 101-102.)
Isosoma vitis Saunders : II, 93. (From Saunders.)
Antigaster mirabilis Walsh : VI, 163. (From the *Am. Ent.*, II, 169-170.)
Pristiphora grossulariæ Walsh : IX, 26-27. (From the *Prac. Ent.*, I, 123.)
Nematus ventricosus (Klug) : IX, 22. (From the *Prac. Ent.*, I, 120-121, and the *Am. Ent.*, II, 16-17.)
Emphytus maculatus Nort. : IX, 28.
Lophyrus LeContei Fitch : IX, 33.

COLEOPTERA.

- Doryphora 10-lineata* Say, var. : IX, 40.
Sphenophorus zeæ Walsh : III, 59. (From Walsh.)
Scolytus caryæ Riley : V, 107. (Female first described in *Prairie Farmer* Feb. 2, 1867.)
 [See Notes.]

LEPIDOPTERA.

- Apatura lycaon* (Fabr.) : VI, 144.
Apatura herse (Fabr.) : VI, 144.
Megathymus yuccæ (Walk.) : VIII, 175-176.
Ægeria polistiformis Harr. : III, 76.
Ægeria acerni Clem. : VI, 110.
Prodenia autumnalis Riley : III, 116-117. (From *Am. Ent.*, II, 365.) [See Notes.]
Leucania unipuncta Haw. : II, 56.
Leucania albilinea Guen. : IX, 56-57.
Acronycta obliterata Sm. & Abb. : III, 71.
Amphipyra pyramidoides Guen. : III, 74.
Celæna renigera Steph. : I, 86.
Hadena subjuncta Gr. & Rob. : I, 85.
Noctua clandestina Harr. : I, 79.

- Agrotis inermis* *Harr.* : I, 74.
Agrotis cochrani *Riley* : I, 75.
Agrotis telifera *Harr.* : I, 81.
Agrotis jaculifera *Guen.* : I, 83.
Anisopteryx pometaria *Harr.* : VIII, 15-17. (From the *Trans. St. Louis Acad. Sc.*)
Paleacrita vernata (Peck) : VIII, 15-17. (From the *Trans. St. Louis Acad. Sc.*)
Asopia costalis (*Fab.*) : VI, 107.
Pempelia grossulariæ (*Pack.*) : I, 141.
Walshia amorphella *Clem.* : II, 133.
Penthina vitivorana *Pack.* : I, 135.
Euryptychia sa igneana *Clem.* : II, 134. (From *Clemens.*)
Tortrix rileyana *Grote* : I, 154.
Walshia amorphella *Clem.* : II, 133.
Holcocera glandulella *Riley* : IV, 145. (From the *Can. Ent.*, IV, 13-19.)
Pronuba yuccasella *Riley* : V, 150, 151, 155; VI, 131-132. (Both from the *Trans. St. Louis Acad. Sc.*)
Ceta compta *Clem.* : I, 153.

HEMIPTERA.

- Micropus leucopterus* (*Say*) : VII, 21, 22.
Mytilaspis pinifoliæ (*Fitch*) : V, 99.
Eriosoma pyri (*Fitch*) : I, 120.
Phylloxera vastatrix *Planchon* : VIII, 159 (From *Trans. St. Louis Acad. Sc.*); VI, 66-67; VII, 93, 99.
Phylloxera *Rileyi* *Licht.* : IV, 66; VI, 64, 86; VII, 118-120.
Phylloxera caryæ-gummosa *Riley* : VII, 118. (From the *Comptes Rendus*, Paris Acad. of Sci., Dec. 14, 1874.)
Phylloxera caryæ-ren *Riley* : VII, 118. (From the *Comptes Rendus*, Paris Acad. of Sci., Dec. 14, 1874.)
Phylloxera caryæ-fallax *Riley* : VII, 118. (From the *Comptes Rendus*, Paris Acad. of Sci., Dec. 14, 1874.)

DIPTERA.

- Erax bastardi* *Macq.* : II, 124.
Pipiza radicum *Walsh & Riley* : I, 121-122. (From the *Am. Ent.* I, 83-84.)
Exorista leucaniæ *Walsh* : II, 51. (From *Walsh.*)
Tachina bifasciata (*Fabr.*) : V, 140.

ORTHOPTERA.

- Caloptenus femur-rubrum* (*DeG.*) : VII, 126-128.
Caloptenus atlanis *Riley* : VIII, 117.
Caloptenus spretus (*Thos.*) : VII, 128-132; VIII, 117.

ACARINA.

- Hoplophora arctata* *Riley* : VI, 81. (From *Trans. St. Louis Acad.*, III, 216.)
Tyroglyphus phylloxeræ *Riley & Planchon* : VI, 81. (From *Trans. St. Louis Acad.*, III, 215.)

LIST OF ILLUSTRATIONS.

The illustrations in the Reports were prepared at the author's expense, neither the State nor the Board of Agriculture making any provision therefor. The wood-engraving was done for the most part in St. Louis, by either Wm. Macwitz, Emile Lampe, or Wittemberg & Sorber. Some of it was done by Van Ingen & Snyder, of Philadelphia. A few of the later illustrations are by photo-engraving, and Figs. 50-52 of the 8th Report show the first attempt to combine this process with lithography. In the following list, all drawings were made from nature by the author unless otherwise stated, and when the figure is enlarged the natural size, unless otherwise apparent or stated in this list, will be found indicated in hair-line. The nomenclature of the Reports is retained.

REPORT I.

PLATE I. (Drawn by D. Wiest and lithographed by Bowen & Co., Philadelphia.)

- FIG. 1. Unarmed Rustic (*Agrotis inermis* Harr.), moth.
- FIG. 2. Variegated Cut-worm (*Agrotis inermis* Harr.).
- FIG. 3. Variegated Cut-worm (*Agrotis inermis* Harr.), head, enlarged.
- FIG. 4. Variegated Cut-worm (*Agrotis inermis* Harr.), one joint, enlarged.
- FIG. 5. Climbing Cut-worm Moth (*Agrotis scandens* Riley), wings spread.
- FIG. 6. Climbing Cut-worm Moth (*Agrotis scandens* Riley), wings closed.
- FIG. 7. Climbing Cut-worm (*Agrotis scandens* Riley).
- FIG. 8. Lance Rustic (*Agrotis telifera* Harr.), moth.
- FIG. 9. Greasy Cut-worm (*Agrotis telifera* Harr.).
- FIG. 10. Greasy Cut-worm (*Agrotis telifera* Harr.), head, enlarged.
- FIG. 11. Dart-bearing Rustic (*Agrotis jaculifera* Guen.), moth.
- FIG. 12. *Prodenia commelinæ*, Sm. & Abb., one joint of larva enlarged.
- FIG. 13. Clandestine Owlet Moth (*Noctua clandestina* Harr.).
- FIG. 14. Subjoined Hadenia (*Hadenia subjuncta* Gr. & Rob.), moth.
- FIG. 15. Speckled Cut-worm (*Hadenia subjuncta* Gr. & Rob.), head, enlarged.
- FIG. 16. Speckled Cut-worm (*Hadenia subjuncta* Gr. & Rob.), one joint, enlarged.
- FIG. 17. Speckled Cut-worm (*Hadenia subjuncta* Gr. & Rob.), anal joint, enlarged.
- FIG. 18. Eight-spotted Forester (*Alypia octomaculata*, Fabr.).
- FIG. 19. Grape-vine Epimenis (*Psychomorpha epimenis*, Drury), larva. (Mentioned on p. 136, but first named in the 3d Rept., p. 63.)

PLATE II. (Drawn by D. Wiest and lithographed by Bowen & Co., Philadelphia.)

- FIG. 1. Solidago Gall Moth (*Gelechia gallæsolidaginis* Riley), wings expanded.
- FIG. 2. Solidago Gall Moth (*Gelechia gallæsolidaginis* Riley), wings closed.
- FIG. 3. Walnut Tortrix (*Tortrix rileyana* Grote), wings expanded.
- FIG. 4. Walnut Tortrix (*Tortrix rileyana* Grote), wings closed.
- FIG. 5. Solidago Gall Moth (*Gelechia gallæsolidaginis* Riley), larva swollen by the cocoons of the Inflating Chalcis-fly within.
- FIG. 6. Inflating Chalcis-fly, enlarged.
- FIG. 7. *Hemiteles* (?) *cressonii* Riley, enlarged.
- FIG. 8. *Eurytoma bolteri* Riley; male antenna, enlarged.

- FIG. 9. *Eurytoma bolteri* Riley; female, enlarged.
 FIG. 10. Bag of Bag-worm (*Thyridopteryx ephemeraeformis* Steph.), cut to show the cocoons of *Hemiteles* (?) *thyridopterygis*.
 FIG. 11. *Hemiteles* (?) *thyridopterygis* Riley, female.
 FIG. 12. *Hemiteles* (?) *thyridopterygis* Riley, male.
 FIG. 13. Thistle Plume (*Pterophorus carduidactylus* Riley), moth.
 FIG. 14. Thistle Plume (*Pterophorus carduidactylus* Riley), chrysalis.
 FIG. 15. Grape-vine Plume (*Pterophorus periscelidactylus* Fitch), moth.
 FIG. 16. Grape-vine Plume (*Pterophorus periscelidactylus* Fitch), chrysalis.
 FIG. 17. Gooseberry Fruit-worm Moth (*Pempelia grossulariae* Pack.).
 FIG. 18. Chickweed Geometer (*Hæmatopis grataria*, Fabr.), moth.
 FIG. 19. Chickweed Geometer (*Hæmatopis grataria*, Fabr.), larva.
 FIG. 20. Chickweed Geometer (*Hæmatopis grataria*, Fabr.), pupa.
 FIG. 21. Chickweed Geometer (*Hæmatopis grataria*, Fabr.), eggs.
 FIG. 22. Ailanthus worm (*Eta compta*, Clem.), moth, with spread wings.
 FIG. 23. Ailanthus worm (*Eta compta*, Clem.), moth, with closed wings.
 FIG. 24. Seed-corn Maggot (*Anthomyia zææ* Riley), fly, enlarged.
 FIG. 25. Raspberry Geometer (*Aplodes rubivora* Riley), moth.
 FIG. 26. Strawberry Leaf-roller (*Anchylopera fragariae* Walsh & Riley), moth, enlarged.
 FIG. 27. Strawberry Leaf-roller (*Anchylopera fragariae* Walsh & Riley), moth, natural size.
 FIG. 28. American Meromyza (*Meromyza americana* Fitch), fly, enlarged.
 FIG. 29. Grape-berry Moth (*Penthina vitivorana* Pack.), moth, enlarged.
 FIG. 30. Grape-berry Moth (*Penthina vitivorana* Pack.), moth, natural size.

WOOD-CUTS.

- FIG. 1. Harris's Bark-louse (*Aspidiotus Harrisii* Walsh).
 FIG. 2. Oyster-shell Bark-louse (*Aspidiotus conchiformis*, Gmélin).
 FIG. 3. Oyster-shell Bark-louse (*Aspidiotus conchiformis*, Gmélin). 1, egg (natural size scarcely .01.) 2, larva, as it appears when running over the twigs (natural size .01.) 3, its appearance after becoming fixed. 4, appearance of scale after the second plate is formed. 5, form of louse (ventral view) soon after losing its members. 6, form of louse (ventral view) when full grown and just about to deposit. 7, fully formed scale, containing louse, as it appears from the under side when raised. 8, highly magnified antenna of larva, showing joints.
 FIG. 4. Twice-stabbed Ladybird (*Chilocorus bivulnerus* Muls.). [From the *Practicæ Entomologist*.]
 FIG. 5. Twice-stabbed Ladybird (*Chilocorus bivulnerus* Muls.), larva.
 FIG. 6. Seventeen-year Cicada (*Cicada septemdecim* Linn.). A, ♂ of typical form; c, d, genital hooks; g, singing apparatus. B, ♂ of the small form (*cassinii*); e, f, genital hooks.
 FIG. 7. Seventeen-year Cicada (*Cicada septemdecim*, Linn.). a, pupa; b, cast pupa shell; c, imago; d, punctured twig; e, two eggs.
 FIG. 8. Seventeen-year Cicada (*Cicada septemdecim* Linn.), galleries made by pupa; a, front view, e, orifice; b, section, c, pupa awaiting time of change, d, pupa ready to transform.
 FIG. 9. Twig punctured by the Seventeen-year Cicada (*Cicada septemdecim* Linn.).
 FIG. 10. Twig healed after the puncture of the Seventeen-year Cicada (*Cicada septemdecim* Linn.).
 FIG. 11. Thirteen-year Cicada (*Cicada tredecim* Linn.), newly hatched larva.
 FIG. 12. *Stizus grandis* Say, ♀.
 FIG. 13. Seventeen-year Cicada (*Cicada septemdecim* Linn.), side view of ♀ to show beak, a, and ovipositor, b.

- FIG. 14. Round-headed Apple-tree Borer (*Saperda bivittata* Say). *a*, larva; *b*, pupa; *c*, imago.
- FIG. 15. Flat-headed Apple-tree Borer (*Chrysobothris femorata*, Fabr.), larva.
- FIG. 16. Flat-headed Apple-tree Borer (*Chrysobothris femorata*, Fabr.), imago.
- FIG. 17. Peach-tree Borer (*Ageria exitiosa* Say); 1, ♀; 2, ♂.
- FIG. 18. Plum Curculio (*Conotrachelus nenuphar*, Herbst); *a*, larva; *b*, pupa; *c*, imago; *d*, plum and curculio, natural size, the plum bearing one of the punctures.
- FIG. 19. Pennsylvania Soldier-beetle (*Chauliognathus pensylvanicus*, DeGeer). *a*, larva, natural size; *b*, head and first segment enlarged; *c*, under lip (*labium*); *d*, upper lip (*labrum*); *e*, leg; *f*, left lower jaw (*maxilla*); *g*, antenna; *h*, left upper jaw (*mandible*).
- FIG. 20. Lacewing (*Chrysopa* sp.); *a*, eggs; *b*, larva; *c*, cocoon, the upper figure showing the lid; *d*, imago. [*a*, *b*, *d* after Westwood.]
- FIG. 21. Subangular Ground-beetle (*Aspidoglossa subangulata* Chaud.).
- FIG. 22. Carabid larva. *A*, natural size; *B*, under side of head, enlarged; *c*, mandible; *e*, antenna; *f*, labium and labial palpi; *g*, maxilla and its palpi; *h*, joint 12 beneath; *i*, joint 11 beneath; *j*, joints 4-10 each beneath—enlarged.
- FIG. 23. Pennsylvania Ground-beetle (*Harpalus pensylvanicus*, DeGeer).
- FIG. 24. Codling-moth (*Carpocapsa pomonella*, Linn.) *a*, apple showing the work of the larva; *b*, point of entrance of the larva; *d*, pupa; *e*, larva; *f*, *g*, moth; *h*, head of larva; *i*, cocoon.
- FIG. 25. Pupa of Cut-worm in earthen cell. [After Curtis.]
- FIG. 26. Dark-sided Cut-worm (*Agrotis Cochranii* Riley). *a*, larva; *b*, moth.
- FIG. 27. W-marked Cut-worm (*Noctua clandestina* Harr.).
- FIG. 28. Lance Rustic (*Agrotis telifera* Harr.), moth.
- FIG. 29. Gothic Dart (*Agrotis subgothica*, Haw.), moth.
- FIG. 30. Glassy Cut-worm (*Agrotis devastator*, Brace). Lower figure represents the side of one of the middle segments.
- FIG. 31. Figure 8 Minor (*Celæna renigera* Steph.). *a*, moth; *b*, larva.
- FIG. 32. *Microgaster militaris* Walsh. [After Walsh.]
- FIG. 33. Spined Soldier-bug (*Arma spinosa* Dallas). *a*, beak magnified; *b*, bug with right wing spread.
- FIG. 34. Fiery Ground-beetle (*Calosoma calidum*, Fabr.); *a*, larva; *b*, beetle.
- FIG. 35. Potato-stalk Borer (*Gortyna nitela* Guen.) 1, moth; 2, larva.
- FIG. 36. Potato-stalk Borer (*Gortyna nitela* Guen.) larva.
- FIG. 37. Potato-stalk Weevil (*Baridius trinotatus*, Say); *a*, larva; *b*, pupa; *c*, beetle, (all enlarged).
- FIG. 38. Potato- or Tomato-worm (*Sphinx 5-maculata* Haw.). *A*, larva; *B*, pupa; *C*, moth. [After Harris.]
- FIG. 39. Striped Blister-beetle (*Lytta vittata* Fabr.). [From *Practical Entomologist*.]
- FIG. 40. *a*, Ash-gray Blister-beetle (*Lytta cinerea* Fabr.), *d*, antennæ; *b*, Black-rat Blister-beetle (*Lytta murina* Lec.), *c*, antennæ.
- FIG. 41. Margined Blister-beetle (*Lytta marginata* Fabr.). [From *Practical Entomologist*.]
- FIG. 42. Three-lined Potato-beetle (*Lema trilineata*, Oliv.); *a*, larva; *b*, tip of its body; *c*, pupa; *d*, eggs. [From *Practical Entomologist*.]
- FIG. 43. Three-lined Potato-beetle (*Lema trilineata*, Oliv.). [From *Practical Entomologist*.]
- FIG. 44. Striped Cucumber-beetle (*Diabrotica vittata*, Fabr.). [From *Practical Entomologist*.]
- FIG. 45. Cucumber Flea-beetle (*Haltica cucumeris* Harr.). [From *Practical Entomologist*.]
- FIG. 46. Colorado Potato-beetle (*Doryphora 10-lineata*, Say); *a*, eggs; *b*, larva, in different stages; *c*, pupa; *d*, imago or beetle; *e*, wing-cover, enlarged; *f*, leg, enlarged.

- FIG. 47. Bogus Colorado Potato-beetle (*Doryphora juncta*, Germar); *a*, eggs; *b*, larva; *c*, beetle; *d*, wing-cover, enlarged; *e*, leg, enlarged.
- FIG. 48. Colorado Potato-beetle Parasite (*Lydella doryphoræ* Riley).
- FIG. 49. Spotted Ladybird (*Hippodamia maculata*, DeGeer). [From *Practical Entomologist*.]
- FIG. 50. Nine-spotted Ladybird (*Coccinella 9-notata* Herbst). [From *Practical Entomologist*.]
- FIG. 51. Thirteen-spotted Ladybird (*Hippodamia 13-punctata*, Linn.).
- FIG. 52. Convergent Ladybird (*Hippodamia convergens* Guer.)
- FIG. 53. Ladybird larva. [After Westwood.]
- FIG. 54. Spined Soldier-bug (*Arma spinosa* Dallas); *a*, beak enlarged; *b*, bug; *c*, enlarged beak of an allied plant-feeder (*Euschistus punctipes*, Say).
- FIG. 55. Common Squash-bug (*Coreus tristis*, DeGeer); *b*, enlarged beak.
- FIG. 56. Bordered Soldier-bug (*Stiretrus fimbriatus*, Say).
- FIG. 57. Many-banded Robber (*Harpactor cinctus*, Fabr.); *b*, enlarged beak.
- FIG. 58. Rapacious Soldier-bug (*Reduvius raptatorius* Say).
- FIG. 59. Virginian Tiger-beetle (*Tetracha virginica* Hope).
- FIG. 60. Fiery Ground-beetle (*Calosoma calidum*, Fabr.).
- FIG. 61. Elongate Ground-beetle (*Pasimachus elongatus* Lec.).
- FIG. 62. Murky Ground-beetle (*Harpalus caliginosus* Say).
- FIG. 63. Pincers for crushing Potato-beetles.
- FIG. 64. Apple-root Plant-louse (*Eriosoma pyri*, Fitch); *a*, affected root; *b*, larva; *c*, winged louse; *d*, leg; *e*, proboscis; *f*, antenna of winged louse; *g*, antenna of larva (all greatly enlarged).
- FIG. 65. Vagabond Plant-louse (*Pemphigus vagabundus*, Walsh).
- FIG. 66. Root-louse Syrphus-fly (*Pipiza radicum* Riley); *a*, larva; *b*, puparium from which the fly has emerged; *c*, fly.
- FIG. 67. Gigantic Grape-root Borer (*Prionus laticollis*, Drury).
- FIG. 68. Gigantic Grape-root Borer (*Prionus laticollis*, Drury); head and thoracic joints.
- FIG. 69. Cylindrical Orthosoma (*Orthosoma cylindricum*, Fabr.).
- FIG. 70. Grape Curculio (*Cæliodes inæqualis*, Say); *a*, infested grape; *b*, larva.
- FIG. 71. Grape Curculio (*Cæliodes inæqualis*, Say). [After Walsh.]
- FIG. 72. Grape Curculio (*Cæliodes inæqualis*, Say); front leg. [After Walsh.]
- FIG. 73. Grape-seed Maggot (*Isosoma vitis* Saunders).
- FIG. 74. Grape-cane Gall-curculio (*Baridius Sesostris* Lec.).
- FIG. 75. Grape-vine Fidia (*Fidia viticida* Walsh). [From *Practical Entomologist*.]
- FIG. 76. Grape Fruit-worm (*Penthina vitivorana* Pack. = *Lobesia botrana* Schiff.); *a*, pupa; *b*, cocoon.
- FIG. 77. Snowy Tree-cricket (*Ecanthus niveus* Harr.), ♂. [From *Practical Entomologist*.]
- FIG. 78. Snowy Tree-cricket (*Ecanthus niveus* Harr.), ♀. [From *Practical Entomologist*.]
- FIG. 79. Gooseberry Fruit-worm (*Pempelia grossulariæ* Pack.); *a*, cocoon; *b*, moth. [After Packard.]
- FIG. 80. Strawberry Leaf-roller (*Anchylopera fragariæ* Walsh & Riley; *a*, larva; *b*, anterior part enlarged; *d*, anal segment; *c*, moth.
- FIG. 81. White-marked Tussock Moth (*Orgyia leucostigma*, Sm. & Abb.); *a*, ♀ on cocoon; *b*, larva; *c*, female pupa; *d*, male pupa.
- FIG. 82. White-marked Tussock Moth (*Orgyia leucostigma*, Sm. & Abb.); female caterpillar.
- FIG. 83. White-marked Tussock Moth (*Orgyia leucostigma*, Sm. & Abb.), male.
- FIG. 84. Bag-worm (*Thyridopteryx ephemeraeformis* Haw.); *a*, larva; *b*, male chrysalis; *c*, female moth; *d*, male moth; *e*, female chrysalis in bag, sectional view; *f*, caterpillar and bag; *g*, very young caterpillars in their bags.
- FIG. 85. Walnut Tortrix (*Tortrix Rileyana* Grote); *a*, larva; *b*, side view of one segment.

- FIG. 86. Seed-corn Maggot (*Anthomyia zea* Riley); *a*, enlarged; *b*, puparium.
- FIG. 87. Seed-corn Maggot (*Anthomyia zea* Riley); kernels of corn containing the maggot.
- FIG. 88. White Grub or May-beetle (*Lachnosterna quercina*, Knoch); 1, pupa; 2, the grub; 3, 4, the beetle.
- FIG. 89. White Grub attacked by fungus.
- FIG. 90. American Meromyza (*Meromyza americana* Fitch); *a*, infested stalk; *b*, maggot; *c*, pupa.
- FIG. 91. Sheep Head Maggot (*Estrus ovis* Linn.); 1 and 2, the Gad-fly; 3, the puparium; 4, larva, dorsal view; 5, larva, ventral view; 6, younger larva; *a*, head; *b*, corneous appendages at anus; *c*, spiracles.
- FIG. 92. Bee-moth (*Galleria cereana* Fabr.); *a*, larva; *b*, cocoon; *c*, pupa; *d*, *e*, moth.
- FIG. 93. Nebraska Bee-killer (*Trupanea apivora* Fitch = *Promachus Fitchii* O. S.).
- FIG. 94. Camel-cricket (*Mantis carolina*, Linn.); *a*, female; *b*, male.
- FIG. 95. Camel-cricket (*Mantis carolina*, Linn.), egg-masses.
- FIG. 96. Solidago Gall of *Gelechia gallæsolidaginis* Riley; *a*, section of gall; *b*, whole gall; *c*, orifice through which the moth escapes; *d*, excrement of the larva; *e*, larva.
- FIG. 97. *Eurytoma Bolteri* Riley; antennæ of ♂ and ♀.
- FIG. 98. Thistle Plume-moth (*Pterophorus carduidactylus* Riley = *Pt. cardui* Zell. emend), anterior and posterior joints of the larva.

REPORT II.

- FIG. 1. Chinch-bug (*Micropus leucopterus*, Say).
- FIG. 2. Chinch-bug (*Micropus leucopterus*, Say), short-winged form.
- FIG. 3. Spotted Ladybird (*Hippodamia maculata*, DeGeer). [From *Practical Entomologist*.]
- FIG. 4. Trim Ladybird (*Coccinella munda* Say).
- FIG. 5. Lacewing (*Chrysopa* sp.). [After Westwood.]
- FIG. 6. Insidious Flower-bug (*Anthocoris insidiosus*, Say).
- FIG. 7. Spined Soldier-bug (*Arma spinosa* Dallas).
- FIG. 8. Ash-gray Leaf-bug (*Piesma cinerea*, Say).
- FIG. 9. Flea-like Negro-bug (*Corimelæna pulicaria*, Germar).
- FIG. 10. Bordered Soldier-bug (*Stiretrus fimbriatus*, Say).
- FIG. 11. Tent-caterpillar of the Forest (*Clisiocampa sylvatica* Harr.).
- FIG. 12. Cotton-worm (*Anomis xyliana*, Say); *a*, egg; *b*, worm, one-third grown; *d*, top view; *c*, side view of full-grown worm; *e*, cocoon; *f*, chrysalis. [Adapted from Glover.]
- FIG. 13. Cotton-worm Moth (*Anomis xyliana*, Say); *a*, with wings expanded; *b*, wings closed.
- FIG. 14. Army-worm (*Leucania unipuncta* Haw.).
- FIG. 15. Army-worm (*Leucania unipuncta* Haw.), chrysalis.
- FIG. 16. Army-worm Moth (*Leucania unipuncta* Haw.).
- FIG. 17. Red-tailed Tachina-fly (*Exorista leucaniæ* Kirk.).
- FIG. 18. Yellow-tailed Tachina-fly (*Exorista flavicauda* Riley).
- FIG. 19. Glassy Mesochorus (*Mesochorus vitreus* Walsh). [After Walsh.]
- FIG. 20. *Pezomachus minimus* Walsh. [After Walsh.]
- FIG. 21. *Pezomachus minimus* Walsh; bunch of cocoons. [After Walsh.]
- FIG. 22. *Chalcis albifrons* Walsh. [After Walsh.]
- FIG. 23. *Microgaster militaris* Walsh. [After Walsh.]
- FIG. 24. *Glyphe viridascens* Walsh. [After Walsh.]
- FIG. 25. *Ophion purgatus* Say.
- FIG. 26. Clubbed Tortoise-beetle (*Deloyala clavata*, Oliv.).
- FIG. 27. Two-striped Sweet-potato Beetle (*Cassida bivittata* Say); 2, larva; 3, pupa; 4, beetle.

- FIG. 28. *Chelymorphe cribraria*, Fabr.; pupa (enlarged). [After Packard.]
- FIG. 29. *Chelymorphe cribraria*, Fabr. (enlarged). [After Packard.]
- FIG. 30. *Physonota quinquepunctata* Walsh & Riley; *a*, larva; *b*, beetle.
- FIG. 31. Golden Tortoise-beetle (*Cassida aurichalcea*, Fabr.), egg.
- FIG. 32. Two-striped Sweet-potato Beetle (*Cassida bivittata* Say), larvæ.
- FIG. 33. Golden Tortoise-beetle (*Cassida aurichalcea*, Fabr.), larva; *a*, natural size; *b*, enlarged and with the dung taken from the fork.
- FIG. 34. Golden Tortoise-beetle (*Cassida aurichalcea*, Fabr.); *a*, pupa; *b*, beetle.
- FIG. 35. Mottled Tortoise-beetle (*Cassida guttata*, Oliv.); *a*, larva; *b*, pupa.
- FIG. 36. Mottled Tortoise-beetle (*Cassida guttata*, Oliv.).
- FIG. 37. Black-legged Tortoise-beetle (*Cassida nigripes* Oliv.); *a*, larva; *b*, larva cleaned and enlarged; *c*, pupa (enlarged).
- FIG. 38. Black-legged Tortoise-beetle (*Cassida nigripes* Oliv.).
- FIG. 39. Striped Cucumber-beetle (*Diabrotica vittata*, Fabr.). [From *Practical Entomologist*.]
- FIG. 40. Striped Cucumber-beetle (*Diabrotica vittata*, Fabr.), larva; *a*, dorsal view; *b*, side view.
- FIG. 41. Striped Cucumber-beetle (*Diabrotica vittata*, Fabr.) pupa; 1, ventral; 2, dorsal view.
- FIG. 42. Twelve-spotted Diabrotica (*Diabrotica 12-punctata*, Oliv.). [From *Practical Entomologist*.]
- FIG. 43. Pickle-worm (*Phacellura nitidalis* Cram.); *a*, natural size; *b*, head and first joints, enlarged; *c*, side view of a joint, enlarged; *d*, cervical shield, enlarged; *e*, side of first joint, enlarged; *f*, 2d joint from above, enlarged; *g*, anal joint, enlarged; *h*, cocoon; *i*, moth, male.
- FIG. 44. Hog-caterpillar of the Vine (*Charocampa pampinatrix*, Sm. & Abb.).
- FIG. 45. Hog-caterpillar of the Vine (*Charocampa pampinatrix*, Sm. & Abb.), chrysalis.
- FIG. 46. Hog-caterpillar of the Vine (*Charocampa pampinatrix*, Sm. & Abb.), moth.
- FIG. 47. Microgaster cocoons or Hog-caterpillar of the Vine (*Char. pampinatrix*, Sm. & Abb.) [After Harris.]
- FIG. 48. *Microgaster* = *Apanteles*. [After Harris.]
- FIG. 49. Achemon Sphinx (*Philampelus achemon*, Drury), caterpillar.
- FIG. 50. Achemon Sphinx (*Philampelus achemon*, Drury), chrysalis.
- FIG. 51. Achemon Sphinx (*Philampelus achemon*, Drury), moth.
- FIG. 52. Satellite Sphinx (*Philampelus satellitia*, Linn.); *a*, full-grown larva; *b*, its position at rest; *c*, young larva.
- FIG. 53. Satellite Sphinx (*Philampelus satellitia*, Linn.), moth.
- FIG. 54. Abbot Sphinx (*Thyreus Abbotii* Swainson); larva and moth.
- FIG. 55. Eight-spotted Forrester (*Alypia octomaculata*, Fabr.); *a*, caterpillar; *b*, side view of one joint; *c*, moth.
- FIG. 56. Beautiful Wood-nymph (*Eudryas grata*, Fabr.).
- FIG. 57. ? Pearl Wood-nymph (*Eudryas unio*, Hüb.); *a*, larva; *b*, side view of one segment enlarged; *c*, hump on 11th joint, enlarged. (See 3d Rep., Fig. 25.)
- FIG. 58. American Procris (*Procris americana* Boisd.); *a*, larva; *b*, chrysalis; *c*, cocoon; *d*, *e*, moth.
- FIG. 59. American Procris (*Procris americana* Boisd.), larvæ.
- FIG. 60. Gigantic Grape-root Borer (*Prionus laticollis*, Drury).
- FIG. 61. Broad-necked Prionus (*Prionus laticollis*, Drury), female.
- FIG. 62. Gigantic Grape-root Borer (*Prionus laticollis*, Drury), pupa.
- FIG. 63. Tile-horned Prionus (*Prionus imbricornis*, Linn.), male.
- FIG. 64. Grape-seed Maggot (*Isosoma vitis* Saunders).
- FIG. 65. Joint-worm Fly (*Isosoma hordei*, Harr.); *a*, female; *b*, male; *c*, ♀ antenna; *d*, ♂ antenna; *e*, ♀ abdomen; *f*, ♂ abdomen.

- FIG. 66. Canker-worm; *a*, eggs of Fall Canker-worm (*Anisopteryx pometaria* Harr.); *b*, five eggs of same, enlarged; *c*, larva of Spring Canker-worm (*Paleacrita vernata*, Peck), *d*, cocoon, *e*, chrysalis, *f*, male moth, *g*, female moth—all probably of *vernata*. (See 6th Rept., p. 29). [*a, b, c, d, e*, after Harris; *f, g*, after Packard.]
- FIG. 67. Spring Canker-worm (*Paleacrita vernata*, Peck), head enlarged.
- FIG. 68. Mite (*Nothrus ovivorus* Pack.), enlarged. [After Packard.]
- FIG. 69. Rummaging Ground-beetle (*Calosoma scrutator*, Fabr.).
- FIG. 70. Fiery Ground-beetle (*Calosoma calidum*, Fabr.).
- FIG. 71. Fraternal Potter-wasp (*Eumenes fraterna* Say); *b*, clay nest; *c*, same cut open.
- FIG. 72. Southern Cabbage-butterfly (*Pieris protodice* Boisd.); *a*, caterpillar; *b*, chrysalis.
- FIG. 73. Southern Cabbage-butterfly (*Pieris protodice* Boisd.), female.
- FIG. 74. Southern Cabbage-butterfly (*Pieris protodice* Boisd.), male.
- FIG. 75. Potherb Butterfly (*Pieris oleracea* Boisd.). [After Harris.]
- FIG. 76. Potherb Butterfly (*Pieris oleracea* Boisd.), chrysalis. [After Harris.]
- FIG. 77. Imported Cabbage-butterfly (*Pieris rapæ* Schrank.); *a*, larva; *b*, chrysalis. [After Curtis.]
- FIG. 78. Imported Cabbage-butterfly (*Pieris rapæ* Schrank.), female.
- FIG. 79. Imported Cabbage-butterfly (*Pieris rapæ* Schrank.), male.
- FIG. 80. Butterfly Net: 5, socket; 6, ring.
- FIG. 81. Cabbage Plusia (*Plusia brassicæ* Riley); *a*, caterpillar; *b*, chrysalis in cocoon; *c*, moth, male.
- FIG. 82. Zebra-caterpillar (*Mamestra picta* Harr.); *a*, caterpillar; *b*, moth.
- FIG. 83. Tarnished Plant-bug (*Capsus oblineatus* Say).
- FIG. 84. Philenor Swallow-tail (*Papilio philenor* Drury), caterpillar.
- FIG. Philenor Swallow-tail (*Papilio philenor* Drury); *a*, chrysalis, back view; *b*, lateral outline.
- FIG. 86. Philenor Swallow-tail (*Papilio philenor* Drury).
- FIG. 87. Cottonwood Dagger (*Acronycta populi* Riley); caterpillar.
- FIG. 88. Cottonwood Dagger (*Acronycta populi* Riley).
- FIG. 89. Missouri Bee-killer (*Asilus missouriensis* Riley).
- FIG. 90. Wing of *Promachus* (*a*), *Asilus* (*b*), *Erax* (*c*).
- FIG. 91. Silky Asilus (*Asilus sericeus* Say). [After Harris.]
- FIG. 92. *Erax bastardi* Macq., larva.
- FIG. 93. *Erax bastardi* Macq.; *a*, fly; *b*, pupa.
- FIG. 94. Goat-weed Butterfly (*Paphia glycerium* Doubl.); *a*, caterpillar; *b*, chrysalis.
- FIG. 95. Goat-weed Butterfly (*Paphia glycerium* Doubl.), male.
- FIG. 96. Goat-weed Butterfly (*Paphia glycerium* Doubl.), female.
- FIG. 97. Black Breeze-fly (*Tabanus atratus* Fabr.); *a*, larva; *b*, pupa shell; *c*, fly.
- FIG. 98. False-indigo Gall-moth (*Walshia amorphella* Clem.); *a*, moth; *b*, caterpillar; *c*, gall; *d*, section of gall, showing larva in burrow.
- FIG. 99. Misnamed Gall-moth (*Euryptychia saligneana* Clem.); *a*, moth; *b*, gall with protruding pupa-shell.

REPORT III.

- FIG. 1. Plum Curculio (*Conotrachelus nenuphar*, Herbst); *a*, larva; *b*, pupa; *c*, curculio, enlarged; *d*, punctured plum with curculio resting on it, natural size.
- FIG. 2. The Hull Curculio-catcher.
- FIG. 3. The Hull Curculio-catcher; viewed from beneath; *a*, slide for closing central hole, *d*; *b, b*, handles; *c, c*, wheels; *e, f*, position of bag.
- FIG. 4. The Hull Curculio-catcher; viewed from above.
- FIG. 5. Strips of sheeting for closing up the tree-way in Hull's Curculio-catcher.
- FIG. 6. The Hooten Curculio-catcher.

- FIG. 7. *Sigalphus Curculio*-parasite (*Sigalphus curculionis* Fitch); *a*, male; *b*, female; *c*, antenna.
- FIG. 8. *Sigalphus Curculio*-parasite (*Sigalphus curculionis* Fitch); *a*, larva; *b*, cocoon; *c*, pupa.
- FIG. 9. Porizon *Curculio*-parasite (*Porizon conotracheli* Riley); *a*, ♀; *b*, ♂; *c*, antenna.
- FIG. 10. Apple *Curculio* (*Anthonomus quadrigibbus* Say); *a*, natural size; *b*, side view; *c*, back view.
- FIG. 11. Apple *Curculio* (*Anthonomus quadrigibbus* Say); *a*, pupa; *b*, larva.
- FIG. 12. Quince *Curculio* (*Conotrachelus crataegi* Walsh.); *a*, side; *b*, back.
- FIG. 13. Plum Gouger (*Anthonomus prunivora* Walsh.).
- FIG. 14. Strawberry Crown-borer (*Analcis fragariae* Riley); *a*, larva; *b*, side view of beetle; *c*, dorsal view.
- FIG. 15. Pea-weevil (*Bruchus pisi* Linn.); *a*, beetle; *b*, injured pea.
- FIG. 16. Pea-weevil (*Bruchus pisi* Linn.), egg enlarged.
- FIG. 17. Pea-weevil (*Bruchus pisi* Linn.); *b*, beetle, side view; *c*, larva; *d*, pupa, dorsal view; *g*, pea, infested. [After Curtis.]
- FIG. 18. Grain *Bruchus* (*Bruchus granarius* Linn.). [After Curtis.]
- FIG. 19. American Bean-weevil (*Bruchus fabae* Riley); *a*, beetle; *b*, bean, infested.
- FIG. 20. New York Weevil (*Ithycerus novæboracensis*, Forster); *a*, excavation made by female to deposit eggs; *b*, larva; *c*, beetle.
- FIG. 21. Imbricated Snout-beetle (*Epicærus imbricatus*, Say).
- FIG. 22. Corn *Sphenophorus* (*Sphenophorus zæ* Walsh); *a*, back view; *b*, outline side view; *c*, enlarged punctures of elytra.
- FIG. 23. Cocklebur *Sphenophorus* (*Sphenophorus pulchellus* Schæn.); *a*, back view; *b*, outline side view.
- FIG. 24. Grape Leaf-folder (*Desmia maculalis* Westw.); 1, caterpillar in folded leaf; 2, enlarged view of head and anterior joints; 3, chrysalis; 4, male moth; 5, female moth.
- FIG. 25. Grape-vine *Epimenis* (*Psychomorpha epimenis*, Drury); *a*, larva; *b*, side view of one segment, enlarged; *c*, hump on 11th joint, enlarged.
- FIG. 26. Grape-vine *Epimenis* (*Psychomorpha epimenis*, Drury), moth, male.
- FIG. 27. Grape-vine Plume (*Pterophorus periscelidactylus* Fitch); *a*, caterpillars in their retreat; *b*, chrysalis; *c*, one of the dorsal processes of chrysalis; *d*, moth; one joint of larva enlarged, side view.
- FIG. 28. Yellow-bear Caterpillar (*Spilosoma virginica*, Fabr.); *a*, caterpillar; *b*, chrysalis; *c*, moth.
- FIG. 29. Smeared Dagger (*Acronycta obliqua*, Sm. & Abb.); *a*, caterpillar; *b*, cocoon; *c*, moth.
- FIG. 30. *Aleiodes Rileyi* Cress.; hardened skin of caterpillar of the Smeared Dagger (*Acronycta obliqua*, Sm. & Abb.) from which the *Aleiodes* has emerged.
- FIG. 31. Pyramidal Grape-vine Worm (*Amphipyra pyramidoides* Guen.), moth.
- FIG. 32. Pyramidal Grape-vine Worm (*Amphipyra pyramidoides* Guen.).
- FIG. 33. Grape-root Borer (*Ægeria polistiformis* Harr.); *a*, male; *b*, female.
- FIG. 34. Spotted Pelidnota (*Pelidnota punctata*, Linn.); *a*, larva; *b*, pupa; *c*, beetle; *d*, anal joint of larva; *e*, antenna of larva; *f*, leg of larva.
- FIG. 35. Grape-vine Flea-beetle (*Haltica chalybea* Illiger); *a*, larvæ on leaf; *b*, larva, enlarged; *c*, earthen cell containing pupa; *d*, beetle. [*d* after Harris.]
- FIG. 36. Grape-vine Flea-beetle (*Haltica chalybea* Illiger). [From *Practical Entomologist*.]
- FIG. 37. Grape-vine *Colaspis* (*Colaspis flavida* Say); 1, enlarged; 2, natural size.
- FIG. 38. Grape-vine *Colaspis* (*Colaspis flavida* Say); *a*, enlarged side view of larva; *b*, terminal joints seen from beneath.
- FIG. 39. Galls of the Grape Phylloxera (*Phylloxera vitifoliae*, Fitch = *vastatrix* Pl.).

- FIG. 40. Grape Phylloxera (*Phylloxera vitifoliae*, Fitch = *Ph. vastatrix* Pl.); *a*, the winged female; *b*, her foot or tarsus—after Signoret; *c*, egg; *d*, newly-hatched gall-inhabiting type; *e*, same, dorsal view; *f*, section of gall; *g*, tubercled root-inhabiting form; *h*, mother gall-louse at height of her fertility; *i*, same, dorsal view; *j*, *k*, differently veined wings of the Oak Phylloxera of Europe.
- FIG. 41. Great Lebia (*Lebia grandis* Hentz.).
- FIG. 42. Boll-worm (*Heliothis armigera* Hübn.) on tomato.
- FIG. 43. Boll-worm (*Heliothis armigera* Hübn.); *a*, egg, side view; *b*, egg, top view; *c*, caterpillar; *d*, chrysalis in earthen cocoon; *e*, moth, wings expanded; *f*, moth, wings closed. [*a*, *b*, *c*, *d* after Glover.]
- FIG. 44. Army-worm (*Leucania unipuncta* Haw.).
- FIG. 45. Fall Army-worm (*Prodenia autumnalis* Riley = *Laphygma frugiperda*, Sm. & Abb.); *a*, natural size; *b*, head magnified; *c*, one segment enlarged, from above; *d*, same, from side.
- FIG. 46. Fall Army-worm (*Prodenia autumnalis* Riley = *Laphygma frugiperda*, Sm. & Abb.); *a*, *b*, *c*, three varieties.
- FIG. 47. Army-worm Moth (*Leucania unipuncta* Harr.).
- FIG. 48. Spiderwort Owlet-moth (*Prodenia commelinæ*, Abb.); *a*, caterpillar; *b*, *c*, dark and light varieties of the moth. [See Notes, etc., p. 56.]
- FIG. 49. Unarmed Rustic (*Agrotis inermis* Harr. = *A. saucia* Hübn.); *a*, egg, enlarged; *b*, batch of eggs, natural size.
- FIG. 50. Apple-tree Tent-caterpillar (*Clisiocampa americana* Harr.); *a*, *b*, caterpillars; *c*, eggs; *d*, cocoon.
- FIG. 51. Apple-tree Tent-caterpillar (*Clisiocampa americana* Harr.) moth.
- FIG. 52. Tent-caterpillar of the Forest (*Clisiocampa sylvatica* Harr.); *a*, eggs; *b*, female moth; *c*, egg enlarged, top view; *d*, enlarged eggs, side view.
- FIG. 53. Tent-caterpillar of the Forest (*Clisiocampa sylvatica* Harr.).
- FIG. 54. Rummaging Ground-beetle (*Calosoma scrutator*, Fabr.).
- FIG. 55. Fall Web-worm (*Hyphantria textor* Harr.); *a*, caterpillar; *b*, chrysalis; *c*, moth.
- FIG. 56. Blue-spangled Peach-worm (*Callimorpha fulvicosta* Clem.); *a*, caterpillar; *b*, moth; *c*, one segment enlarged, side view; *d*, same, top view.
- FIG. 57. Ash-gray Pinion (*Xylina cinerea* Riley); *a*, worm in fruit; *b*, moth.
- FIG. 58. Glassy-winged Soldier-bug (*Campyloneura vitripennis*, Say).
- FIG. 59. Glassy-winged Soldier-bug (*Campyloneura vitripennis*, Say), pupa.
- FIG. 60. White-lined Morning Sphinx (*Deilephila lineata*, Fabr.), moth.
- FIG. 61. White-lined Morning Sphinx (*Deilephila lineata*, Fabr.), caterpillar, light form.
- FIG. 62. White-lined Morning Sphinx (*Deilephila lineata*, Fabr.); caterpillar, dark form.
- FIG. 63. Archippus Butterfly (*Danais archippus*, Fabr.).
- FIG. 64. Archippus Butterfly (*Danais archippus*, Fabr.); *a*, egg, greatly enlarged; *c*, natural size; *e*, *f*, lateral and dorsal views of a segment of the larva in its first stage, enlarged; *b*, larva in act of casting its skin, to show how the flexible horns are folded (*d*).
- FIG. 65. Archippus Butterfly (*Danais archippus*, Fabr.), caterpillar.
- FIG. 66. Archippus Butterfly (*Danais archippus*, Fabr.); *a*, *b*, *c*, successive stages in changing from caterpillar to chrysalis.
- FIG. 67. Archippus Butterfly (*Danais archippus*, Fabr.), chrysalis.
- FIG. 68. Disippus Butterfly (*Limenitis disippus*, Godt.), showing upper surface of left wing, and under surface on the right. [After Harris.]
- FIG. 69. Disippus Butterfly (*Limenitis disippus*, Fabr.); *a*, egg greatly enlarged; *c*, natural size; *d*, one cell of the egg-shell, greatly magnified; *b*, one segment of the larva, in its first stage.
- FIG. 70. Disippus Butterfly (*Limenitis disippus*, Fabr.); *a*, caterpillar; *b*, chrysalis; *c*, hibernaculum; *d*, leaf cut for hibernaculum.

- FIG. 71. Disippus Butterfly (*Limenitis disippus*, Fabr.); a leaf eaten by the caterpillar.
- FIG. 72. Disippus Egg-parasite (*Trichogramma? minuta* Riley); *a*, fly with wings folded; *b*, front wing; *c*, hind wing; *d*, leg; *e*, antenna— all enlarged.
- FIG. 73. *Microgaster militaris* Walsh. [After Walsh.]

REPORT IV.

- FIG. 1. Perforated tin box for sifting paris green.
- FIG. 2. Creighton's "Improved Patent Insect Destroyer."
- FIG. 3. Grand-Daddy-Long-Legs (*Phalangium dorsatum* Say).
- FIG. 4. Fifteen-spotted Ladybird (*Mysia 15-punctata*, Oliv.); *a*, larva; *b*, pupa; *c*, first joint of larva, enlarged; *d*, *e*, *f*, *g*, different varieties of the beetle.
- FIG. 5. Icy Ladybird (*Hippodamia glacialis* Fabr.).
- FIG. 6. Ring-banded Soldier-bug (*Perillus circumcinctus* Stål); *b*, antenna; *c*, beak (enlarged).
- FIG. 7. Dotted-legged Plant-bug (*Euschistus punctipes*, Say); *c*, beak (enlarged).
- FIG. 8. Spined Soldier bug (*Arma spinosa* Dallas); *a*, beak (enlarged).
- FIG. 9. Spined Soldier-bug (*Arma spinosa* Dallas); *a*, pupa; *b*, larva; *c*, egg (all enlarged).
- FIG. 10. Rove-beetle (*Philonthus apicalis*, Say).
- FIG. 11. Rove-beetle larva (*Goërius olens*). [After Westwood.]
- FIG. 12. Rove-beetle (*Quedius molochinus*, Grav.), pupa.
- FIG. 13. Wier's Apple-worm Trap.
- FIG. 14. Pennsylvania Soldier-beetle (*Chauliognathus pensylvanicus* DeG.); *a*, larva; *b*, head and prothorax, enlarged; *c*, labium; *d*, labrum; *e*, leg; *f*, maxilla; *g*, antenna; *h*, mandible.
- FIG. 15. Two-lined Soldier-beetle (*Telephorus bilineatus*, Say); *a*, larva; *b*, anterior joints enlarged; *c*, beetle.
- FIG. 16. Grape-vine Colaspis (*Colaspis flavida* Say); one joint of larva, viewed from beneath and enlarged; *b*, head of larva, from beneath; *c*, same, from above, enlarged.
- FIG. 17. Harlequin Cabbage-bug (*Strachia histrionica* Hahn); *a*, larva; *b*, pupa; *c*, eggs; *d*, eggs enlarged, side view; *e*, same, top view; *g*, bug; *h*, same, with wings expanded.
- FIG. 18. Rascal Leaf-crumpler (*Phycita nebulo* Walsh); *a*, case, containing caterpillar; *b*, cases in winter; *c*, head and thoracic joints of larva, enlarged; *d*, moth.
- FIG. 19. Larval cases of the Rascal Leaf-crumpler (*Phycita nebulo* Walsh) in winter.
- FIG. 20. Walnut Case-bearer (*Acrobasis juglandis* LeBaron); *a*, case between two leaflets; *b*, case; *c*, wings of *nebulo* for comparison; *d*, wings of moth; *e*, wings of a variety of same from the crab-apple.
- FIG. 21. Apple-leaf Skeletonizer (*Pempelia Hammondi* Riley); *a*, larva; *b*, middle joint, enlarged; *c*, anterior joints, enlarged; *d*, moth.
- FIG. 22. Green Apple-leaf-tyer (*Tortrix cinderella* Riley); *a*, caterpillar; *b*, chrysalis; *c*, moth; *d*, pupal case.
- FIG. 23. Apple-leaf Bucculatrix (*Bucculatrix pomifoliella* Clem.); *a*, cocoons on twig; *b*, cocoon, enlarged; *c*, moth.
- FIG. 24. Apple-twigg Borer (*Bostrichus bicaudatus*, Say). [After Walsh.]
- FIG. 25. Apple-twigg Borer (*Bostrichus bicaudatus*, Say); twigs bored by this insect.
- FIG. 26. Red-shouldered Sinoxylon (*Sinoxylon basilare*, Say); *a*, larva; *b*, pupa; *c*, beetle.
- FIG. 27. Red-shouldered Sinoxylon (*Sinoxylon basilare*, Say); *a*, head and thoracic joints of larva greatly enlarged; *b*, labrum and mandibles; *c*, anterior leg; *d*, intermediate leg; *e*, posterior leg.

- FIG. 28. Grape Phylloxera (*Phylloxera vitifoliae*, Fitch = *Ph. vastatrix* Pl.); *a*, shows a healthy root; *b*, one on which the lice are working, representing the knots and swellings caused by their punctures; *c*, a root that has been deserted by them, and where the rootlets have commenced to decay; *d, d, d*, shows how the lice are found on the larger roots; *e*, female pupa, dorsal view; *f*, same, ventral view; *g*, winged female, dorsal view; *h*, same, ventral view; *i*, magnified antenna of winged insect; *j*, side view of the wingless female, laying eggs on roots; *k*, shows how the punctures of the lice cause the larger roots to rot.
- FIG. 29. Mulberry Silkworm (*Bombyx mori* Linn.), larva.
- FIG. 30. Mulberry Silkworm (*Bombyx mori* Linn.), cocoon.
- FIG. 31. Mulberry Silkworm (*Bombyx mori* Linn.), moth.
- FIG. 32. Mulberry Silkworm (*Bombyx mori* Linn.), cocoons; *a*, White French Annual; *b*, Yellow French Annual; *c*, Green Japanese Annual; *d*, White Japanese Annual; *e*, White Chinese Annual.
- FIG. 33. Cecropia Silkworm Moth (*Attacus Cecropia* Linn.).
- FIG. 34. Cecropia Silkworm (*Attacus Cecropia* Linn.), cocoon.
- FIG. 35. Cecropia Silkworm (*Attacus Cecropia* Linn.), chrysalis.
- FIG. 36. Cecropia Silkworm (*Attacus Cecropia* Linn.).
- FIG. 37. *Ophion macrurum*, Linn. [After Packard.]
- FIG. 38. *Ophion macrurum*, Linn., larva.
- FIG. 39. Mary Chalcis-fly (*Chalcis maria* Riley).
- FIG. 40. Cecropia Cryptus (*Cryptus samiae* Pack.), cocoons within the larger Cecropia cocoon.
- FIG. 41. Cecropia Cryptus (*Cryptus samiae* Pack.); *a*, female; *b*, female abdomen of *C. nuncius*; *c*, male abdomen; *d*, highly magnified piece of wing.
- FIG. 42. Ailanthus Silkworm (*Attacus cyntia*, Hübn.); 1, caterpillar; 2, moth; 3, cocoon; 4, chrysalis; 5, eggs.
- FIG. 43. Promethia Silkworm (*Attacus promethea* Drury); *a*, third stage; *b*, head in fourth stage, enlarged; *c*, lateral view of a joint in fourth stage, enlarged; *d*, full-grown caterpillar.
- FIG. 44. Promethia Silkworm (*Attacus promethea* Drury), cocoon.
- FIG. 45. Promethia Moth (*Attacus promethea* Drury), male. [After Harris.]
- FIG. 46. Promethia Moth (*Attacus promethea* Drury), female. [After Harris.]
- FIG. 47. Luna Moth (*Attacus Luna* Linn.). [After Harris.]
- FIG. 48. Luna Silkworm (*Attacus Luna* Linn.).
- FIG. 49. Luna Silkworm (*Attacus Luna* Linn.), cocoon. [After Harris.]
- FIG. 50. Polyphemus Moth (*Attacus Polyphemus* Linn.), male.
- FIG. 51. Polyphemus Moth (*Attacus Polyphemus* Linn.), female. [After Harris.]
- FIG. 52. Polyphemus Silkworm (*Attacus Polyphemus* Linn.). [After Trouvelot.]
- FIG. 53. Polyphemus Silkworm (*Attacus Polyphemus* Linn.), cocoon. [After Trouvelot.]
- FIG. 54. Polyphemus Silkworm (*Attacus Polyphemus* Linn.), chrysalis. [After Trouvelot.]
- FIG. 55. Yama-mai Moth (*Attacus yama-mai*, Guér.-Mén.), male.
- FIG. 56. Yama-mai Silkworm (*Attacus yama-mai*, Guér.-Mén.); egg, natural size and enlarged; young caterpillar on leaf; full grown caterpillar at rest on twig.
- FIG. 57. Yama-mai Silkworm (*Attacus yama-mai*, Guér.-Mén.), at rest on leafy twig, at *a*. [After Adams.]
- FIG. 58. Yama-mai Silkworm (*Attacus yama-mai*, Guér.-Mén.), cocoon.
- FIG. 59. Cage for receiving the deposition of the eggs of Yama-mai Moth. [After Adams.]
- FIG. 60. Pernyi Moth (*Attacus Pernyi*, Guér.-Mén.).
- FIG. 61. Pernyi Silkworm (*Attacus Pernyi*, Guér.-Mén.); egg, natural size and enlarged cocoon.

- FIG. 62. Horned Passalus (*Passalus cornutus* Fabr.); *a*, larva; *b*, pupa; *c*, beetle; *d*, under side of three thoracic joints of larva, showing legs; *e*, metathoracic leg of larva.
- FIG. 63. Great Leopard-moth (*Ecpantheria scribonia*, Stoll.), *a*, caterpillar; *b*, one hair, enlarged.
- FIG. 64. Great Leopard-moth (*Ecpantheria scribonia*, Stoll.); *a*, female; *b*, male.
- FIG. 65. Isabella Tiger-moth (*Arctia isabella*, Smith); *a*, caterpillar; *b*, chrysalis; *c*, moth.
- FIG. 66. Acorn-moth (*Holcocera glandulella* Riley); *a*, caterpillar in acorn; *b*, perforated acorn; *c*, head and thoracic joints of caterpillar, enlarged; *d*, *e*, lateral and dorsal views of one segment of larva; *f*, moth; *g*, base of antenna of male.

REPORT V.

- FIG. 1. Pyramid, showing the nature of the mouth, the relative rank of the Orders and the affinities of the Sub-orders of Insects.
- FIG. 2. Bald-faced Hornet (*Vespa maculata* Linn.). [After Sanborn.]
- FIG. 3. Goldsmith-beetle (*Cotalpa lanigera*, Linn.).
- FIG. 4. *Deïopeïa bella*, Drury.
- FIG. 5. Dotted-legged Plant-bug (*Euschistus punctipes*, Say).
- FIG. 6. Buffalo Tree-hopper (*Ceresa bubalus*, Fabr.); *a*, side view; *b*, view from above.
- FIG. 7. Missouri Bee-killer (*Asilus missouriensis* Riley).
- FIG. 8. Differential Locust (*Caloptenus differentialis* Walk.).
- FIG. 9. Dragon-fly (*Libellula trimaculata*, DeGeer.) [After Sanborn.]
- FIG. 10. Hull's Curculio-catcher.
- FIG. 11. Butterfly net; *b*, hinge in the ring; *c*, ring folded; *d*, nut sunk and soldered into brass tube at end of handle; *e*, screw; *f*, tip of handle, showing attachment of the ring.
- FIG. 12. Butterfly net; *a*, ring; *b*, socket; *c*, cork plug.
- FIG. 13. Butterfly net, head for attaching the ring to the rod.
- FIG. 14. Poison-bottle for killing insects; *a*, wadding to keep the cyanide grains in place.
- FIG. 15. Chloroform in stoppered bottle with brush.
- FIG. 16. Chloroform in bottle with tube passing through the cork.
- FIG. 17. Method of pinning insects; *a*, beetle; *b*, bug.
- FIG. 18. Method of carding small insects.
- FIG. 19. Method of "setting" Lepidoptera on a spreading board.
- FIG. 20. Setting-needle.
- FIG. 21. Sections of framework of glass-covered volume to display showy insects; *a*, ends; *b*, front; *c*, back.
- FIG. 22. Forceps for pinning insects.
- FIG. 23. Forceps for pinning insects.
- FIG. 24. Forceps for pinning insects.
- FIG. 25. Breeding-cage; *a*, bottom board; *b*, four-sided frame, with glass sides and door, fitting over a zinc pan (*ff*) attached to the bottom board; *c*, cover fitting to the frame and having a wire gauze top; *d*, zinc tube attached in centre of the pan, to contain a bottle for the reception of the food plant; *e*, sand in the pan; *gg*, cross pieces for supporting the cage and to prevent warping.
- FIG. 26. Ring-legged Pimpla (*Pimpla annulipes* Br.), female; to the right a figure of the ovipositor to show the two inner rods; to the left the abdomen of the male.
- FIG. 27. Delicate Longsting (*Macrocentrus delicatus* Cress.); to the right the abdomen of the male.
- FIG. 28. Rust-red Social Wasp (*Polistes rubiginosus* St. Farg.); *b*, nest, the natural position being with the mouths of the cells down.

- FIG. 29. Apple-tree Tent-caterpillar (*Clisiocampa americana* Harr.), eggs.
- FIG. 30. Grape Phylloxera (*Phylloxera vastatrix* Plan.); *a, b*, peculiar pedunculated galls; *c*, gall just forming; *d*, same from beneath.
- FIG. 31. Oyster-shell Bark-louse (*Mytilaspis pomicorticis* Riley); *a*, male louse from beneath; *b*, same from above and with wings expanded; *c*, male scale; *d*, leg of male; *e*, portion of wing very highly magnified; *f*, one joint of male antennæ (all highly magnified).
- FIG. 32. Oyster-shell Bark-louse (*Mytilaspis pomicorticis* Riley); anal joint of louse, with a more highly magnified segment of edge at *b*, and of a single pore at *c*; *d*, female louse; *e*, a section of its proboscis more highly magnified; *g h f*, female scale, *h*, first scale, *g*, second scale, *f*, third scale.
- FIG. 33. Mite (*Dermaleichus*?).
- FIG. 34. *Aphelinus mytilaspidis* LeBaron.
- FIG. 35. Pine-leaf Scale-insect (*Mytilaspis pinifoliae*, Fitch.); *a*, scales on leaves of white pine; *b*, male scale; *c*, female scale from white pine; *d*, female scale from broader leaved pine (*b, c* and *d*, enlarged).
- FIG. 36. Pine-leaf Scale-insect (*Mytilaspis pinifoliae*, Fitch); male, highly magnified.
- FIG. 37. Painted Ladybird (*Coccinella picta* Randall); *a*, larva; *b*, beetle; *c*, beetle, enlarged.
- FIG. 38. Hickory Bark-borer (*Scolytus caryæ* Riley); 1, view of its galleries on the inside of the bark, showing the beetle in the central gallery and the larvæ at the ends of the side galleries; 2, burrows made by larger larvæ; 3, beetle, magnified and natural size; 4, larva, magnified and natural size; 5, pupa, magnified; 6, sculpture of elytra, magnified.
- FIG. 39. Rose Chafer (*Macrodactylus subspinosus*, Fabr.), with the enlarged anterior tibia at the left.
- FIG. 40. Chinch-bug (*Micropus leucopterus*, Say).
- FIG. 41. False Chinch-bug (*Nysius destructor* Riley); *a*, potato leaf showing some effects of its punctures; *b*, pupa; *c*, mature bug.
- FIG. 42. Grape-vine Apple-gall (*Vitis-pomum* Walsh & Riley); *a*, exterior; *b*, section.
- FIG. 43. Gall-gnat (*Cecidomyia salicis-strobiloides* Walsh), *a*, female; *b*, male antennæ.
- FIG. 44. Grape-vine Filbert-gall (*Vitis-coryloides* W. & R.); *a*, anterior joints of larva, showing breast-bone; *b*, cluster of galls; *c*, section of single gall.
- FIG. 45. Grape-vine Tomato-gall (*Vitis-tomatos* Riley = *Lasioptera vitis* O. S., gall); *a*, section of a single swelling.
- FIG. 46. Grape-vine Trumpet-gall (*Vitis-viticola* Riley = *Cecidomyia viticola* O. S.)
- FIG. 47. Jumping Tree-cricket (*Orocharis saltator* Uhler) eggs in grape twig; *a*, eggs; *b*, punctures; *c*, egg, enlarged.
- FIG. 48. Jumping Tree-cricket (*Orocharis saltator* Uhler); *a*, female; *b*, male.
- FIG. 49. Snowy Tree-cricket (*Ecanthus niveus* Harr.) eggs; *a*, punctures in twig; *b*, section of twig showing the eggs within; *c*, egg, enlarged; *d*, granulations at rounded end of egg, more highly magnified.
- FIG. 50. Buffalo Tree-hopper (*Ceresa bubalus* Fabr.) eggs in slits in the bark of a tree; *a*, one slit enlarged; *b*, natural size.
- FIG. 51. Buffalo Tree-hopper (*Ceresa bubalus*, Fabr.); *a*, side; *b*, dorsal view.
- FIG. 52. Buffalo Tree-hopper (*Ceresa bubalus*, Fabr.); *a*, larva; *b*, pupa; *c*, ovipositor of the female, all enlarged.
- FIG. 53. Egg-punctures of Tree-hopper (?) on apple twigs; *a*, natural size; *b*, enlarged.
- FIG. 54. Frosted Lightning-hopper (*Pæciloptera pruinosa*, Say) eggs; *a*, enlarged; *b*, in position within twig, enlarged; *c*, natural size.
- FIG. 55. Frosted Lightning-hopper (*Pæciloptera pruinosa*, Say).
- FIG. 56. Egg-punctures of (?) *Orchelimum glaberimum* (Burm.).
- FIG. 57. Eggs of the Angular-winged Katydid (*Microcentrus retinervis*, Burm.); *a*, front; *b*, side view, just before hatching.

- FIG. 58. Eggs of the Angular-winged Katydid (*Microcentrus retinervis*, Burm.); *a*, front; *b*, side view, soon after laid.
- FIG. 59. Eggs of the Broad-winged Katydid (*Platyphyllum concavum* Harr.); *a*, side; *b*, front view, enlarged; *c*, *d*, natural size.
- FIG. 60. Buck Moth (*Hemileuca maia*, Drury).
- FIG. 61. Buck Moth (*Hemileuca maia*, Drury) eggs.
- FIG. 62. Buck Moth (*Hemileuca maia*, Drury); *a*, full-grown larva; *b*, pupa; *c*, ordinary form of spine of larva in the first stage; *d*, branched spine on thoracic joints of same; *e*, form of spines in second stage of larva; *f*, *g*, spines of full-grown larva.
- FIG. 63. Io Moth (*Hyperchiria Io*, Fabr.), male.
- FIG. 64. Io Moth (*Hyperchiria Io*, Fabr.), female.
- FIG. 65. Io Moth (*Hyperchiria Io*, Fabr.), caterpillar.
- FIG. 66. Io Moth (*Hyperchiria Io*, Fabr.), spines in 1st (*c*), 2d (*b*), and 5th (*a*) stages of caterpillar.
- FIG. 67. Green-striped Maple-worm (*Dryocampa rubicunda*, Fabr.); *a*, caterpillar; *b*, chrysalis; *c*, female moth.
- FIG. 68. *Belvoisia bifasciata*, Fabr.
- FIG. 69. Hellgrammite Fly (*Corydalis cornutus*, Linn.); *a*, larva; *b*, pupa; *c*, male fly; *d*, outline of head and prothorax of female.
- FIG. 70. Hellgrammite Fly (*Corydalis cornutus*, Linn.); supposed eggs.
- FIG. 71. Hellgrammite Fly (*Corydalis cornutus*, Linn.), pupa.
- FIG. 72. Goat-weed Butterfly (*Paphia glycerium* Doubl.); *a*, leaf eaten by the larva (natural size); *b*, head of larva in the first stage; *c*, larva in third stage; *d*, head in second stage; *e*, head in fourth stage—all enlarged.
- FIG. 73. Painted-wing Digger-wasp (*Ammophila pictipennis* Walsh).
- FIG. 74. Yucca-moth (*Pronuba yuccasella* Riley); *a*, head with pollen mass (1), (2) the maxillary tentacle, (3) the maxillae, (4) maxillary palpi, (5) antenna; *b*, maxillary palpi with tentacle; *c*, single spine from maxillary tentacle; *d*, maxillary palpus of male; *e*, wing scale; *f*, anterior leg; *g*, labial palpus; *h*, venation of anterior wing; *i*, venation of posterior wing, male; *j*, last joint of the abdomen of the female with the ovipositor exerted—all enlarged.
- FIG. 75. Yucca-moth (*Pronuba yuccasella* Riley); *a*, larva; *b*, moth with wings folded; *c*, female moth with wings expanded, (all natural size); *d*, side view of one joint of larva; *e*, head of larva from below; *f*, same from above; *g*, leg of larva; *h*, maxilla; *i*, mandible; *j*, labial palpi and spinneret; *k*, antenna—all enlarged.

REPORT VI.

- FIG. 1. Potato-beetle Catcher. Made of five barrel hoops and four (BB. EE) barrel staves, covered with cotton cloth.
- FIG. 2. Grape Phylloxera (*Phylloxera vastatrix* Planchon), galls on the leaf, seen from beneath.
- FIG. 3. Grape Phylloxera (*Phylloxera vastatrix* Plan.); *a*, *b*, pedunculated galls; *c*, gall just forming; *d*, same from beneath.
- FIG. 4. Grape Phylloxera (*Phylloxera vastatrix* Plan.)—Type Gallicola; *a*, *b*, newly-hatched larva, ventral and dorsal view; *c*, egg; *d*, section of gall; *e*, swelling of tendril; *f*, *g*, *h*, mother gall-louse—lateral, dorsal and ventral views; *i*, her antenna; *j*, her two-jointed tarsus.
- FIG. 5. Grape Phylloxera (*Phylloxera vastatrix* Plan.)—Type Radicicola; *a*, roots of Clinton vine, showing relation of swellings to leaf galls, and power of resisting decomposition; *b*, larva as it appears when hibernating; *c*, *d*, antenna and leg of same; *e*, *f*, *g*, forms of more mature lice; *h*, granulations of skin; *i*, tubercle; *j*, transverse folds at border of joints; *k*, simple eyes.

- FIG. 6. Grape Phylloxera (*Phylloxera vastatrix* Plan.)—Type Radicicola; *a*, shows a healthy root; *b*, one on which the lice are working, representing the knots and punctures caused by their punctures; *c*, a root that has been deserted by them, and where the rootlets have commenced to decay; *d, d, d*, show how the lice are found on the larger roots; *e*, female pupa, dorsal view; *h*, same, ventral view; *i*, magnified antenna of winged insect; *j*, side view of the wingless female, laying eggs on roots; *k*, shows how the punctures of the lice cause the larger roots to rot.
- FIG. 7. Grape Phylloxera (*Phylloxera vastatrix* Plan.). Pterogostic characters; *a, b*, different venation of front wing; *c*, hind wing; *d, e, f*, showing development of wings.
- FIG. 8. Grape Phylloxera (*Phylloxera vastatrix* Plan.)—Type Radicicola; *a, b*, pupa and imago of a problematical individual or supposed male; *c, d*, its antenna and leg; *e*, vesicles found in abdomen.
- FIG. 9. Thrips, enlarged, wings at right more highly enlarged.
- FIG. 10. Lace-wing fly (*Chrysopa* sp.); *a*, eggs; *b*, larva; *c*, cocoon, the upper figure with the lid open after the fly has escaped; *d*, fly, the wings omitted on the left. [*a, b, d*, after Westwood.]
- FIG. 11. Ladybird (*Hippodamia convergens* Gué.); larva, pupa and beetle.
- FIG. 12. Syrphus larva; *b*, one joint enlarged.
- FIG. 13. Syrphus-fly (*Helophilus latifrons* Loew).
- FIG. 14. Insidious Flower-bug (*Anthecoris insidiosus*, Say).
- FIG. 15. Root-louse Syrphus-fly (*Pipiza radicum* W. & R.); *a*, larva; *b*, pupa; *c*, fly.
- FIG. 16. Phylloxera Mite (*Tyroglyphus phylloxerae* Planchon & Riley); *a*, dorsal; *b*, ventral view of female; *c*, mouth parts; *d, f, g, h*, forms of tarsal appendages; *e*, ventral tubercles of male.
- FIG. 17. *Hoplophora arctata* Riley; *a, b, c, d, e*, different attitudes assumed by it; *f*, strongly magnified leg.
- FIG. 18. American Oak Phylloxera (*Phylloxera Rileyi* Lichtn.); *a*, pupa; *b*, winged females; *c*, antenna greatly enlarged; *d*, portion of infested leaf, under side.
- FIG. 19. American Oak Phylloxera (*Phylloxera Rileyi* Lichtn.); *a, b*, dorsal and ventral views of larva as seen hibernating; *c, d*, highly magnified leg and antenna of same.
- FIG. 20. Grape-vine Epimenis (*Psychomorpha epemenis*, Drury); *a*, larva; *b*, one joint, enlarged, side view; *c*, hump on joint 11.
- FIG. 21. Grape-vine Epimenis (*Psychomorpha epimenis*, Drury), male moth.
- FIG. 22. Beautiful Wood-nymph (*Eudryas grata*, Fabr.); *a*, full grown larva; *b*, one joint, enlarged, side view; *c*, cervical shield from behind; *d*, anal hump from behind; *e, f*, top and side views of egg.
- FIG. 23. Beautiful Wood-nymph (*Eudryas grata*, Fabr.), female moth.
- FIG. 24. Pearl Wood-nymph (*Eudryas unio*, Hübn.), male moth.
- FIG. 25. Eight-spotted Forester (*Alypia octomaculata*, Fabr.); *a*, larva; *b*, one joint, enlarged, side view; *c*, female moth.
- FIG. 26. Red-legged Ham-beetle (*Corynetes rufipes*, Fabr.); *a*, larva; *b*, pupa; *c*, cocoon; *d*, beetle, enlarged; *e*, same, natural size; *f*, leg of larva; *g*, mandible, *h*, labium, *i*, maxilla, *j*, antenna, of larva—all enlarged.
- FIG. 27. Larder-beetle (*Dermestes lardarius* Linn.); *a*, larva; *b*, one of its barbed hairs; *c*, beetle.
- FIG. 28. Clover-hay Worm (*Asopia costalis*, Fabr.); 1, 2, larva; 3, cocoon; 4, chrysalis; 5, 6, moth with wings expanded, and closed; 7, worm covered with silken web.
- FIG. 29. Legged Maple Borer (*Æjeria acerni*, Clem.); *a, a*, larva, dorsal and lateral views; *b, b, b*, cocoons exposed by detachment of bark; *c*, moth; *d*, chrysalis skin as it is often left remaining in the hole of exit.

- FIG. 30. Raspberry-root Borer (*Ægeria rubi* Riley); *a*, male moth; *b*, female moth.
- FIG. 31. Northern Brentian (*Eupsalis minuta*, Drury); *a*, larva; *b*, pupa; *c*, female beetle; *d*, head of male do.; *f*, leg of larva; *g*, head of larva, from in front; *h*, labium; *i*, labrum; *j*, mandible; *k*, maxilla; *l*, head from beneath, all of larva and enlarged; *m*, end of body of pupa, dorsal view.
- FIG. 32. Larva of Tenebrionid (?); *b*, front view of head; *c*, mandible; *f*, antenna; *g*, maxilla; *h*, labium; *d, e*, concave end of the body, full and side views.
- FIG. 33. Sumach Flea-beetle (*Blepharida rhois*, Forst.); *a*, egg; *b, b*, egg-masses, covered with excrement; *c, c, c, c*, larva; *d*, cocoon; *e*, pupa; *f*, beetle; *g*, antenna of larva; *h*, maxilla do.; *i*, mandible do.; *j*, labium do.; *k*, labrum do.; *l*, leg do.
- FIG. 34. *Tiphia inornata* Say; *a*, perfect wasp; *b*, head of larva, enlarged; *c*, larva, ventral view; *d*, cocoon cut open.
- Report VI, p. 122. Jiggers (*Leptus irritans* Riley, to the right; *L. americanus* Riley, to the left).
- FIG. 35. White-grub Fungus (*Torrubia ravenelii*, Berk.).
- FIG. 36. Dominican Case-bearer (*Coscinoptera dominicana*, Fabr.); *a*, larva extracted from case; *b*, do. with case; *c*, beetle, showing punctures; *d*, same, natural size; *e*, egg, enlarged; *i*, eggs, natural size; *g*, head of male beetle, enlarged; *h*, mandible of same, more enlarged; *j*, leg of larva, with the claw joint more enlarged; *f*, under side of larva; *k*, its mandible; *l*, maxilla, all enlarged.
- FIG. 37. *Chlamys plicata*, Oliv.; *a*, larva extracted from case, the figure at the right showing the larva in the case. [After Packard.]
- FIG. 38. Yucca-moth (*Pronuba yuccasella* Riley); *m*, female chrysalis; *l*, male chrysalis, the apical joints more highly enlarged and viewed from the side in lower figure.
- FIG. 39. Eyed Emperor (*Apatura lycaon*, Fabr.); *a*, eggs; *b*, larva; *c, d*, chrysalis, dorsal and lateral views; *e*, imago, male, the dotted line showing form of female — all natural size.
- FIG. 40. Eyed Emperor (*Apatura lycaon*, Fabr.); *f*, egg, magnified; *g*, larva, lateral view; *h*, imago, under side — natural size; *i, j, k, l, m*, the five different larval heads; *n, o*, dorsal and lateral views of one joint of larva — enlarged.
- FIG. 41. Tawny Emperor (*Apatura herse*, Fabr.); *a*, eggs; *b*, larva; *c*, chrysalis; *d*, imago, male, the dotted line showing form of female — all natural size.
- FIG. 42. Tawny Emperor (*Apatura herse*, Fabr.); *g*, larva, half grown, dorsal view; *h*, imago, male, under side — natural size; *i, j, k, l, m*, the five different heads of larva; *n, o*, dorsal and lateral views of one joint of larva; *p*, egg — enlarged; *g*, larvæ as when hibernating — natural size.
- FIG. 43. Eggs of the Angular-winged Katydid (*Microcentrus retinervis*, Burm.); *a*, front; *b*, side view, just before hatching.
- FIG. 44. Eggs of Angular-winged Katydid (*Microcentrus retinervis*, Burm.); *a*, front; *b*, side view, soon after laid.
- FIG. 45. Angular-winged Katydid (*Microcentrus retinervis*, Burm.); male wings closed.
- FIG. 46. Angular-winged Katydid (*Microcentrus retinervis*, Burm.); *a*, ovipositor of female, nat. size; *b*, tip of same, enlarged.
- FIG. 47. Angular-winged Katydid (*Microcentrus retinervis*, Burm.); female ovipositing.
- FIG. 48. Back-rolling Wonder (*Antigaster mirabilis* Walsh); *a*, female, wings expanded; *b*, same, side view, partly rolled up; *c*, same nearly rolled up; *d*, antenna of same.
- FIG. 49. Back-rolling Wonder (*Antigaster mirabilis* Walsh); *a*, eggs of *Microcentrus* from which it has issued; *b*, female pupa, ventral view; *c*, male fly; *d*, his antenna.
- FIG. 50. Narrow-winged Katydid (*Phaneroptera curvicauda*, DeGeer); female. [After Harris.]

- FIG. 51. Narrow-winged Katydid (*Phaneroptera curvicauda* DeGeer); *a*, ovipositor of female, nat. size; *d*, end of same, enlarged; *c*, anal appendage of male, side view; *b*, same, back view.
- FIG. 52. Broad-winged Katydid (*Platyphyllum concavum* Harr.); male (after Harris). [Adapted from Harris.]
- FIG. 53. Broad-winged Katydid (*Platyphyllum concavum* Harr.); *a*, ovipositor of female, nat. size; *b*, end of same, enlarged.
- FIG. 54. Eggs of Broad-winged Katydid (*Platyphyllum concavum* Harr.); *a*, side; *b*, front view—enlarged; *c*, *d*—natural size.
- FIG. 55. Oblong-winged Katydid (*Phylloptera oblongifolia*, DeGeer), outline of female [adapted from Harris]; *b*, end of ovipositor, enlarged.

REPORT VII.

- FIG. 1. Gray's Improved Sprinkler, for the use of Paris Green water. [From inventor.]
- FIG. 2. Chinch-bug (*Micropus leucopterus*, Say).
- FIG. 3. Chinch-bug (*Micropus leucopterus*, Say); *a*, *b*, eggs; *c*, newly hatched larva; *d*, its tarsus; *e*, larva after first molt; *f*, same after second molt; *g*, pupa, the natural sizes indicated at sides; *h*, enlarged leg of perfect bug; *j*, tarsus of same still more enlarged; *i*, proboscis or beak, enlarged.
- FIG. 4. Chinch-bug (*Micropus leucopterus*, Say), short-winged form.
- FIG. 5. Spotted Ladybird (*Hippodamia maculata*, DeGeer). [From *Practical Entomologist*.]
- FIG. 6. Trim Ladybird (*Coccinella munda* Say).
- FIG. 7. Insidious Flower-bug (*Anthocoris insidiosus*, Say).
- FIG. 8. Many-banded Robber (*Harpactor cinctus*, Fabr.); *a*, bug; *b*, its beak, enlarged.
- FIG. 9. False Chinch-bug (*Nysius destructor* Riley); *b*, pupa; *c*, mature bug.
- FIG. 10. Ash-gray Leaf-bug (*Piesma cinerea* Say).
- FIG. 11. Flea-like Negro-bug (*Corimelæna pulicaria*, Germar); natural size and enlarged.
- FIG. 12. Flat-headed Apple-tree Borer (*Chrysobothris femorata*, Fabr.); *a*, larva, dorsal view; *b*, pupa; *c*, swollen thoracic joints of larva from beneath; *d*, beetle.
- FIG. 13. Cherished Bracon (*Bracon charus* Riley).
- FIG. 14. Spring Canker-worm (*Anisopteryx vernata*, Peck); *a*, full grown larva; *b*, egg, enlarged, the natural size shown in the small mass at the side; *c*, *d*, one joint enlarged, side and dorsal views.
- FIG. 15. Spring Canker-worm (*Anisopteryx vernata*, Peck); *a*, male moth; *b*, female do.—natural size; *c*, joints of her antennæ; *d*, joint of her abdomen, showing spines; *e*, her ovipositor—enlarged.
- FIG. 16. Spring Canker-worm (*Anisopteryx vernata*, Peck); front view of head.
- FIG. 17. Fall Canker-worm (*Anisopteryx pometaria* Harr.); *a*, *b*, egg, side and top views; *c*, *d*, side and top views of one joint of larva,—enlarged; *e*, batch of eggs; *f*, full grown larva; *g*, female chrysalis—natural size; *h*, top view of anal tubercle of chrysalis.
- FIG. 18. Fall Canker-worm (*Anisopteryx pometaria* Harr.); *a*, male moth; *b*, female do.—natural size; *c*, joints of her antennæ; *d*, joint of her abdomen—enlarged.
- FIG. 19. Phylloxera, Male (*Phylloxera caryæcaulis*, Fitch?).
- FIG. 20. Grafting; *a*, *b*, incisions to receive the scion; *d*, scion; *c*, string to secure scion—to prevent phylloxera injury.
- FIG. 21. Grafting—to prevent phylloxera injury.
- FIG. 22. American Oak Phylloxera (*Phylloxera rileyi* Licht.); *a*, male, ventral view; *b*, genital organ; *c*, tarsus—all greatly enlarged.

- FIG. 23. Rocky Mountain Locust (*Caloptenus spretus* Thomas); *a, a, a*, female in different positions, ovipositing; *b*, egg-pod extracted from ground, with the end broken open, showing how the eggs are arranged; *c*, a few eggs lying loose on the ground; *d, e*, shows the earth partially removed, to illustrate an egg-mass already in place, and one being placed; *f*, shows where such a mass has been covered up.
- FIG. 24. Rocky Mountain Locust (*Caloptenus spretus* Thomas); anal characters of female, showing horny valves of ovipositor; *b*, an upper valve; *c*, a lower valve—all enlarged.
- FIG. 25. Rocky Mountain Locust (*Caloptenus spretus* Thomas); *a, a*, newly hatched larva; *b*, full grown larva; *c*, pupa.
- FIG. 26. Red-legged Locust (*Caloptenus femur-rubrum*, DeG.).
- FIG. 27. Rocky Mountain Locust (*Caloptenus spretus* Thomas).
- FIG. 28. Rocky Mountain Locust (*Caloptenus spretus* Thomas); *a*, tip of abdomen of male, side view; *b, c*, hind and top views of tip—all enlarged.
- FIG. 29. Red-legged locust (*Caloptenus femur-rubrum* DeGeer); *a*, tip of abdomen of male, side view; *b, c*, hind and top view—all enlarged.
- FIG. 30. Migratory Locust of Europe (*Ædipoda migratoria* Linn.).
- FIG. 31 (p. 142). Map of North America, illustrating the country east of the Rocky Mountains subject to the Ravages of the Rocky Mountain Locust.
- (Opposite p. 144.) Map of Missouri, illustrating the Locust Invasion of 1874.
- FIG. 32. Swarm of Locusts falling upon and devouring a wheat-field.
- FIG. 33. Differential Locust (*Caloptenus differentialis*, Walk.).
- FIG. 34. Two-striped Locust (*Caloptenus bivittatus*, Say).
- FIG. 35. Silky Mite (*Trombidium sericeum* Say); natural size shown at side.
- FIG. 36. Locust Mite (*Astoma gryllaria* LeBaron); greatly enlarged.
- FIG. 37. Mite parasitic on the House-fly (*Trombidium muscarum* Riley); enlarged.
- FIG. 38. Red-tailed Tachina-fly (*Exorista militaris* Kirkp.)
- FIG. 39. Flesh-fly (*Sarcophaga sarraceniæ* Riley); *a*, larva; *b*, pupa; *c*, fly; *d*, head and prothoracic joints of larva, showing curved hooks, lower lip (more enlarged at *g*), and prothoracic spiracles; *e*, end of body of larva, showing stigmata (more enlarged at *f*), prolegs and vent; *h*, tarsal claws of fly with protecting pads; *i*, antenna of fly—all enlarged.
- FIG. 40. Seventeen-year Locust (*Cicada septemdecim* Linn.); one wing removed so as to show ovipositor, *b*; *a*, beak.

REPORT VIII.

- FIG. 1. *Lebia grandis* Hentz.
- FIG. 2. Peck's Spray Machine in operation. [From inventor.]
- FIG. 3. Spring Canker-worm (*Paleacrita vernata*, Peck); *a*, caterpillar; *b*, eggs, natural size, one enlarged; *c*, one joint of larva, enlarged, side view; *d*, same, dorsal view.
- FIG. 4. Fall Canker-worm (*Anisopteryx pometaria* Harr.); *a, b*, egg enlarged, side and top views; *c, d*, joint of larva, enlarged, side and dorsal views; *e*, eggs, natural size; *f*, caterpillar; *g*, female chrysalis; *h*, tip of chrysalis, enlarged.
- FIG. 5. Spring Canker-worm (*Paleacrita vernata*, Peck), female chrysalis, enlarged.
- FIG. 6. Fall Canker-worm (*Anisopteryx pometaria* Harr.); *a*, male, *b*, female chrysalis, enlarged; a dorsal view of the tip of each, shown beneath.
- FIG. 7. Spring Canker-worm (*Paleacrita vernata*, Peck); *a, b*, venation of wings; *c*, one joint of male antennæ, greatly enlarged.
- FIG. 8. Fall Canker-worm (*Anisopteryx pometaria* Harr.); *a, b*, venation of wings; *c, d*, one joint of male antennæ, greatly enlarged, side and under views.
- FIG. 9. Spring Canker-worm (*Paleacrita vernata*, Peck); *a*, male moth; *b*, female moth—nat. size; *c*, portion of antenna of female; *d*, one segment of female abdomen; *e*, ovipositor—enlarged.

- FIG. 10. Fall Canker-worm (*Anisopteryx pometaria* Harr.); *a*, male moth; *b*, female moth—nat. size; *c*, joints of female antenna; *d*, one joint of female abdomen—enlarged.
- FIG. 11. Canker-worm Trap, consisting of a band of tin attached to a circle of muslin.
- FIG. 12. Canker-worm Trap, of tin and muslin; section. [From *Country Gentleman*.]
- FIG. 13. Canker-worm Trap, of tin and muslin; section to show the mode of union of the tin and muslin. [From *Country Gentleman*.]
- FIG. 14. Canker-worm Trap, at base of tree—Section. [From *Country Gentleman*.]
- FIG. 15. Canker-worm Trap, at base of tree. [From *Country Gentleman*.]
- FIG. 16. Tent-caterpillar of the Forest (*Clisiocampa sylvatica* Harr.); *a*, eggs; *b*, female moth; *c*, egg, enlarged, top view; *d*, same, side view.
- FIG. 17. Tent-caterpillar of the Forest (*Clisiocampa sylvatica* Harr.).
- FIG. 18. Army-worm (*Leucania unipuncta* Haw.), male genitalia; *A*, end of body denuded of hairs, showing the upper clasps protruding, and the natural position of the hidden organs by dotted lines; *B*, the organs extended; *c*, upper valves; *d*, lower valves; *e*, upper intermediate organ; *f*, penis; *g*, back view of upper intermediate organ; *h*, inner surface of upper valves—all enlarged.
- FIG. 19. Army-worm (*Leucania unipuncta* Haw.); *a*, *b*, end of abdomen of female denuded of scales, showing the ovipositor withdrawn and exerted; *c*, terminal joint of ovipositor; *d*, striations representing folds of the membrane, to facilitate expansion; *e*, *f*, retractile subjoints; *h*, eggs—all enlarged; *g*, eggs, natural size.
- FIG. 20. Army-worm (*Leucania unipuncta* Haw.), natural size when full grown.
- FIG. 21. Army-worm (*Leucania unipuncta* Haw.), chrysalis.
- FIG. 22. Army-worm (*Leucania unipuncta* Haw.); *a*, male moth; *b*, abdomen of female—nat. size; *c*, eye, *d*, base of male antenna; *e*, base of female antenna—enlarged.
- FIG. 23. Stalk-Borer (*Gortyna nitela* Guen.); *a*, terminal joints of female abdomen denuded to show the exerted ovipositor; *b*, view of the ovipositor from above.
- FIG. 24. Unarmed Rustic (*Agrotis saucia* Treit.); *a*, top view of egg, enlarged; *b*, batch of eggs enlarged. [See Notes, etc., p. 55.]
- FIG. 25. Unarmed Rustic (*Agrotis saucia* Treit.); *a*, ovipositor as it appears at the end of the abdomen; *b*, same when extended.
- FIG. 26. Fall Army-worm (*Laphygma frugiperda*, Sm. & Abb.); *a*, full grown worm, nat. size; *b*, head, front view; *c*, one joint of body, dorsal view; *d*, do., side view—enlarged. [See Notes, etc., p. 56.]
- FIG. 27. Fall Army-worm (*Laphygma frugiperda*, Sm. & Abb.); *a*, the typical form; *b*, *c*, variations of wings.
- FIG. 28. Elongate Ground-beetle (*Pasimachus elongatus* Lec.).
- FIG. 29. Murky Ground-beetle (*Harpalus caliginosus*, Fabr.).
- FIG. 30. Fiery Ground-beetle (*Calosoma calidum*, Fabr.); *a*, larva; *b*, beetle.
- FIG. 31. Rummaging Ground-beetle (*Calosoma scrutator*, Fabr.). [After Harris.]
- FIG. 32. Red-tailed Tachina-fly (*Exorista leucaniæ* Kirk.). [After Walsh.]
- FIG. 33. Yellow-tailed Tachina-fly (*Exorista flavicauda* Riley).
- FIG. 34. *Microgaster militaris* Walsh. [After Walsh.]
- FIG. 35. Glassy Mesochorus (*Mesochorus vitreus* Walsh). [After Walsh.]
- FIG. 36. *Pezomachus minimus* Walsh. [After Walsh.]
- FIG. 37. *Pezomachus minimus* Walsh, bunch of cocoons. [After Walsh.]
- FIG. 38. *Ophion purgatus* Say.
- FIG. 39. Rocky Mountain Locust (*Caloptenus spretus* Thomas): process of acquiring wings; *a*, pupa with skin just split on the back; *b*, the imago extending; *c*, do., nearly out; *d*, do. with wings expanded; *e*, do. with all parts perfect.
- FIG. 40. *Acridium americanum*, Drury.
- FIG. 41. Coral-winged Locust (*Edipoda phænicoptera* Germ.).

- FIG. 42. White-lined Morning Sphinx (*Deilephila lineata* Fabr.), green larva.
 FIG. 43. White-lined Morning Sphinx (*Deilephila lineata* Fabr.), black larva.
 FIG. 44. White-lined Morning Sphinx (*Deilephila lineata* Fabr.).
 FIG. 45. Lubber Locust (*Brachypeplus magnus* Gir.).
 FIG. 46. Green-striped Locust (*Tragocephala viridifasciata*); *a*, pupa; *b*, perfect insect.
 FIG. 47. Granulated Grouse-locust (*Tettix granulata* Scudder).
 FIG. 48. Grape Phylloxera (*Phylloxera vastatrix* Plan.); *a*, female, ventral view, showing egg through transparent skin; *b*, do. dorsal view; *c*, greatly enlarged tarsus; *d*, shrunken anal joints as they appear after oviposition; *e*, male of *Ph. caryæcaulis*, Fitch?, dorsal view—the dots in circle indicating natural size.
 FIG. 49. Yucca Borer (*Megathymus yuccæ*, Walk.); *a*, funnels made by the larva; *b*, under ground stem, showing tunnelings of larva.
 FIG. 50. Yucca Borer (*Megathymus yuccæ*, Walk.), female moth.
 FIG. 51. Yucca Borer (*Megathymus yuccæ*, Walk.); *a*, egg, side view, enlarged; *b*, egg from which the larva has hatched; *bb*, *bbb*, unhatched eggs—natural size; *c*, newly-hatched larva, enlarged; *c c*, full-grown larva, natural size; *d'*, underside of head of same, enlarged to show the trophi.
 FIG. 52. Yucca Borer (*Megathymus yuccæ*, Walk.), pupa.
 FIG. 53. Yucca Borer (*Megathymus yuccæ*, Walk.), moth walking.
 FIG. 54. Yucca Borer (*Megathymus yuccæ*, Walk.); *a*, *b*, venation of front and hind wings; *c*, labial palpus denuded; *d*, club of antenna; *e*, *f*, *g*, front, middle and hind legs,—all but wings enlarged.
 FIG. 55. *Castnia phalaris* (Fabr.), venation.

REPORT IX.

- FIG. 1. Gooseberry Span-worm (*Eufitchia ribearia*, Fitch.); *a*, *b*, larvæ; *c*, pupa.
 FIG. 2. Gooseberry Span-worm (*Eufitchia ribearia*, Fitch), female moth.
 FIG. 3. Gooseberry Span-worm (*Eufitchia ribearia*, Fitch); *a*, egg, enlarged; *b*, *b*, eggs, natural size.
 FIG. 4. Imported Currant-worm (*Nematus ventricosus* Klug); currant leaf showing eggs (1), and the holes which the young worms make (2, 3). [From *Practical Entomologist*.]
 FIG. 5. Imported Currant-worm (*Nematus ventricosus* Klug); *a*, *a*, *a*, larvæ; *b*, side view of one joint, enlarged, showing black tubercles.
 FIG. 6. Imported Currant-worm (*Nematus ventricosus* Klug); *a*, male fly; *b*, female fly.
 FIG. 7. Soldier-bug (*Podisus placidus* Uhler); *a*, enlarged; *b*, natural size.
 FIG. 8. Ovipositors of Sawflies; *a*, Willow-apple Sawfly (*Nematus salicis-pomum* Walsh); *b*, Currant-worm Sawfly (*Nematus ventricosus* Klug.), enlarged.
 FIG. 9. Native Currant-worm (*Pristiphora grossulariæ* Walsh); *a*, larva, nat. size; *b*, fly enlarged.
 FIG. 10. Strawberry-worm (*Emphytus maculatus* Nort.); 1, 2, ventral and lateral views of pupa; 3, enlarged sketch of perfect fly, the wings on one side detached; 4, larva crawling, natural size; 5, perfect fly with wings folded, natural size; 6, larva at rest; 7, cocoon; 8, antenna, enlarged; 9, egg, enlarged.
 FIG. 11. Abbot's Pine-worm (*Lophyrus Abbotii* Leach); 1, perfect fly, magnified; the left wings removed; 2, 3, ventral and lateral views of pupa, enlarged; 4, larvæ in different positions, nat. size; 5, cocoon, nat. size; 6, antenna of male, enlarged; 7, antenna of female, enlarged.
 FIG. 12. Map showing the distribution of the Colorado Potato-beetle (*Doryphora decemlineata*, Say).

- FIG. 13. *Uropoda americana* Riley; *a*, Colorado Potato-beetle attacked by it—nat. size; *b*, the mite, ventral view, showing the penetrating organ lying between the legs; *c*, the organs extended; *d*, the claw; *e*, the excrematitious filament—all greatly enlarged.
- FIG. 14. Wheat-head Army-worm (*Leucania albilinea* Guen.); *a*, *a*, larvæ; *b*, eggs—nat. size; *c*, *d*, egg, top and side view—enlarged.
- FIG. 15. Wheat-head Army-worm moth (*Leucania albilinea* Guen.).
- FIG. 16. Map of North America, illustrating the country east of the Rocky Mountains overrun by the Rocky Mountain Locust in 1876.
- FIG. 17. *Acridium americanum*, Drury.
- FIG. 18. Rocky Mountain Locust (*Caloptenus spretus* Thomas); *a*, *a*, *a*, female in different positions, ovipositing; *b*, egg-pod extracted from ground, with the end; *c*, a few eggs lying loose on the ground; *d*, *e*, shows the earth partially removed, to illustrate an egg-mass already in place and one being placed; *f*, shows where such a mass has been covered up.
- FIG. 19. Rocky Mountain Locust (*Caloptenus spretus* Thomas); Anal characters of female, showing horny valves of ovipositor; *b*, an upper valve; *c*, lower valve—all enlarged.
- FIG. 20. Rocky Mountain Locust (*Caloptenus spretus* Thomas); oviposition—*i*, superanal plate; *h*, sponge-like exsertile organ—the egg passing through the horny valves of the ovipositor, *g*.
- FIG. 21. Rocky Mountain Locust (*Caloptenus spretus* Thomas), egg-mass, enlarged; *a*, side view within burrow, the line of exit of the young locusts shown at *d* and *e*; *b*, egg-mass from beneath; *c*, same from above.
- FIG. 22. Rocky Mountain Locust (*Caloptenus spretus* Thomas); *a*, egg, enlarged to show sculpture of outer shell; *b*, portion of same very highly magnified; *c*, the inner shell, just before hatching; *d*, *e*, points where it ruptures.
- FIG. 23. Anthomyia Egg-parasite (*Anthomyia radicum*, Linn., var. *calopteni* Riley); fly; *b*, puparium; *c*, larva, side view; *d*, head of same, from above—enlarged.
- FIG. 24. Bombyliid larva (*Systæchus* sp.); *a*, enlarged; *b*, head, side view, more enlarged; *c*, do., front view; *d*, posterior spiracle. [See Notes, etc., p. 60.]
- FIG. 25. Harpalus? larva; *a*, from above; *b*, head, from beneath; *c*, leg—enlarged; *d*, antenna; *e*, maxilla; *f*, labium.
- FIG. 26. Harpalus? larva; A, natural size; B, under side of head, enlarged; *c*, mandible; *e*, antenna; *f*, labium and labial palpi; *g*, maxilla and maxillary palpi; *h*, joint 12 beneath; *i*, joint 11 beneath; *j*, joints 4–10 each beneath—enlarged.
- FIG. 27. Pennsylvania Ground-beetle (*Harpalus pensylvanicus*, DeGeer).
- FIG. 28. *Erax bastardi* Macq.; *a*, larva; *b*, pupa.
- FIG. 29. *Amblychila cylindriciformis* Say.
- FIG. 30. Hellgrammite (*Corydalis cornutus*, Linn.); *a*, larva; *b*, pupa; *c*, fly, male; *d*, head of female fly.
- FIG. 31. Hellgrammite (*Corydalis cornutus*, Linn.); *a*, *a*, egg-masses attached; *b*, one detached, showing lower surface,—all rather below average size; *c*, a few eggs of the outer row; *d*, the newly-hatched larva; *e*, labium; *f*, antenna; *g*, maxilla; *h*, mandible; *i*, tarsal claw; *j*, anal hooks—all enlarged.
- FIG. 32. Eggs of *Belostoma*?
- FIG. 33. Gigantic Water-bug (*Belostoma grandis* Linn.).

CLASSIFIED LIST OF ILLUSTRATIONS.

The following list of illustrations, brought together in classificatory order, will prove serviceable to entomologists, as it will enable such to readily ascertain whether or not any particular insect of a particular Order has been figured in the Reports. The explanations to the figures are omitted, since they are already given in the preceding list. The nomenclature of the Reports is here, also, retained, and references to figures other than those of insects or their products are omitted. The number of the Report is indicated in Roman and of the figure in Arabic numerals.

HYMENOPTERA.

- Ovipositors of Sawflies: IX, 8.
Pristiphora grossulariæ *Walsh*: IX, 9.
Nematus ventricosus *Klug*: IX, 4, 5, 6.
Emphytus maculatus *Nort.*: IX, 10.
Lophyrus Abbotii *Leach.*: IX, 11.
Aphelinus mytilaspidis *LeBaron*: V, 34.
Trichogramma ? *minuta* *Riley*: III, 72.
Antigaster mirabilis *Walsh*: VI, 48, 49.
Chalcid sp.: I, pl. 2, Fig. 6.
Glyphe viridascens *Walsh*: II, 24.
Isosoma vitis *Saunders*: I, 73; II, 64.
Isosoma hordei *Harr.*: II, 65.
Eurytoma bolteri *Riley*: I, pl. 2, Figs. 8, 9; I, 97.
Chalcis albifrons *Walsh*: II, 22.
Chalcis mariæ *Riley*: IV, 39.
Microgaster (= *Apanteles*): II, 48.
Microgaster cocoons on Hog-caterpillar of the Vine (*Chær. pampinatrix*, *Sm.* & *Abb.*): II, 47.
Microgaster militaris *Walsh*: I, 32; II, 23; III, 73; VIII, 34.
Aleiodes Rileyi *Cress.*: III, 30.
Bracon charus *Riley*: VII, 13.
Macrocentrus delicatus *Cress.*: V, 27.
Sigalphus curculionis *Fitch*: III, 7, 8.
Pimpla annulipes *Br.*: V, 26.
Cryptus samiæ *Pack.*: IV, 40, 41.
Hemiteles (?) *cressonii* *Riley*: I, pl. 2, Fig. 7.
Hemiteles (?) *thyridopterygis* *Riley*: I, pl. 2, Figs. 10, 11, 12.
Pezomachus minimus *Walsh*: II, 20, 21; VIII, 36, 37.
Porizon conotracheli *Riley*: III, 9.
Mesochorus vitreus *Walsh*: II, 19; VIII, 35.
Ophion macrurum (*Linn.*): IV, 37, 38.
Ophion purgatus *Say*: II, 25; VIII, 38.
Tiphia inornata *Say*: VI, 34.
Ammophila pictipennis *Walsh*: V, 73.
Stizus grandis *Say*, ♀: I, 12.
Eumenes fraterna *Say*: II, 71.
Polistes rubiginosus *St. Farg.*: V, 28.
Vespa maculata *Linn.*: V, 2.

COLEOPTERA.

- Amblychila cylindriciformis* *Say*: IX, 29.
Tetracha virginica *Hope*: I, 59.
Calosoma scrutator (*Fabr.*): II, 69; III, 54; VIII, 31.
Calosoma calidum (*Fabr.*): I, 34, 60; II, 70; VIII, 30.
Pasimachus elongatus *Lec.*: I, 61; VIII, 28.
Aspidoglossa subangulata *Chaud.*: I, 21.
Lebia grandis *Hentz*: III, 41; VIII, 1.
Harpalus caliginosus *Say*: I, 62; VIII, 29.
Harpalus pensylvanicus (*DeGeer*): I, 23; IX, 27.
Harpalus ? larva: I, 22; IX, 25, 26.
Quedius molochinus (*Grav.*): IV, 12.

- Gœrius olens*: IV, 11.
Philonthus apicalis (Say): IV, 10.
Dermestes lardarius Linn.: VI, 27.
Hippodamia maculata (DeGeer): I, 29; II, 3; VII, 5.
Hippodamia convergens Guer.: I, 52; VI, 11.
Hippodamia glacialis (Fabr.): IV, 5.
Hippodamia 13-punctata (Linn.): I, 51.
Coccinella 9-notata Herbst: I, 50.
Coccinella munda Say: II, 4; VII, 6.
Coccinella picta Randall: V, 37.
Mysia 15-punctata (Oliv.): IV, 4.
Chilocorus bivulnerus Muls.: I, 4, 5.
 Ladybird larva: I, 53.
Passalus cornutus Fabr.: IV, 62.
Macroductylus subspinosus (Fabr.): V, 39.
Lachnosterna quercina (Knoch): I, 88.
 White Grub attacked by fungus: I, 89.
Pelidnota punctata (Linn.): III, 34.
Cotalpa lanigera (Linn.): V, 3.
Chrysobothris femorata (Fabr.): I, 15, 16; VII, 12.
Chauliognathus pensylvanicus (DeGeer): I, 19; IV, 14.
Telephorus bilineatus (Say): IV, 15.
Corynetes rufipes (Fabr.): VI, 26.
Sinoxylon basilare (Say): IV, 26, 27.
Bostrichus bicaudatus (Say): IV, 24.
Orthosoma cylindricum (Fabr.): I, 69.
Prionus laticollis (Drury): I, 67; II, 60, 61, 62.
Prionus imbricornis (Linn.): II, 63.
Saperda bivittata Say: I, 14.
Bruchus pisi Linn.: III, 15, 16, 17.
Bruchus granarius Linn.: III, 18.
Bruchus fabæ Riley: III, 19.
Lema trilineata (Oliv.): I, 42, 43.
Coscinoptera dominicana (Fabr.): VI, 36.
Chlamys plicata (Oliv.): VI, 37.
Fidia viticida Walsh: I, 75.
Colaspis flavida Say: III, 37, 38; IV, 16.
Doryphora 10-lineata Say: I, 46.
Doryphora juncta (Germar): I, 47.
Diabrotica 12-punctata (Oliv.): II, 42.
Diabrotica vittata (Fabr.): I, 44; II, 39, 40, 41.
Haltica chalybea Illiger: III, 35, 36.
Haltica cucumeris Harr.: I, 45.
Blepharida rhois (Forst.): VI, 33.
Chelymorpha cribraria (Fabr.): II, 28, 29.
Physonota quinquepunctata Walsh & Riley: II, 30.
Cassida nigripes Oliv.: II, 37, 38.
Cassida bivittata Say: II, 27, 32.
Cassida aurichalcea (Fabr.): II, 31, 33, 34.
Cassida guttata Oliv.: II, 35, 36.
Deloyala clavata (Oliv.): II, 26.
 Larva of Tenebrionid (?): VI, 32.
Lytta cinerea Fabr.: I, 40.
Lytta vittata Fabr.: I, 39.
Lytta marginata Fabr.: I, 41.
Epicærus imbricatus (Say): III, 21.
Ithycerus novæboracensis (Forster): III, 20.
Anthonomus prunicida Walsh: III, 13.
Anthonomus quadrigibbus Say: III, 10, 11.
Conotrachelus nenuphar (Herbst): I, 18; III, 1.
Conotrachelus cratægi Walsh: III, 12.
Analcis fragariæ Riley: III, 14.
Cœliodes inæqualis (Say): I, 70, 71, 72.
Baridius trinotatus Say: I, 37.
Baridius sesostris Lec.: I, 74.
Eupsalis minuta (Drury): VI, 31.
Sphenophorus zeæ Walsh: III, 22.
Sphenophorus pulchellus Schœn.: III, 23.
Scolytus caryæ Riley: V, 38.

LEPIDOPTERA.

- Papilio philenor* Drury: II, 84, 85, 86.
Pieris protodice Boisd.: II, 72, 73, 74.
Pieris oleracea Boisd.: II, 75, 76.
Pieris rapæ Schrank.: II, 77, 78, 79.
Danaïs archippus (Fabr.): III, 63, 64, 65, 66, 67.
Limenitis disippus (Godt.): III, 68, 69, 70.
Apatura lycaon (Fabr.): VI, 39, 40.
Apatura herse (Fabr.): VI, 41, 42.
Paphia glycerium Doubl.: II, 94, 95, 96; V, 72.
Megathymus yuccæ Walk.: VIII, 51, 52, 53, 54.
Castnia phalaris (Fabr.) venation: VIII, 35.
Thyreus abboti Swainson: II, 54.
Deilephila lineata (Fabr.): III, 60, 61, 62; VIII, 42, 43, 44.
Chærocampa pampinatrix (Sm. & Abb.): II, 44, 45, 46.
Philampelus satellitia (Linn.): II, 52, 53.
Philampelus achemon (Drury): II, 49, 50, 51.
Sphinx 5-maculata Haw.: I, 38.
Ægeria exitiosa Say: I, 17.
Ægeria polistiformis Harr.: III, 33.
Ægeria rubi Riley: VI, 30.
Ægeria acerni (Clem.): VI, 29.
Alypia octomaculata (Fabr.): I, pl. 1, Fig. 18; II, 55; VI, 25.

- Psychomorpha epimenis* (*Drury*): I, pl. 1, Fig. 19; III, 26; VI, 20, 21.
Eudryas unio (*Hübner*): II, 57; VI, 24.
Eudryas grata (*Fabr.*): II, 56; VI, 22, 23.
Procris americana *Boisd.*: II, 58, 59.
Deiopeia bella (*Drury*): V, 4.
Callimorpha fulvicosta *Clem.*: III, 56.
Arctia isabella (*Smith*): IV, 65.
Spilosoma virginica (*Fabr.*): III, 28.
Hyphantria textor *Harr.*: III, 55.
Ecpantheria scribonia (*Stoll*): IV, 63, 64.
Orgyia leucostigma (*Sm. & Abb.*): I, 81, 82, 83.
Thyridopteryx ephemeræformis *Haw.*: I, 81.
Bombyx mori *Linn.*: IV, 29, 30, 31, 32.
Attacus polyphemus *Linn.*: IV, 50, 51, 52, 53, 54.
Attacus luna *Linn.*: IV, 47, 48, 49.
Attacus yama-mai (*Guér-Mén.*): IV, 55, 56, 57, 58.
Attacus Pernyi (*Guér-Mén.*): IV, 60, 61.
Attacus cynthia (*Hübner*): IV, 42.
Attacus promethea *Drury*: IV, 43, 44, 45, 46.
Attacus cecropia *Linn.*: IV, 33, 34, 35, 36.
Hemileuca maia (*Drury*): V, 60, 61, 62.
Hyperchiria io (*Fabr.*): V, 63, 64, 65, 66.
Dryocampa rubicunda (*Fabr.*): V, 67.
Clisiocampa americana *Harr.*: III, 50, 51; V, 29.
Clisiocampa sylvatica *Harr.*: II, 11; III, 52, 53; VIII, 16, 17.
Acronycta populi *Riley*: II, 87, 88.
Acronycta obliterata (*Sm. & Abb.*): III, 29.
Agrotis subgothica (*Haw.*): I, 29.
Agrotis jaculifera *Guen.*: I, pl. 1, Fig. 11.
Agrotis scandens *Riley*: I, pl. 1, Figs. 5, 6, 7.
Agrotis cochrani *Riley*: I, 26.
Agrotis inermis *Harr.*: I, pl. 1, Figs. 1, 2, 3, 4.
Agrotis sanciana *Treit.*: III, 49; VIII, 24, 25.
Agrotis telifera *Harr.*: I, pl. I, Figs. 8, 9, 10; I, 28.
Noctua clandestina *Harr.*: I, pl. I, Fig. 13; I, 27.
Hadena subjuncta *Gr. & Rob.*: I, pl. I, Figs. 14, 15, 16, 17.
Agrotis devastator (*Brace*): I, 30.
Pupa of Cut-worm in earthen cell: I, 25.
Mamestra picta *Harr.*: II, 82.
Celaena renigera *Steph.*: I, 83.
Prodenia commelinæ (*Sm. & Abb.*): I, pl. I, Fig. 12; III, 48.
Gortyna nitela *Guen.*: I, 35, 36; VIII, 23.
Leucania albilinea *Guen.*: IX, 14, 15.
Leucania unipuncta *Haw.*: II, 14, 15, 16; III, 47; VIII, 18, 19, 20, 21, 22.
Prodenia autumnalis *Riley*: III, 45, 46; VIII, 26, 27.
Amphipyra pyramidoides *Guen.*: III, 31, 32.
Anomis xyliana (*Say*): II, 12, 13.
Xyliana cinerea *Riley*: III, 57.
Plusia brassicæ *Riley*: II, 81.
Heliothis armigera *Hübner*: III, 42, 43.
Anisopteryx pometaria *Harr.*: VII, 17, 18; VIII, 4, 6, 8, 10.
Paleacrita vernata (*Peck*): VII, 14, 16; VIII, 3, 5, 7, 9.
Aplodes rubivora *Riley*: I, pl. 2, Fig. 25.
Hæmatopsis grataria (*Fabr.*): I, pl. 2, Figs. 18, 19, 20, 21.
Eufitchia ribearia (*Fitch*): IX, 1, 2, 3.
Asopia costalis (*Fabr.*): VI, 28.
Pempelia Hammondi *Riley*: IV, 21.
Pempelia grossulariæ *Pack.*: I, pl. 2, Fig. 17; I, 79.
Phycita nebulo *Walsh*: IV, 18, 19.
Acrobasis juglandis *LeBaron*: IV, 20.
Galleria cereana *Fabr.*: I, 92.
Desmia maculalis *Westw.*: III, 24.
Phacellura nitidalis *Cram.*: II, 43.
Tortrix cinderella *Riley*: IV, 22.
Tortrix rileyana *Grote*: I, pl. 2, Figs. 3, 4; I, 85.
Penthina vitivorana *Pack.*: I, pl. 2, Figs. 29, 30; I, 76.
Euryptychia saligneana *Clem.*: II, 99.
Carpocapsa pomonella (*Linn.*): I, 24.
Anchylopera fragariæ *Walsh & Riley*: I, pl. 2, Figs. 26, 27; I, 80.
Pronuba yuccasella *Riley*: V, 74, 75; VI, 38.
Walshia amorphella *Clem.*: II, 98.
Gelechia gallæsolidaginis *Riley*: I, pl. 2, Figs. 1, 2; I, 96.
Holcocera glandulella *Riley*: IV, 66.
Bucculatrix pomifoliella *Clem.*: IV, 23.
Pterophorus periscelidactylus *Fitch*: I, pl. 2, Figs. 15, 16; III, 27.
Pterophorus cardui *Riley*: I, pl. II, Figs. 13, 14; I, 98.

HETEROPTERA.

- Corimelæna pulicaria* (*Germar*): II, 9; VII, 11.
Stiretrus fimbriatus (*Say*): I, 5, 6; II, 10.
Perillus circumcinctus *Stål*: IV, 6.

Arma spinosa Dallas : I, 33, 54; II, 7; IV, 8, 9.
Podisus placidus Uhler : IX, 7.
Euschistus punctipes, Say : IV, 7; V, 5.
Strachia histrionica Hahn : IV, 17.
Coreus tristis (DeGeer) : I, 55.
Nysius destructor Riley : V, 41; VII, 9.
Micropus leucopterus (Say) : II, 1, 2; V, 40; VII, 2, 3, 4.
Campyloneura vitripennis (Say) : III, 58, 59.
Capsus oblineatus Say : II, 83.
Anthocoris insidiosus (Say) : II, 6; VI, 14; VII, 7.
Reduvius raptatorius Say : I, 58.
Harpactor cinctus (Fabr.) : I, 57; VII, 8.
Piesma cinerea (Say) : II, 8; VII, 10.
Belostoma grandis Linn. : IX, 33.

HOMOPTERA.

Cicada septemdecim Linn. : I, 6, 7, 13; VII, 40.
Cicada tredecim Linn. : I, 11.
Ceresa bubalus (Fabr.) : V, 6, 50, 51, 52.
Pæcilopectera pruinosa (Say) : V, 54, 55.
Pemphigus vagabundus (Walsh) : I, 65.
Eriosoma pyri Fitch : I, 64.
Phylloxera vastatrix Planchon : III, 39, 40; IV, 28; V, 30; VI, 2, 3, 4, 5, 6, 7, 8; VIII, 48.
Phylloxera caryæcaulis (Fitch)? : VII, 19.
Phylloxera rileyi Lichn. : VI, 18, 19; VII, 22.
Aspidiotus conchiformis (Gmêlin) : I, 2, 3; V, 31.
Mytilaspis pomicorticis Riley : I, 2, 3; V, 31, 32.
Aspidiotus harrisii (Walsh) : I, 1.
Mytilaspis pinifoliæ (Fitch) : V, 35, 36.

DIPTERA.

Gall, *Vitis-tomatos Riley*=*Lasioptera vitis O. S.* : V, 45.
Gall, *Vitis-viticola Riley*=*Cecidomyia viticola O. S.* : V, 46.
Gall, *Cecidomyia salicis-strobiloides Walsh* : V, 43.
Gall, *Vitis-coryloides W. & R.* : V, 44.
Gall, *Vitis-pomum W. & R.* : V, 42.
Tabanus atratus Fabr. : II, 97.
Wing of *Promachus (a)*, *Asilus (b)*, *Erax (c)* : II, 90.
Trupanea apivora Fitch : I, 93.
Erax bastardi Macq. : II, 92, 93; IX, 28.

Asilus missouriensis Riley : II, 89; V, 7.
Asilus sericeus Say : II, 91.
Bombyliid larva (*Systæchus sp.*) : IX, 24.
Pipiza radicum W. & R. : I, 66; VI, 15.
Syrphus larva : VI, 12.
Helophilus latifrons Loew : VI, 13.
Æstrus ovis Linn. : I, 91.
Exorista leucaniæ Kirk. : II, 17; VII, 38; VIII, 32.
Exorista flavicauda Riley : II, 18; VIII, 33.
Lydella doryphoræ Riley : I, 48.
Belvoisia bifasciata (Fabr.) : V, 68.
Sarcophaga sarraceniae Riley : VII, 39.
Anthomyia radicum (Linn.), var. *calopteni Riley* : IX, 23.
Anthomyia zeæ Riley : I, pl. II, Fig. 24; I, 86, 87.
Meromyza americana Fitch : I, pl. II, Fig. 28; I, 90.

ORTHOPTERA.

Æcanthus niveus Harr : I, 77, 78; V, 49.
Orocharis saltator Uhler : V, 47, 48.
Phaneroptera curvicauda (DeGeer) : VI, 50, 51.
Phylloptera oblongifolia (DeGeer) : VI, 55.
Microcentrus retinervis (Burm.) : V, 57, 58; VI, 43, 44, 45, 46, 47.
Platyphyllum concavum Harr : V, 59; VI, 52, 53, 54.
Orchelimum glaberrimum (Burm.) : V, 56.
Aceridium americanum (Drury) : VIII, 40; IX, 17.
Caloptenus spretus Thomas : VII, 23, 24, 25, 27, 28; VIII, 39; IX, 18, 19, 20, 21, 22.
Caloptenus femur-rubrum, DeG. : VII, 26, 29.
Caloptenus differentialis Walk. : V, 8; VII, 33.
Caloptenus bivittatus (Say) : VII, 34.
Brachypeplus magnus Gir. : VIII, 45.
Ædipoda migratoria Linn. : VII, 30.
Ædipoda phœnicoptera Germ. : VIII, 41.
Tragocephala viridifasciata Harr. : VIII, 46.
Tettix granulata Scudder : VIII, 47.
Mantis carolina (Linn.) : I, 94, 95.

NEUROPTERA.

Libellula trimaculata (DeGeer) : V, 9.
Corydalus cornutus (Linn.) : V, 69, 70, 71; IX, 30, 31.
Chrysopa sp. : I, 20; II, 5; VI, 10.
Thrips sp. : VI, 9.

ARACHNIDA.

Phalangium dorsatum Say : IV, 3.

ACARINA.

Trombidium muscarum Riley : VII, 37.

Trombidium sericeum Say : VII, 35.

Tyroglyphus phylloxeræ Planchon & Riley :
VI, 16.

Nothrus ovivorus Pack. : II, 68.

Leptus irritans Riley and L. americanus
Riley : VI, p. 122.

Uropoda americana Riley : IX, 13.

Dermaleichus? : V, 33.

Astoma gryllaria LeBaron : VII, 36.

Hoplophora arctata Riley : VI, 17.

GENERAL INDEX.

In this general index each report is referred to in Roman and the page in Arabic numerals. The index to the new matter of this Bulletin is also included and referred to by the abbreviation "Supp."

A.

Abbot Sphinx, II, 78
Abbotii, *Lophyrus*, IX, 32, Supp., 65
Thyreus, II, 78
 Abbots' White Pine Worm, IX, 29
 Descriptive, IX, 32.
 Natural History, IX, 30
 Natural Enemies, IX, 31
 Remedies, IX, 32
Abraxas grossulariata, IX, 5
Acarus mali, II, 6
 malus, I, 16, V, 87
 scabiei, VI, 61
 Walshii, V, 87
acericola, *Acronycta*, II, 121
acerni, *Aegeria*, VI, 107, 108, Supp., 55
 Trochilium, VI, 108
achemon, *Philampelus*, II, 74
 Achemon Sphinx, II, 74, 78
Achreioptera, a proposed order of insects, V, 16
Acidalia persimilata, VI, 138
Acoloithus falsarius, II, 86
 Acorn Moth, IV, 144
Acrididæ, Stridulation of, VI, 153
Acridii, VIII, 115, 128
Acridium americanum, VII, 173, 174, VIII, 103, 104, IX, 84
 peregrinum, VII, 133, VIII, 144, 145
 spretis, VII, 128
 spretum, VIII, 128
 Acridophagi, VIII, 144
Acrobasis, IV, 46
 consociella, IV, 45
 Hammondii, III, 7
 juglandis, IV, 42, 43, V, 49, Supp., 67, 80
 nebulo, IV, 38, 47, Supp., 79, 80
 nebulella, IV, 42, Supp., 80
Acronycta, II, 119
 americana, II, 121, Supp., 73, 74
 interrupta, II, 121, Supp., 73
 leporina, II, 121, Supp., 73
 lepusculina, II, 121, Supp., 73, 74
 leutiocoma, V, 126
 oblinita, III, 70, 71, V, 126
 occidentalis, II, 121, V, 126, Supp., 73
 populi, II, 119, 120, Supp., 72, 74
 psi, II, 121, Supp., 73
 tridens, II, 121, Supp., 73
 xylinoides, V, 126
acronyctæ, *Microgaster*, II, 120
 Act to provide for the destruction of Locusts in Minnesota, IX, 114

Act to encourage the destruction of Locusts in Missouri, IX, 111
 Acts to provide for the destruction of Locusts in Kansas, IX, 112, 113
Actias luna, IV, 123
Aculeata, a section of Hymenoptera, V, 9
 Address before the Farmers of Cass County, Missouri, VIII, 66
 Address before the National Agricultural Congress, VI, 17
 Adephaga, a section of carnivorous beetles, V, 11
 Adkins, F. D., Experience of, with Rocky Mountain Locust, VIII, 126
 Adkins, James, Report on Rocky Mountain Locust, IX, 74
Adoneta, VI, 140
 spinuloides, V, 126
adonidum, *Coccus*, III, 96
Ægeria, VI, 108
 acerni, VI, 107, Supp., 55
 cucurbitæ, II, 64
 exitiosa, I, 47
 polistiformis, I, 127, III, 75, 76
 rubi, VI, 111, 113, Supp., 72
 tipuliformis, II, 10, VI, 108, IX, 2
Ægeridæ, V, 41
Ægiale, VIII, 170
 indecisa, VIII, 179
 Kollari, VIII, 179
ænea, *Lytta*, III, 6
æqua, *Agrotis*, I, 74
Æschna, Oviposition of, VIII, 37
æscularia Anisopteryx, VIII, 17, Supp., 56
æstiva, *Dendroica*, VI, 27
 Affleck, Thos., on Cotton Worm, II, 38, 40, VI, 24
 Rocky Mountain Locust, VII, 139, 191
affusana, *Pædisca*, Supp., 57
 Agassiz, Professor, on mimicry, III, 73
Agelaius phœniceus, VI, 27
Ageratum, VI, 138
Aglaope americana, II, 85
 Agriculture, relation of insects to, V, 18
Agrion, oviposition of, VIII, 36
Agrotis, I, 68, Supp., 55
 æqua, I, 74
 Cochranii, I, 74, Supp., 76, 77
 cursoria, I, 78
 devastator, I, 83, Supp., 56
 herilis, Supp., 55
 inermis, I, 72, 74, II, 50, III, 15, 114, 129
 VII, 37, Supp., 55

- Agrotis jaculifera*, I, 82, 83, Supp., 56
lycarum, Supp., 77
maizi, I, 81
messoria, Supp., 77
murænula, I, 78
nigricans, I, 81, 83, 87
ortonii, Supp., 55
repentis, Supp., 77
saucia, I, 74, Supp., 55
scandens, I, 76, 78, III, 6, Supp., 75
subgothica, I, 81, 83, III, 151, Supp., 55, 56
suffusa, Supp., 55
telifera, I, 80, Supp., 55
triosa, Supp., 55
ypsilon, Supp., 55
Ailanthus Silkworm, IV, 112
 Best method of raising, IV, 119
 Difference between Castor bean and *Ailanthus* Worms, IV, 112
 Larval changes, IV, 117
 Natural History of, IV, 117
 Retrospective History of, IV, 113
 Thoroughly acclimated in America, IV, 115
 Value of the Cocoon, IV, 115
 When introduced into America, IV, 114
Ailanthus Worm, I, 151
alabamæ, *Chrysobothris*, VII, 71
Albany Argus, article from, on Army Worm, II, 43, VIII, 26
albifrons, *Chalcis*, II, 52, VIII, 54
albilinea, *Leucania*, IX, 55
albivenosus, *Micropus*, VII, 22
albolineata, *Synchlora*, Supp., 79
alboscuteclatus, *Bruchus*, Supp., 71
alceæ, *Erynnis*, VIII, 182
Aleochara anthomyiæ, IV, 22
Aleiodes Rileyi, III, 71
Aletia argillacea, VIII, 23, Supp., 56
alicia, *Apatura*, VI, 145, 150
Alkalies for Grape-vine Root-lice, IV, 69
Allen, T. R., Experience of, with Grass Cut-worm, I, 80
 on Army Worm, II, 47, VIII, 52
 Wheat Cut Worm, I, 87
Allen, G. W., on Rocky Mountain Locust, VIII, 102
Altica virginica, VI, 122
Alucitidæ, III, 67
Alypia octomaculata, I, 136, II, 80, 82, VI, 88, 94, 95
Amara angustata, VIII, 52
 obesa, Supp., 52
ambiguella, *Conchylis*, Supp., 57
Amblychila cylindriciformis, IX, 98
Ambulatoria, a division of Orthopterous insects, V, 14
americana, *Acronycta*, II, 121, Supp., 73, 74
 Aglaope, II, 85
 Apatela, II, 121, Supp., 73, 74
 Olisiocampa, II, 7, III, 117
 Clostera, II, 19, VII, 27
 Ctenucha, II, 85
 Meromyza, I, 59
 Parula, VI, 27
 Procris, II, 85
 Silpha, VI, 100
 Uropoda, IX, 41
American Acridium, VIII, 103, IX, 84
 Agriculturist, article from, on grass-hoppers, VII, 172
 article from, on Remedy for Currant Worm, IX, 15
 article from, on Yucca fertilization, V, 159
Bacon-beetle, VI, 100
Bean-weevil, III, 52
Blight, III, 95, IX, 43
Carion-beetle, VI, 100
Copper Underwing, III, 72
Cuckoos, III, 121
Entomologist, article from, on Bean weevil, III, 53
 article from, on Colorado Potato-beetle, III, 97
 article from, on Curculio extermination, III, 15
 article from, on imported and native insects, II, 8
 article from, on Strawberry Leaf-roller, I, 142
Meat Worm, IX, 43
Meromyza, I, 159, II, 16
Naturalist, article from, on Birds destroying Canker Worm, VI, 27
 article from, on injury caused by *Alypia*, II, 81
Oak Phylloxera, VI, 64, VII, 99, 118, VIII, 158
Plants and Insects acclimated in Europe, IX, 43
Procris, II, 85, 86, 87
Silk-worm, IV, 104
Tent-caterpillar, eggs of, V, 55
Timber-beetle, III, 7
vines, first suggestion to use in France as a remedy for Phylloxera, IV, 62
vines in France, VI, 79
americanum, *Acridium*, VII, 173, 174, VIII, 103, 104, IX, 84
americanus, *Chauliognathus*, Supp., 53
 Coccyzus, III, 121, VIII, 124
 Hylecætus, III, 7
Ammophila pictipennis, V, 149
amœnum, *Callidium*, IV, 54
amorphella, *Walshia*, II, 132, 133
Ampelis cedrorum, VI, 27
 garrulus, VII, 90
Ampelogypter sesostris, Supp., 71
ampelopsis, *Madarus*, I, 132
Amphicerus bicaudatus, IV, 51, V, 54
Amphipyra conspersa, III, 75, Supp., 75
 inornata, III, 75
 pyramidea, III, 73, 74
 pyramidoides, III, 72, 74, Supp., 75
Amphydasis cognataria, IX, 7
Amputating Brocade-moth, I, 87
amputatrix, *Hadena*, I, 87
Amydria, V, 151
Analcis, III, 44
 fragariæ, III, 42, 44, Supp., 71
Anaphora, V, 151

- Anasa*, Supp., 58
Anatis, Supp., 53
Anchylopera comptana, I, 143
 fragariae, I, 142, Supp., 57
 Andrews, W. V., on Eight-spotted Forester, II, 81
Angerona crocataria, IX, 7
 Angular-winged Katydid, VI, 155
 Description of immature Stages, VI, 161
 Natural Enemies, VI, 162
angulifera, *Callosamia*, IV, 122, 128, Supp., 55
 Angus, Jas., on Bean weevil, III, 52
angustata, *Amara*, VIII, 52
angustatus, *Nysius*, Supp., 85
angustifrons, *Anthomyia*, Supp., 89
 Animal Kingdom, Classification of, V, 6
Anisonyx rufa, VI, 143
Anisopteryx æscularia, VIII, 17, Supp., 56
 pometaria, II, 97, VI, 29, VII, 80, 83,
 86, VIII, 13, Supp., 56
 vernata, I, 109, II, 94, VI, 28, VII, 80,
 86
Anisota rubicunda, V, 137, 140
 stigma, V, 126
Anobium, III, 7, VI, 101
Anomalon apicale, IX, 55
 flavicorne, III, 69
Anomis, VI, 24
 xylina, II, 37, V, 19, VI, 17, VIII, 23,
 Supp., 56
anonyma, *Tachina*, IV, 129, V, 133, VII, 178, VIII,
 179
 Anonymous Tachina-fly, VII, 178
Annelida, V, 6, 7.
annulata, *Brochymena*, IV, 20
annulipes, *Pimpla*, IV, 43
Antheræa, IV, 114
 Pernyi, IV, 137
 Paphia, IV, 138
 yama-mai, IV, 130
Anthocoris, VII, 47
 insidiosus, II, 27, 32, VI, 51, VII, 41,
 Supp., 58
Anthomyia, III, 150
 angustifrons, Supp., 89
 brassicæ, I, 156, IV, 22, 35, IX, 95
 ceparum, I, 155, II, 9, IX, 95
 radicum, IX, 92, Supp., 88
 var. *calopteni*, IX, 92, Supp.,
 88
 raphani, IX, 95
 zææ, I, 154, Supp., 89
Anthomyia Egg-parasite of Locust, IX, 92
anthomyiæ, *Aleochara*, IV, 22
Anthonomus pomorum, III, 11
 prunicida, III, 39, Supp., 54
 pyri, III, 11
 quadrigibbus, III, 29, 35
 scutellaris, Supp., 54
 signatus, V, 154
 suturalis, III, 60
 Anthony, J. M., on Army Worm, VIII, 39
Anthribus varius, III, 10
 Antidote for Paris green, IV, 13
Antigaster mirabilis, VI, 162, 163, Supp., 52
 Ants, V, 9
Apanteles, Supp., 52, 66
Apatela americana, II, 121, Supp., 73, 74
Apatura alicia, VI, 145, 150
 celtis, VI, 139, 142, 150
 clyton, 142, 150
 herse, VI, 136, 137, 140, 148, 150
 idyja, VI, 145
 iris, VI, 136
 lycaon, VI, 136, 137, 146, 150
 proserpina, VI, 145, 150
Aphaniptera, V, 15
Aphelinus mytilaspidis, V, 88, 100
Aphidæ, V, 68, 85, VII, 27
 Aphidian Hickory galls, V, 154
Aphididæ, VI, 31, 33
Aphis, II, 19, III, 96, V, 149
 avenæ, II, 5, 6, 10
 brassicæ, II, 10, IV, 36
 mali, II, 6, 10, III, 6
 ribesii, VI, 46
 ribis, II, 10, IX, 2
 vitis, I, 13
Aphis, Currant, VI, 46
 Wooly, IV, 100
Aphodius, VI, 124
apiarius, *Clerus*, VI, 101
apicale, *Anomalon*, IX, 55
apicalis, *Philonthus*, IV, 21
Apiomerus, IX, 98
Apion apricans, III, 11
Apis mellifica, V, 18
apivora, *Trupanea*, I, 168, II, 122, Supp., 60
Aplodontia leporina, VI, 144
Aplodes rubivora, I, 139, Supp., 79
 Apple Bud-moth, III, 6
 Curculio, III, 6, 29, 30, 32, 33, 39
 Its natural history, III, 30
 It transforms in the fruit, III, 31
 Remedies, III, 34
 Season during which it works, III, 34
 The amount of damage it does, III, 33
 gall, Grape-vine, V, 114
 growing on Grape-vine, V, 115
 insects, III, 6
 leaf Bucculatrix, IV, 49
 remedies for, IV, 50
 Folder, The Lesser, IV, 47
 Skeletonizer, IV, 44, 47
 remedies for, IV, 45
 Tyer, IV, 46
 Maggot, III, 6, 9
 Maggot Fly, I, 108
 Micropteryx, III, 7
 Midge, III, 6
 Plant-louse, III, 6
 Root-lice, III, 95, IV, 68, 69
 Plant-louse, I, 118
 Artificial remedies for, I, 123
 Natural remedies for, I, 121
 Syrphus fly, I, 121
 tree Bark-louse, I, 7, III, 93
 borer, Flat-headed, I, 46, VII, 71
 Remedies for, I, 47
 Round-headed, I, 42, IV, 124
 VII, 27
 borer, Remedies for, I, 45
 Borers, I, 42

- Apple-tree Plant-louse, II, 6, 10
 Root-louse, II, 15, VI, 52
 Tent-caterpillar, II, 11, III, 117, 121
 twig Borer, IV, 51, V, 54
 Worm, I, 62, II, 6, 19, III, 6, 33, 90, 102, VII, 27
 Again, IV, 22
 Attacks peaches, V, 49
 False doctrines about, V, 51
 In California, V, 49
 Natural Enemies, V, 49
 Parasites, V, 49
 Remedies, I, 65
 Traps, V, 46
apricans, *Apion*, III, 11
apterus, *Micropus*, VII, 22
aquana (*Spilonota*), Supp., 57
Arachnida, V, 6, 7
archippivora, *Tachina*, III, 116, 150
Archippus, III, 168, 169, V, 148
 Butterfly, II, 125, III, 143
 How the larva becomes a chrysalis, III, 146
 It often congregates in immense swarms, III, 151
 Its natural history, III, 143
 The larva enjoys immunity from the attack of predaceous animals, III, 148.
 Tachina-fly, III, 150
archippus, *Danaus*, III, 143, 167, IV, 129, Supp., 55
arctata, *Hoplophora*, VI, 53, 81, VII, 106
Arctia Isabella, IV, 143, Supp., 55
Arctomys rufa, VI, 143
arcuata, *Ortalis*, II, 9
argillacea, *Aletia*, VIII, 23, Supp., 56
Argynnis, III, 103
 diana, III, 169, 171
 Arkansas, Locusts in, IX, 76
Arma modesta, V, 133
 spinosa, I, 77, 89, 113, II, 32, IV, 19, V, 51, Supp., 58
armigera, *Heliothis*, III, 45, 104, IV, 129
 Arment, A. B., on Rocky Mountain Locust, IX, 119
 Arnott, M. A., on Rocky Mountain Locust, IX, 117
 Arrenotoky in the Imported Currant-worm, IX, 18
Arrhenodes septentrionis, VI, 116
 Arsenious acid for Potato-bugs, IV, 14
arthemis, *Limenitis*, III, 171
Arthropoda, V, 6
Articulata, V, 6
 Army Grasshopper, VII, 194
 Worm, I, 89, 109, II, 5, 37, 70, 103, 110, III, 110, 125, 128, V, 22, 25, 68, VIII, 22, 182, IX, 49
 Additional Notes on Mode, Place, and Time of Oviposition, VIII, 182.
 Are there one or two Broods each year?, VIII, 47
 Completion of its Natural History, VIII, 182
 Correspondents quoted, III, 109, VIII, 39
 Description of the Egg, VIII, 34
 Larval Stages, VIII, 184
 Fall Army-worm, VI, 17, VIII, 23, 35, 49
 Army Worm, Further Notes and Experiments on, IX, 47
 Habits of the Worm, VIII, 45
 Ichneumon Fly parasitic, on II, 53, 54
 Its History in 1875, VIII, 28
 Missouri in 1875, VIII,
 Its Sudden Appearance and Disappearance, II, 45, VIII, 50
 Microgaster parasite of, III, 158
 Natural Enemies, VIII, 52
 History, II, 47, VIII, 32
 Nomenclature, VIII, 22
 Northern, VIII, 24
 Parasites, II, 50
 Parent Moth, II, 11, 48
 Past History, II, 41, VIII, 24
 Plants Preferred by the, VIII, 49
 Remedies, VIII, 54
 Sexual Differences, VIII, 30
 Summary of Natural History, VIII, 56, IX, 49
 Tachina-Fly parasitic on, IV, 109, VIII, 53
 the term "Army Worm" applied to various Insects, VIII, 23
 time of Appearance, VIII, 46
 Wheat-head Army-worm, IX, 50
 When are the Eggs laid?, VIII, 40
 Where are the Eggs laid?, VIII, 38
 Army Worms, The three, II, 37
 Ashes and air-slacked lime for potato-bugs, IV 14
 Ash-gray Blister-beetle, I, 97, 115
 Leaf-bug, II, 32, VII, 47
 Pinion, III, 134
 Asilus-fly, II, 123, III, 161
Asilus missouriensis, II, 121, 122, IV, 21, V, 13, Supp., 87
 sericeus, II, 123
Asopia costalis, VI, 102, 106
 olinalis, VI, 103, 107
asparagi, *Crioceris*, II, 10, 13, 19, VII, 5
 Asparagus Beetle, II, 10, 13, VII, 5
Aspidoglossa subangulata, I, 58
Aspidiotus, I, 14
 conchiformis, I, 7, II, 9, 10, V, 91, 94, Supp., 86
 harrisii, I, 7, II, 9, Supp., 60
 pinifoliae, III, 92
asterias, *Papilio*, III, 169
Astoma, VI, 52
 gryllaria, VII, 175, Supp., 63
 parasiticum, VII, 176
Astyci, VIII, 176
Atchison (Kans.) *Champion*, article from, on Rocky Mountain Locust, VIII, 108
atalanta, *Pyrameis*, III, 167
atlanis, *Caloptenus*, VII, 169, VIII, 113, 114, 115, 116, 117, 118, 153, Supp., 89
 Atlantic Locust, VIII, 150
 Atomizer for applying Paris green water, VI, 20, VIII, 5
atrata, *Lytta*, I, 98, Supp., 54
atratus, *Tabanus*, II, 128, 129, 130
atricapillus, *Parus*, IV, 107, VI, 27
atriventris, *Lebia*, VIII, 3, Supp., 52
atrox, *Oedipoda*, VII, 124

- Attacus cecropia*, III, 129, 170, IV, 103, 138
cynthia, III, 170, IV, 112, 121, 138
luna, IV, 123, 138
mylitta, IV, 138
pernyi, IV, 137, 138
polyphemus, III, 170, IV, 110, 125, 138
promethea, IV, 110, 121, 138
selene, IV, 125
yama-mai, IV, 130, 138
 Aughey, Prof. Saml., on Rocky Mountain Locust, VIII, 114
aurichalcea, *Cassida*, II, 62, Supp., 53
 Ooptocycla, Supp., 53
auricincta, *Tachina*, V, 140
aurocapillus, *Seiurus*, VI, 27
Aurocorisa, a group of Heteroptera, V, 12
autumnalis, *Prodenia*, III, 13, 109, 116, IV, 129, VIII, 48
avenæ, *Aphis*, II, 5, 6, 10
 Avery, Wm. H., on Army Worm, VIII, 39
 Chinch Bug, VII, 44
 Rocky Mountain Locust, IX, 69
 Ayres, E. J., on Tarnished Plant-bug, II, 114

B.

- bacchus*, *Rhynchites*, III, 11
 Backbone animals, V, 6
 Bacon-beetle, VI, 100
 Bag-worm, I, 147, III, 160, IX, 17
 parasite, I, 150
 Bailey, S. S., on Army Worm, VIII, 39
Balaninus, III, 10
 cerasorum, III, 11
 nucum, III, 11
 rectus, IV, 144
Baltimore American, article from, on Wheat-head Army-worm, IX, 51
baltimore, *Icterus*, VI, 27
 Baltimore Oriole destroying Canker Worm, VI, 27
 Katydid, VI, 162
 Pea-weevil, III, 50
Banchus fugitivus, IV, 41
 Bandage for Apple Worm, IV, 23
 Canker Worm, VI, 26, 27
 Banded Borer, III, 7
barbara, *Colaspis*, III, 82
 Barber, A. W., on White Pine-worm, IX, 30
Barbicornis, VIII, 170
 basalis, VIII, 170
Baridius sesostris, III, 60, Supp., 71
 trinotatus, I, 93, III, 60, Supp., 54
Baris chlorizans, III, 11
 Bark-beetles, III, 6
 -borer of hickory, V, 103
 -lice, II, 15, 25, III, 6, 10, V, 16, VI, 33
 classification of, V, 92
 of the Apple-tree, I, 7
 -louse, III, 85, V, 18
 on Apple, I, 7, V, 73
 remedies for, I, 16
 Currant, I, 15
 Pear, I, 15
 Persian lilac, I, 15
 Pine, V, 97
 Plum, I, 15

- Barker's Canker Worm trap, VIII, 21
 Barnet, W. N., on Phylloxera, VI, 82
 Barret, W. H., on Rocky Mountain Locust, VIII, 65
 Barron, W. H., on Rocky Mountain Locust, IX, 70
 Barter, A., on Grape-root Borer, I, 125
 Bartlett, Dr. L., on Army Worm, II, 49
basalis, *Micropus*, VII, 22
basillare, *Sinoxylon*, IV, 52, 53, 54
 Basket-worm, I, 147
 Bassett, S. C., on Rocky Mountain Locust, IX, 117
bassiana, *Botrytis*, IV, 88
bastardi, *Erax*, II, 124
 Promachus, II, 122, IV, 21, Supp., 60
 Bates, H. W., on Danais Butterflies, III, 161
 Bat-ticks, V, 13
 Baxter, C., on Phylloxera, VI, 83
 Bazille, L., on Phylloxera in France, VII, 104
 Beach, H. P., on Locusts, VIII, 152
 Beal, M., on Chinch Bug, II, 17
 Bean-weevil, III, 45, 52, 54, 55
 Beautiful Wood Nymph, II, 83, 84, III, 64, VI, 88, 91, 95
 Bed Bug, II, 15, 31, V, 12
 Beech-twigg Plant-louse, I, 121
 Bee-fly larva, Supp., 60
 Bee, Honey, IV, 84
 Bee-killer, I, 168, II, 121, IV, 21
 moth, I, 166, II, 10, III, 68
 parasite, V, 15
 Bees, V, 9
 Beetles, preparing of, for cabinet, V, 34
 Bell & Gruelle on Rocky Mountain Locust, IX, 92
Belostoma grandis, IX, 128
 probable eggs of, IX, 128
 Belt, J., on Rocky Mountain Locust, IX, 73
Belvoisia bicincta, V, 140
 bifasciata, V, 140
Bembex fasciata, V, 9
 Beneficial insects, I, 169, III, 137, IV, 72, VI, 123
 Bennet, W. H., on Mimicry, III, 172
 Benson's machine for catching Potato-beetle, I, 117
 Berckmans, P. J., on Phylloxera, VII, 103, VIII, 164, 165
berenice, *Danaïis*, III, 143
 Bessey, Prof. C. E., on Colorado Potato-beetle, V, 54
 Common Flesh-fly, VII, 180
 Locust Mite, VII, 175
 Strawberry Worm, IX, 27
bethunei, *Xylina*, III, 136, Supp., 75
betuleti, *Rhynchites*, III, 11
bicarinata, *Polysphincta*, III, 71
bicaudatus, *Amphicerus*, IV, 51, V, 54
 Bostrichus, III, 6, IV, 51, 53
 Bichromate of potash for Potato bugs, IV, 14
bicincta, *Belvoisia*, V, 140
 Senometopia, V, 140
bicolor, *Botys*, III, 61
bifasciata, *Belvoisia*, V, 140
 Musca, V, 140
bilineatus, *Ophion*, III, 69
 Telephorus, IV, 29, 30
bipunctata, *Xabea*, Supp., 61
bipunctatus, *Æcanthus*, Supp., 61
 Birds that destroy the Canker Worm, VI, 27, 28
 Birds vs. insects, III, 169, VI, 29

- Bismarck Tribune*, article from, on Rocky Mountain Locust, IX, 59
- bistriga*, *Cryptoblabes*, IV, 46, Supp., 81
- Bi-sulphide of carbon for Phylloxera, VI, 56
- bivittata*, *Cassida*, II, 61, Supp., 53
Saperda, I, 42, II, 19, III, 6, VII, 27, Supp., 53
- bivittatus*, *Caloptenus*, VII, 124, 173
- Bivoltin Silkworms, IV, 85
- bivulnerus*, *Chilocorus*, I, 16, V, 100
- Black-bear caterpillar, The Large, IV, 141
 -bellied Lebia, VIII, 3
- Blackberry fruit-worm, I, 139
- Blackbird destroying Katydid, VI, 162
 , Red-winged, destroying Chinch Bug, VII, 41
- Black Blister-beetle, I, 98
 Breeze-fly, II, 128, 129, 132
 larva of White-lined Morning Sphinx, III, 142, VIII, 122
 -legged Tortoise-beetle, II, 63
 -rat Blister-beetle, I, 98
 Stinger of the Oak, V, 126
- Blanchard, Stephen, on Rocky Mountain Locust, VII, 139
- Blatta orientalis*, II, 10
- Blepharida rhois*, I, 100, II, 58, VI, 118, 121
- Bliss, N. W., on Hickory Bark-borer, V, 103
- Blissus*, Supp., 58
- Blister-beetle, The Ash-gray, I, 97, 115
 Black, I, 98
 Black-rat, I, 98
 Brazen, III, 6
 Margined, I, 98
 Striped, I, 96, 115
- beetles, I, 115, Supp., 54
 remedies for, I, 99
- Blow-fly, II, 19, VII, 27
- Blue-bird destroying Canker Worm, VI, 27, 28
 Codling Moth, IV, 28
- Blue Caterpillars of the Vine, I, 136, II, 79, III, 63, 65, VI, 87
 Jay hiding corn in cocoons, IV, 107
 -spangled Peach-worm, III, 132
 Stone for Potato-beetles, IV, 14
- Boardman, S. P., on Sheep Bot-fly, I, 164
- Bob-o-link destroying Canker Worm, VI, 27
- Bogus Chinch Bug, II, 31, V, 112, VII, 45
 Colorado Potato-beetle, I, 105
- Boissière, E. V., on Silkworm raising, IV, 82
- Boissière's silk establishment, IV, 82
- Boll-worm, III, 45, 104, 105, VI, 20
 remedies for, III, 108
- Bolteri*, *Eurytoma*, I, 177, Supp., 68
- Bombycidae*, IV, 85, V, 127
- Bombyx*, IV, 114
graminis, II, 44
mori, IV, 75, 84, 183
- Book-lice, V, 15
- Bordered Soldier-bug, I, 114, II, 34
- borealis*, *Epilachna*, IV, 18
- Borer, The Apple-twigg, IV, 51, V, 54
 Flat-headed Apple-tree, I, 46, VII, 71
 Grape-root, III, 75
 Hickory Bark, V, 103
 Legged Maple, VI, 107
- Borer, The New Grape-root, I, 124, II, 87
 Peach, I, 47
 Potato Stalk, I, 92
 Raspberry Root, VI, 111
 Round-headed Apple-tree, I, 42
 Squash, II, 64
 Strawberry Crown, III, 42
 Yucca, VIII, 169, IX, 129
- Boston Daily Advertiser*, note from, on Army Worm, VIII, 29
- Boston Journal*, article from, on Canker Worm, II, 96
 Rocky Mountain Locust, III, 15
- Bostrichidae*, IV, 53
- Bostrichus*, IV, 53
bicaudatus, III, 6, IV, 51, 53
- botrana*, *Eudemis*, Supp., 57
Lobesia, Supp., 57
Tortrix, Supp., 57
- Botrytis bassiana*, IV, 88
viticola, VI, 36
- Bottom, R., on Rocky Mountain Locust, VIII, 62
- Botys bicolor*, III, 61
- bovinus*, *Tabanus*, II, 129
- Bowen, H., on destroying Potato-beetle, VI, 14
- Bowles, G. J., on imported Cabbage Butterfly, II, 107
- Box-turtle, IX, 98
- Boxes for insects, V, 37
- Brachinus*, Supp., 59
kansanus, IV, 21
- Brachista*, VI, 142
- Brachocera*, a division of Diptera, V, 13
- Brachypeplus magnus*, VIII, 148
- Brachypterus micropterus*, IX, 17
- Brachyscelides*, V, 92
- Bracket, G. C., on remedies for Canker Worm, II, 100
 Strawberry Crown-borer, III, 42
 Tent-caterpillar of the Forest, III, 127
- Bracon charus*, VII, 75, Supp., 67
scolytivorus, V, 106, Supp., 67
- Braconides*, III, 27, V, 106
polymorphi, I, 96
- brassicæ*, *Anthomyia*, I, 156, IV, 22, 35
Aphis, II, 10, IV, 36
Pieris, III, 167
Plusia, II, 110, 111
- Braula cæca*, V, 74
- Brazen Blister-beetle, III, 6
- Breast-bone in Cecidomyid larvæ, V, 114.
- Breck, Joseph, on remedies for Canker Worm, II, 100
- bredowii*, *Limenitis*, III, 171
- Breeding Insects, V, 41
- Breeze-fly, V, 13, VI, 123
- Brenthian, Northern, VI, 113
- Brenthus*, III, 10
maxillosus, VI, 116
septentrionis, VI, 116
- brevis*, *Phygadeuon*, IV, 28
- brevipennis*, *Eudryas*, VI, 91
- Brewer, F. A., on Phylloxera, V, 73
- Brimstone for Bark-lice, I, 17
- Briggs, A. A., on Flat-headed Borer, VII, 79

- Brinkerhoff, M., on Locusts, VIII, 154
 British America, Rocky Mountain Locust in, IX, 59
 Broadhead, G. C., on Pickle Worm, II, 69
 Rocky Mountain Locust, IX, 74
 Broad-necked Prionus II, 9, 87, 88, 89, V, 56
 -winged Katydid, V, 123, VI, 167
Brochymena annulata, IV, 20
 Brous, H. A., on Rocky Mountain Locust, IX, 66, 98, 99
 Brown Colaspis, III, 82
 Brown, A. M., on Apple Curculio, III, 34
 Brown, E. R., on Army Worm, VIII, 39
 Brown, L. A., on Army Worm, VIII, 39
 Brown, Major, on Colorado Potato-beetle, VII, 6
Bruchi, III, 51
Bruchides, III, 45
Bruchus alboscuteUellatus, Supp., 71
 discoideus, III, 45
 erythrocerus, III, 55, 56, Supp., 70
 fabæ, III, 52, 55, Supp., 69, 70
 flavimanus, III, 56, Supp., 70
 granarius, II, 11, 14, III, 50, 51
 hibisci, Supp., 70, 71
 obsoletus, III, 54, 56, Supp., 70, 71
 pisi, II, 11, III, 44, IX, 43, Supp., 53
 pisorum, Supp., 53, 71
 rufimanus, III, 56, Supp., 70
 serratus, III, 56, Supp., 70
 transversus, Supp., 70
 varicornis, III, 55, 56, Supp., 69, 70, 71
 Bruhl, Henry, on Army Worm, VIII, 39
 Bruner, Uriah, on Locusts, VII, 139
brunnea, *Colaspis*, III, 82
 Bryant, Arthur, on Hickory Bark-borer, V, 104
 Bryning, J. J., on Rocky Mountain Locust, IX, 69
bubalus, *Ceresa*, V, 12, 121
Bucculatrix pomifoliella, IV, 49, 51
 thuiella, IV, 51
 Buck-bug, V, 145
 Moth, V, 127
 Egg of, V, 128
 Issuing of, V, 132
 Larva of, V, 129
 Natural enemies of, V, 132
 Pupa of, V, 131
 Sting of larva of, V, 131
 Buffalo-gnat, V, 13
 Tree-hopper, V, 121
 Bugs, how to pin, V, 34
Buprestidæ, VII, 72
Buprestis gigantea, IV, 141
 Burns, A. M., on Locusts, VII, 138
 Burrows, J. H., on Rocky Mountain Locust, IX, 71
 Burt, Huron, on Codling Moth, III, 103
 Gooseberry Spanworm, IX, 5
 Plum Gouger, III, 40
 Tortoise-beetle, II, 57
 Bush & Son, list of grape-vine cuttings, showing relative ease of propagating, IV, 65
 Butcher-bird destroying Canker Worm, VI, 27
 Butterflies, classification of, V, 12
 preparing of, for cabinet, V, 35, 36
 pupation of, III, 146, IV, 55, VI, 138, VIII, 179, Supp., 55
 Two of our common, III, 142
 Butterflies, swarming of, III, 151
 Byfield, Jno., on Rocky Mountain Locust, VII, 152
- C.**
- Cabbage-bug, the Harlequin, IV, 35
 Cabbage Butterflies, II, 111, V, 26
 Butterfly, the Potherb, II, 105
 Rape, II, 107
 Southern, II, 104
 caterpillar, the Zebra, II, 112
 -fly, IV, 35
 -maggot, V, 13
 Plant-louse, II, 10
 Plusia, II, 110, 112
 Tinea, II, 10, IV, 36
 Worms, II, 104, 123
 Remedies for, II, 109
 Cabinet and boxes for insects, V, 37
 Cabinet of insects prepared for Missouri State University, VII (preface, p. 5).
 Caddice-flies, V, 16
cagurus, *Pollyxerus*, VII, 106
calidum, *Calosoma*, I, 89, 115, II, 103, VIII, 52
 California, Phylloxera ravages in, VIII, 163
 Silk-growing in, IV, 79
californica, *Ecanthus*, Supp., 61
caliginosus, *Harpalus*, I, 115, VIII, 52
Callidium amoenum, IV, 54
Callimorpha clymene, III, 134
 fulvicosta, III, 132, 134, VI, 92, Supp., 56
 lecontei, III, 134, VI, 92, Supp., 55
 vestalis, III, 133
Callidryas, III, 151
Callochloa viridis, III, 150
Callosamia angulifera, IV, 122, 128
 promethea, IV, 121, Supp., 55
calmariensis, *Galeruca*, II, 10, 95, VII, 5, 86
Calocampa exoleta, VIII, 23
calopteni, *Anthomyia*, IX, 92, 95, Supp., 89
Caloptenus atlantis, VII, 169, VIII, 113, 114, 116, 117, 118, 153, IX, 86, Supp., 89
 bivittatus, VII, 124, 173
 differentialis, VII, 124, 171, 173, 180, VIII, 150, 153
 femur-rubrum, VII, 126, 128, 170, VIII, 114, 115, 116, 117, 118, 153, IX, 86, Supp., 89, 90
 italicus, VII, 133, VIII, 140
 occidentalis, VIII, 116
 spretus, VII, 121, 128, 138, 170, 180, VIII, 57, 109, 114, 115, 116, 117, 118, IX, 57, Supp., 89, 90
 viridis, VIII, 117
Calosoma, III, 129
 calidum, I, 89, 115, II, 46, 103, VIII, 52
 externum, VIII, 52
 obsoletum, IX, 98
 scrutator, II, 103, III, 129, VIII, 52
 wilcoxi, VIII, 52
 Cambre, Eugene, on preventing Phylloxera, VII, 113
 Camel-cricket, I, 169, III, 68
 Campbell, G. W., on Grape Phylloxera, VII, 100
 Campbell, W. A., on Rocky Mountain Locust, IX, 73

- Campoplex fugitivus*, I, 139
Campyloneura vitripennis, III, 137
Canada Warbler destroying Canker Worm, VI, 27
Canadian Entomologist, article from, on Hellebore for Currant Worm, IX, 14
Canaday, Elihu, on Army Worm, VIII, 40
canadensis, *Myiodioctes*, VI, 27
candida, *Saperda*, Supp., 53
Cane Curculio of the Grape, I, 131
caniculi, *Cuterebra*, I, 164
Canker worms, I, 9, 109, II, 11, 15, 54, 94, 95, 99, 101, 103, III, 128, 160, IV, 23, 40, VI, 24, VII, 80, VIII, 12, Supp., 56
 Bandages for, VI, 26, 27
 Birds that devour the worms, VI, 27, 28
 Destroyed by plowing, II, 100
 Distinction between two species of, VI, 27
 Enemies of, II, 102
 Extract from the original Essay on, by W. D. Peck, VII, 89
 Fall canker-worm, VII, 83, VIII, 18
 Oviposition of the two different species, VIII, 37
 Origin of, II, 96
 Paris green for, VI, 26
 Practical considerations, VII, 85
 Remedies against, II, 98, VI, 26, VIII, 17, 20
 Spring canker-worm, VII, 80, VIII, 18
 Traps, VI, 26, VIII, 20, 21
 Trough remedy for, VI, 26
 Two species defined, VIII, 13
Cantharides, V, 18
capax, *Xylina*, III, 136, Supp., 75
Capers, Dr., on Cotton Worm, II, 38
Capsus oblineatus, II, 113, VII, 27
 vitripennis, III, 139
Carabidae, VI, 123
Carabid larvæ, I, 59, IX, 97, Supp., 52
Carbolate of lime for Potato bugs, IV, 14
Carbolic acid for Grape-vine Root-lice, IV, 68
cardui, *Cynthia*, III, 151, IV, 129
 Pterophorus, Supp., 83
carduidactylus, *Pterophorus*, I, 180, III, 67
Caris, VI, 52
carnaria, *Sarcophaga*, VII, 180, IX, 95, Supp., 60
Carolina Locust, VII, 179, IX, 92
carolina, *Mantis*, I, 169, III, 68
 Edipoda, VII, 175, 179
 Sphinx, I, 96, IV, 129
carolinensis, *Mimus*, VI, 27, VIII, 124
Carpenter, D., on Rocky Mountain Locust, VIII, 102, IX, 70
Carpet Moth, II, 10
Carpocapsa, V, 50
 pomonella, I, 62, 108, II, 10, III, 6, 101, IV, 27, V, 154
 vitisella, I, 133
Carr, Wm., on Army Worm, VIII, 39
Carrion-beetle, American, VI, 100
 -feeders, V, 11
Carson, J., on Rocky Mountain Locust, IX, 75
caryæ, *Scolytus*, V, 103, 107, Supp., 54
caryæcaulis, *Phylloxera*, VII, 91, 97, 99, 117
caryæ-fallax, *Phylloxera*, VII, 118
caryæfolia, *Phylloxera*, IV, 66, VI, 45, VII, 117
caryæ-globuli, *Phylloxera*, VII, 117
caryæ-gummosa, *Phylloxera*, VII, 118
 -*ren*, *Phylloxera*, VII, 118
 -*semen*, *Phylloxera*, VII, 117
 -*septa*, *Phylloxera*, VII, 118
caryævenæ, *Phylloxera*, VII, 117
Case-bearer, Walnut, IV, 42
 -bearing Coleoptera, VI, 127, 128
casei, *Piophila*, II, 10
Caskie, Robt. E., on Army Worm, VIII, 39
Cass County (Mo.) *Courier*, article from, on Rocky Mountain Locust, VIII, 65
Cassida, I, 100, II, 58, 59
 aurichalcea, II, 62, Supp., 53
 bivittata, II, 61, Supp., 53
 cruciata, II, 63
 guttata, II, 60, 63
 nigripes, II, 63, Supp., 53
 pallida, II, 62, Supp., 53
 signifer, II, 63
 texana, Supp., 54
 trabeata, II, 63
cassinii, *Cicada*, I, 20, 21, IV, 33, Supp., 59
castaneæ, *Phylloxera*, VII, 118
Castnia licus, VIII, 178
 linus, VIII, 178
 yuccæ, VIII, 173
Castnioides, a proposed tribe of butterflies, VIII, 179
Castor-bean Silkworm, IV, 112
Cat Bird destroying Rocky Mountain Locust, VIII, 124
 destroying Canker Worm, VI, 27
Caterpillars of the Vine—The blue, VI, 87
Catocala, VIII, 178
 phalanga, III, 166
Cecidomyia, Supp., 59
 destructor, II, 10, 19, V, 25, VII, 27
Cecidomyiidae, larval characters of, V, 114
cecropia, *Attacus*, III, 129, 170, IV, 74, 103, 138
 Hyalophora, IV, 103
 Platysamia, IV, 103
 Samia, IV, 103, Supp., 55
Cecropia Cryptus, IV, 110
 Silkworm, IV, 103
 Food-plants, 104
 Larval Changes, IV, 106
 Parasites of, IV, 107
 Tachina-fly, IV, 108
 worm, III, 7
cecropiæ, *Exorista*, IV, 108, Supp., 60
Cedar-bird destroying Canker Worm, VI, 27
cedrorum, *Ampelis*, VI, 27
Celæna, I, 68
 egens, Supp., 56
 herbimacula, I, 86
 infecta, Supp., 56
 murcimaculata, Supp., 56
 oblonga, III, 136, Supp., 75
 renigera, I, 86, Supp., 56
 subcadens, Supp., 56
celtis, *Apatura*, VI, 139, 142
Ceutorhynchus napi, III, 11
ceparum, *Anthomyia*, I, 155, II, 9
cerasi, *Selandria*, II, 19, VII, 27
Cerasphorus cinctus, III, 7, VII, 76
Ceratocampinae, V, 127

- Ceratopogon*, Supp., 59
cereana, *Galleria*, I, 166, II, 10, Supp., 57
Ceresa bubalus, V, 12, 121
Ceretes, VIII, 178
Certhiadæ, IV, 28
cervicalis, *Scymnus*, I, 122
Chærocampa, II, 71
 pampinatrix, II, 71
Chalcididæ, V, 88
Chalcis, II, 92, III, 158
 albifrons, II, 52, VIII, 54
 marice, IV, 109, 110, 123
Chalcis-fly, II, 52, IV, 51, V, 89
 The inflating, I, 176
chalybea, *Haltica*, I, 101, III, 79, 81, Supp., 53
 Chambers, V. T., on Cicada, I, 20, 28
 Change of habit, III, 91
 Chapin, Oliver, on Codling Moth, IV, 26
charus, *Bracon*, VII, 75, Supp., 75
 Chatterer of Carolina destroying Canker Worm,
 VII, 90
Chauliognathus, IV, 30
 americanus, Supp., 53
 marginatus, V, 154
 pennsylvanicus, I, 57, IV, 28, V,
 154, Supp., 53
 Cheese Fly, II, 19, VII, 27
 Maggot, II, 10
Chelymorpha, II, 58, 59
 cribraria, II, 58
chenopodii, *Hadena*, Supp., 76, 77
 Cherished Bracon, VII, 75
Chermes pinicorticis, V, 100
 Cherry-bird destroying Canker Worm, VII, 90.
 Chestnut-sided Warbler destroying Canker
 Worm, VI, 27.
 Chickadee storing corn in cocoons, IV, 107
 destroying Canker Worm, VI, 27
 Chickweed Geometer, I, 179. [See Knotweed]
Chicago Times, article from, on Rocky Mountain
 Locust, VIII, 73
Chicago Tribune, articles from, on Rocky Mount-
 ain Locust, VII, 155, VIII, 82, 107
 Child, A. J., on Rocky Mountain Locust, VIII, 91
Chilocorus bivulnerus, I, 16, V, 100
 Chinch Bug, II, 2, 6, 11, 15, 16, 19, 20, 32, 35, 70, 114, V,
 12, 19, 62, VII, 19, 190, VIII, 22, 120,
 142, 143
 Abstaining from the cultivation of the grains
 upon which the insect feeds, VII, 38
 Amount of damage done by, II, 28
 Appearance and transformations of, VII, 20
 Appendix to the article on, VII, 51
 Bogus Chinch Bug, II, 31, V, 112
 Burning as remedy for, VII, 32
 Cannibal foes of, II, 25
 Destructive powers of, II, 22, VII, 24
 Direct remedies against, VII, 31
 False Chinch Bug, V, 111
 Flight of, VII, 29
 Food plants, VII, 26
 Heavy rains destructive to, II, 24, VII, 30
 Importance of winter work and combined
 action, VII, 36
 Injurious to stock, VII, 43
 Injuries in Missouri in 1874, VII, 25
 Chinch Bug—Continued.
 Injuries in 1874, VII, 24
 Invigorating the plant by manure, early sow-
 ing, etc., VII, 34
 List of correspondents who made returns, VII,
 51
 Migration on foot, VII, 30
 Mixing seed or protecting one plant by another,
 VII, 34
 Mode of reproduction and hibernation, VII, 27
 Natural enemies, VII, 38
 Natural history, II, 18
 Past history, II, 17, VII, 22
 in Missouri, VII, 22
 Possible remedial and preventive measures
 that need further and thorough trial, VII,
 41
 Preventing the migration of, from one field to
 another, VII, 35
 Preventive measures, VII, 32
 Prognosticating, VII, 24
 Questions answered by correspondents, VII,
 52
 Recapitulation of its natural history, II, 36,
 VII, 49
 Remedies against, II, 28
 Rolling as preventive, VII, 33
 Unnecessary fears, VII, 44
 Where the eggs are laid, VII, 28
 Chipping Sparrow destroying Canker Worm, VI
 27
Chlamys, III, 159
 plicata, VI, 128, 130
chlorizans, *Baris*, III, 11
Chlorops, I, 160
 Chronological history of Periodical Cicada, I, 30
Chrysobothris femorata, I, 46, III, 6, VII, 71, Supp.,
 67
 var. *alabamæ*, VII, 71
 fastidiosa, VII, 71
 lesueuri, VII, 71
 misella, VII, 71
 obscura, VII, 71
 4-impressa, VII, 71
 semisculpta, VII, 71
 soror, VII, 71
Chrysomela, II, 57, 59, III, 45, VII, 18
 decem-lineata, VII, 16, 18
 meticulosa, VI, 122
 rhois, VI, 122
 stolida, VI, 122
Chrysomelidæ, III, 14
Chrysopa, I, 57, III, 150, IV, 45
 eriosomatis, I, 123
 illinoensis, II, 26, VII, 39, 40
 plorabunda, II, 26, VI, 51, VII, 40
 tabida, VII, 106
Chrysops vittatus, II, 129
Cicada, VI, 37, VIII, 38
 cassinii, I, 20, 21, IV, 33, Supp., 59
 pruinosa, I, 27
 septemdecim, I, 18, 19, 20, II, 19, III, 6, IV,
 31, VII, 27, Supp., 58, 59
 tredecim, I, 19, II, 19, III, 6, VII, 27,
 Supp., 58, 59
 Cicada, The Periodical, I, 18, IV, 30

- Cicadae*, II, 131
Cicindela circumpecta, IX, 98
 formosa, IX, 98
 fulgida, IX, 98
 punctulata, IX, 98
 pulchra, IX, 98
 repanda, VIII, 52
 scutellaris, IX, 98
 sexguttata, IX, 98
 vulgaris, IX, 98
Cincinnati Gazette, article from, on Rocky Mountain Locust, IX, 84
cinctus, *Cerasphorus*, III, 7, VII, 76
 Harpactor, I, 114, VII, 41, Supp., 58
 Tabanus, II, 129
cinderella, *Tortrix*, IV, 47, Supp., 82
cinerea, *Epicauta*, Supp., 54
 Lytta, I, 97, Supp., 54
 Piesma, II, 32, VII, 47
 Xylina, III, 134, 135, Supp., 75
cinereopunctella, *Elachista*, VI, 138
cinerosa, *Xylina*, III, 136, Supp., 75
cingulatus, *Oncideres*, III, 6
circumcinctus, *Perillus*, IV, 19
Citheronia regalis, III, 151, IV, 129, V, 141
clandestina, *Noctua*, I, 79, Supp., 55
Clandestine Owlet-moth, I, 79
Clark, Rufus, Machine for destroying Locusts, VIII, 129
Clark, William H., on Rocky Mountain Locust, IX, 74
Classification of insects, V, 8
Classification, remarks on, I, 98, 99, II, 71, III, 94, 95, 96, 133, 143, IV, 46, V, 9, VII, 143, 170, VIII, 170, 178, 179
clavata, *Deloyala*, II, 57, Supp., 54
Claxton & Stevens, Curculio Catcher of, III, 22
Clelland, J. L., on Rocky Mountain Locust, IX, 73
Clemens, Dr. B., description by, of *Callimorpha fulvicosta*, III, 133
Cleonymus clisiocampæ, III, 120
Clerus apiarus, VI, 101
Cleveland *Herald*, article from, on Grape-vines, V, 59
Climbing cut-worms, I, 76
Clisiocampa americana, II, 7, III, 117, V, 56
 disstria, Supp., 55
 sylvatica, II, 7, 37, III, 121, IV, 41, Supp., 55
clisiocampæ, *Cleonymus*, III, 120
 Semiotellus, III, 120
Clostera americana, II, 19, VII, 27
Clothes Moth, II, 10
Clover-hay Worm, VI, 102
 Natural History, VI, 105
 Remedies, VI, 105
Clover Worms, VI, 103
 in the State of New York, VI, 104
Clubbed Tortoise-beetle, II, 56, 57
Clumsy Locust, VIII, 148
clymene, *Callimorpha*, III, 134
Clythra, VI, 128, 130
clyton, *Apatura*, VI, 142, 145
Clytus pictus, III, 7, VI, 101
Cnethocampa processionea, V, 126
cnotus, *Otus*, II, 71
Cobalt, for Potato-beetle, IV, 14
Coccidæ, II, 15, V, 16, 92, VI, 33
Coccinella, VI, 51
 munda, II, 25, VII, 39, Supp., 52
 novemnotata, I, 112
 picta, V, 101, Supp., 52
Coccinellidæ, V, 11, 27
Coccotorus, Supp., 54
Coccus adonidum, III, 96, V, 80
 cacti, V, 18
 lacca, V, 18
Coccyzus americanus, III, 121
 erythrophthalmus, III, 121, VI, 27
Cochineal, IV, 84, V, 18
Cochylis hilarana, I, 175, II, 135
cochranii, *Agrotis*, I, 74, Supp., 76, 77
Cochran, J. W., on Dark-sided Cut-worm, I, 75
 on Climbing cut-worms, I, 69
Cochran Rustic, I, 74
Cocklebur *Sphenophorus*, III, 60
Cockroach, II, 10, V, 14
Cocoon, issuing of moth from, IV, 105, 127
Codling Moth, I, 62, II, 6, 10, III, 32, 101, 118, IV, 22, 27, 48, V, 26, 47, VI, 9
 Again, IV, 22
 Attacks peaches, IV, 22, V, 49
 Best kind of bandage for, IV, 23
 False doctrines about, V, 51
 Fires, lights, bottles of liquid as remedies, IV, 27
 Found in California, V, 49
 Jarring as remedy, IV, 25, V, 48
 Natural enemies of, IV, 28, V, 49
 New methods of trapping, IV, 23
 Remedies for, I, 65, IV, 25, V, 48
 Sweetened water as remedy for, IV, 138
 Time of year that the first moths appear, IV, 22
 Wier's Trap, IV, 23, V, 47, VI, 10
Cœliodes inæqualis, I, 128, III, 60, Supp., 54
Colaspis barbara, III, 82
 brunnea, III, 82
 flavida, III, 44, 63, 81, 84, IV, 34, V, 108
 suilla, III, 82
Colby, Lewis, on Locust ravages, VII, 168
Cole, M., on Phylloxera, VIII, 166
Coleoptera, classification of, V, 10
 preparing of, for cabinet, V, 34
Collecting insects, V, 29
Colman's *Rural World*, article from, on Phylloxera, VI, 83, 84
Colorado Farmer, article from, on Rocky Mountain Locust, VIII, 84, 156, IX, 62
Colorado Locust, VII, 188
Colorado Potato-beetle, I, 101, 102, II, 6, 19, 25, 32, 59, III, 80, 98, IV, 5, V, 26, 52, 62, VI, 11, 17, 18, VII, 1, 29, 39, VIII, 1, 137, IX, 17, 34
 Alarm about it abroad, VII, 3
 Amount of damage caused by it in Missouri, IV, 7
 An addition to its natural enemies, IX, 40
 Area invaded by, IX, 38
 Arsenious acid as remedy, IV, 14
 Artificial remedies for, I, 116
 Best means of fighting it, III, 97
 Bichromate of potash as remedy, IV, 14

Colorado Potato-beetle—Continued.

- Bogus experiments, III, 100
 Carbolate of lime as remedy, IV, 14
 Causes which limit the spread of, IX, 38
 Dog-fennel as remedy, IV, 15
 Damage during the year 1875, VIII, 1
 Enemies of, I, 111, 112, 113, 114, 115, IV, 16, V, 52, VIII, 3, IX, 40
 Further experience with Paris green, VIII, 5
 Gray's Improved Sprinkler, VII, 15
 Hellebore as remedy, IV, 14
 How it affected the price of potatoes, IX, 39
 How it traveled, IX, 37
 Is it poisonous? VII, 6
 It passes the winter in the beetle state, VII, 14
 It reaches the Atlantic, VII, 1
 It spreads, but does not travel in the sense of leaving one district for another, IV, 9
 Its habits, I, 107
 Its hibernation, IV, 11
 Its injuries in 1871, IV, 5
 Its introduction to Europe, IX, 43
 Its past history and future progress, I, 101
 Its progress eastward, V, 52, VI, 12
 Its scientific name, VIII, 2
 Machine for sweeping it off vines, VIII, 4
 Mandrake or May-apple as remedy, IV, 15
 Mechanical means of destroying, IV, 15, VI, 14
 Modification it has undergone in habits, IX, 40
 Native home of, VIII, 8
 Natural checks increasing, III, 100
 Natural remedies, I, 109
 New food plants of, IV, 10, V, 52, VI, 11, VII, 14
 New means of destruction, VII, 15
 New territory invaded, IV, 8
 Occurrence in the Atlantic States, VIII, 1
 Parasite of, I, 111
 Paris green as remedy, III, 99, IV, 11, V, 52, VI, 13, VII, 8, VIII, 3
 Peck's Spray Machine, VIII, 4
 Placard published by the German Government, IX, 44
 Poisonous qualities of, VIII, 11
 Potato Pest Poison as remedy, IX, 45
 Powdered Hellebore as remedy, IV, 14
 Preparing for it in Europe, VI, 15
 Rate at which it traveled, IX, 37
 Remedies for, I, 109, 116, III, 99, IV, 11, 13, 14, 15, V, 52, VI, 13, VII, 3, 7, 8, VIII, 3, IX, 45
 Spread of the insect during the year 1876, IX, 34
 Sulphate of copper as remedy, IV, 14
 The beetle eats as well as the larva, VII, 14
 The Bogus Colorado potato-beetle, I, 105
 The proper scientific name of the beetle, VII, 16
 The true remedy, III, 101
 Use of straw as a preventive, VIII, 4
 Colorado Potato Bug (see Colorado Potato-beetle).
 Colorado, Rocky Mountain Locust in, VIII, 84, IX, 62
columbia, *Samia*, IV, 107, 111, 128, Supp., 55
comma, *Leucania*, VIII, 43
commelinæ, *Prodenia*, I, 88, III, 13, Supp., 56
 Commission, National Entomological VIII, 133
 Common Currant Plant-louse, IX, 2
 Flesh-fly, VII, 180, IX, 95
 May Beetle, III, 8

- Common Pruner, III, 6
 Quail destroying Chinch Bug, II, 28
 Yellow Bear, III, 67
communis, *Melanotus*, III, 6
compta, *Oeta*, I, 151, Supp., 58
 Pæcilopectera, I, 152
comptana, *Anchylopera*, I, 143
 Phoxopteris, Supp., 57
concavum, *Platyphyllum*, V, 124, VI, 167
conchiformis, *Aspidiotus*, I, 7, II, 9, 10, III, 93, Supp., 86
Conchylis ambiguella, Supp., 57
 Cone-like willow-gall, VI, 155
conformis, *Xylina*, III, 136, Supp., 75
conica, *Phylloxera*, VII, 118
conicus, *Rhynchites*, III, 11
Conocephalus, VI, 155
conotracheli, *Porizon*, III, 28, Supp., 64
Conotrachelus, V, 154
 cratægi, III, 35, 39
 juglandis, Supp., 54
 nenuphar, I, 50, III, 11, 28, 127, Supp., 54, 65, 68
conquisitor, *Pimpla*, IV, 43
consociella, *Acrobasis*, IV, 45
conspersa, *Amphipyra*, III, 75, Supp., 75
 Consumptive Lace-wing, VII, 106
Contopus virens, VI, 27
contracta, *Meracantha*, VI, 118
contractilis, *Hoplophora*, VI, 54
convergens, *Hippodamia*, I, 112
 Convergent Lady-bird, I, 112
convolutella, *Myelois*, Supp., 57
 Zophodia, Supp., 57
 Copper, sulphate of, for Potato-beetle, IV, 14
Coptocycla, II, 58, 59, 63, Supp., 53, 54
 aurichalcea, Supp., 53
 guttata, Supp., 53
 Coral-winged Locust, VIII, 104
Coreus linearis, II, 113
 tristis, I, 113, II, 31, VII, 46, Supp., 58
Corimelæna lateralis, II, 35
 pulicaria, II, 33, VII, 48
 unicolor, II, 35
 Cornaby, Samuel, on Silkworms, IV, 101
 Corn Anthomyia, I, 155
 Cut-worm, I, 87
 Maggot, I, 154
 Rustic, I, 81
 Sphenophorus, III, 59
 Worm, III, 45, 104, 105, 111
cornutus, *Corydalus*, V, 143, IX, 125, Supp., 63
 Passalus, IV, 139, 140
Corpodacus purpureus, VI, 27
Corydalus cornutus, V, 143, IX, 125, Supp., 63
Corynetes rufipes, VI, 96, 101
 violaceus, VI, 101
Coscinoptera dominicana, VI, 127
Cossus, VIII, 177
costalis, *Asopia*, VI, 102
 Tabanus, II, 128
Cotalpa lanigera, V, 10
 Cotton Army-worm, II, 41, VIII, 23
 Boll-worm, III, 111
 moth, II, 40
 Position of when alighting, VI, 24

- Cotton Worm, II, 38, III, 105, V, 19, 68, VI, 17, VII, 9
 Hibernation of, VI, 22
 Natural enemies of, VI, 23
 Paris green as remedy, VI, 20
 Range of, VI, 23
 worms, II, 37
- Cottonwood Dagger, II, 119
 Gall plant-louse, I, 120
- Country Gentleman*, article from, on Apple Worm
 remedy, V, 48
 article from, on Clover Worm,
 VI, 104
 article from, on Curculio, IV,
 26
 article from, on Grape-vine
 Colaspis, III, 82
 article from, on Locusts, VII,
 172, VIII, 152
 article from, on Phylloxera,
 V, 59, VI, 82
 article from, on Potato-beetle,
 I, 111
 article from, on Tent-caterpil-
 lar, III, 125
- County reports (Mo.) on Rocky Mountain Locust,
 IX, 68
- Crabro stirpicola*, IX, 95, Supp., 89
- crabro*, *Vespa*, IV, 22
- Cranberry-weevil, III, 60
- Crandall, O. A., on Rocky Mountain Locust, IX, 74
- Crane-flies, I, 180, II, 132
- crantor*, *Pholus*, II, 74
Sphinx, II, 74
- Craponius*, Supp., 54
- cratægi*, *Conotrachelus*, III, 35, 39
- Cratoparis lunatus*, III, 10
- Cream Callimorpha, III, 133
- Creepers destroying Codling Moth, IV, 28
 Black and White, destroying Canker
 Worms, VI, 27
- Creighton, Samuel, Insect-destroyer invented by,
 IV, 15
- cressonii*, *Hemiteles*, I, 177, Supp., 65
- Craig, Wm. G. M., on Rocky Mountain Locust, IX,
 73
- cribraria*, *Chelymorpha*, II, 58
- Crickets (*Gryllidæ*), V, 14, VI, 154
- crinitus*, *Myiarchus*, VIII, 124
- Crioceris asparagi*, II, 10, 13, 19, VII, 5
merdigera, II, 58
- Croton Bug, II, 10
- Crow destroying Katydid, VI, 162
 Potato-beetles, VIII, 3
- Crow, J. H., on Rocky Mountain Locust, IX, 72
- Crow Blackbird destroying Canker Worms, VI, 27
 Codling Moths, IV, 28
 Locusts, VIII, 124
 Pea-weevils, III, 50
- Crown-borer of the Strawberry, III, 44, 83
- cruciata*, *Cassida*, II, 63
- cruciferarum*, *Plutella*, II, 10, IV, 36
- Crustacea, a class of Segmented animals, V, 6, 7
- Cryptoblabes bistriga*, IV, 46, Supp., 87
- Cryptocephalus*, VI, 128, 130
- Cryptus extrematis*, IV, 110, 111, 123, Supp., 52
grallator, VII, 75
inquisitor, I, 150
- Cryptus nuncius*, IV, 110, 111, 123, Supp., 52
samiæ, IV, 110, 111, Supp., 52
smithii, IV, 111
- Otenucha americana*, II, 85
- Cuckoo, Yellow-billed, destroying Canker Worm,
 VI, 27
 Locusts, VIII,
 124
- Cucumber-beetle, I, 100, II, 62, 65
 flea-beetle, I, 101, II, 57, V, 112
- cucumeris*, *Haltica*, I, 101, II, 57, Supp., 53
- cunea*, *Hyphantria*, Supp., 55
- Curculio, II, 11, 16, 92, III, 13, 16, 25, 29, V, 22
 Apple, III, 29
 Natural history of, III, 30
 Remedies for, III, 34
- Catcher, Hooten's, III, 23
 Hull's, III, 19, V, 25
 Ward's, III, 20
- Grape, I, 128, III, 60
- Grape-cane Gall, I, 131
- Grape-seed, I, 129
- Plum, I, 50, III, 11
 Enemies of, I, 57
 Parasites of, III, 24
 Points in its natural history, I, 50, III,
 11
 Remedies for, I, 60, III, 15
 Quince, III, 35
- Curculionidæ*, characteristics of, III, 9
 creaking noise produced by, III, 14
- curculionis*, *Sigalphus*, III, 25, 27, Supp., 67
- cucurbitæ*, *Ægeria*, II, 64
- Currant Aphis, VI, 46
 Fruit-worm, I, 140
 Plant-louse, II, 10
 -stem Borer, VI, 108
- Currant Worm, II, 8, 9, 96, IV, 14, IX, 1
 the imported, IX, 7
 the native, IX, 23
 fly, IX, 19
- Cursoria, a section of Orthoptera, V, 14
- cursoria*, *Agrotis*, I, 78
- curvicauda*, *Phaneroptera*, VI, 164
- Curvirostra leucoptera*, VI, 27
- Cuterebra*, III, 150
caniculi, I, 164
- Cut-worm lion, I, 90
 The climbing, I, 69, 76
 The Dark-sided, I, 74
 The Dingy, I, 82
 The Glassy, I, 83
 The Greasy, I, 80
 The small White Bristly, I, 86
 The Speckled, I, 84
 The Variegated, I, 72
 The Western Striped, I, 81
 The Wheat, I, 87
 The W-marked, I, 79
- Cut-worms, I, 67, II, 16, 45, III, 6
 Natural history of twelve distinct species, I, 67
 Remedies against, I, 89, 90
- cyanea*, *Cyanopiza*, VI, 27
- Cyanopiza cyanea*, VI, 27
- Cycloneda sanguinea*, Supp., 52
- Cylindrical Orthosoma, I, 126, II, 91
- cylindricum*, *Orthosoma*, I, 124, II, 87

Cyllene pictus, VI, 101
Cyllocoris scutellatus, V, 154
Cynipidæ, VI, 70
Cynips, II, 135
 gallæ-tinctoriæ, V, 18
 quercus-aciculata, Supp., 59
 quercus-inanis, I, 14
 quercus-spongifica, I, 14
cynosbana, (*Spilonota*), Supp., 57
cynthia, *Attacus*, III, 170, IV, 74, 112, 138
 Samia, IV, 112
Cynthia cardui, III, 151, IV, 129
Cyrtophyllus, V, 123

D.

Dactylopius longispinus, IV, 70, VI, 63
Dactylosphæra, III, 93, 94
 caryæ-magnum, VII, 117
 caryæ-semen, VII, 117
 caryæ-septum, VII, 117
 conicum, VII, 118
 coniferum, VII, 118
 depressum, VII, 118
 forcatum, VII, 118
 globosum, VII, 117
 hemisphericum, VII, 117
 spinsum, VII, 118
 subellipticum, VII, 117
 vitifoliæ, I, 13
Dactylosphæridæ, III, 85, VI, 31
Dade County Advocate, article from, on Rocky Mountain Locust, VIII, 68
Daggy, E., on False Army-worm, III, 112
 on Apple Curculio, III, 33
Dakota, Rocky Mountain Locust in, VIII, 85, IX, 59
Dakrura turbatella, Supp., 57
Danaïs, II, 125, III, 161, 168, 169, V, 146
 archippus, III, 143, 167, IV, 129, V, 149, Supp., 55
 berenice, III, 143
 erippus, III, 143
Daphne pandorus, II, 76
Darapsa myron, II, 71
Dark-sided Cut-worm, I, 74
 tipped Anomalon, IX, 55
Dart-bearing Rustic, I, 82
Darwin, Charles, on Evolution, III, 172, 173
Darwinism, argument in favor of, III, 173
Datana ministra, III, 124, 127, 129, IV, 129
Davis, C. K., on Rocky Mountain Locust, VII, 154
Dawson, G. M., on Rocky Mountain Locust, VII, 155, IX, 78
Dean, J. J., on Colorado Potato-beetle, VIII, 2
decemlineata, *Chrysomela*, VII, 18
 Doryphora, I, 101, 103, IV, 8, VI, 12, 18, VII, 1, 16, 18
Deer Fly, V, 127
Defakaugh, David, report on Rocky Mountain Locust, IX, 70
Definition of Entomology, V, 5
Defunctionation of special parts in the imported Currant Worm, IX, 19
Deilephila lineata, III, 140
Deiopeia bella, V, 11
Delicate Long-sting, V, 50

Deloyala, II, 59
 clavata, II, 57, Supp., 54
Dendroica æstiva, VI, 27
 discolor, VI, 27
 maculosa, VI, 27
 pennsylvanica, VI, 27
 striata, VI, 27
Department of Agriculture, inefficiency of, VII (preface, p. 5).
depressa, *Phylloxera*, VII, 118
Dermaleichus, V, 87
Dermestes, IV, 97, V, 41
 lardarius, VI, 100
Desmia maculalis, III, 61
Descriptive entomology, comments on, III, 123
Destination of departing Locusts, VIII, 107
Destitution in Missouri from Locust injuries, VIII, 91
Destructive powers of the Chinch Bug, II, 22
destructor, *Cecidomyia*, II, 10, 19, VII, 27
 Nysius, VII, 190, Supp., 84, 85
Devastating Dart, I, 83
devastator, *Agrotis*, I, 83, Supp., 56
Devil's Riding-horse, I, 169
Diabrotica vittata, I, 100, II, 62, 64, III, 6
 12-punctata, II, 66
diadema, *Sinea*, Supp., 58
diana, *Argynnis*, III, 69, 171
Diaspides, a subfamily of *Coccidæ*, V, 92
Diaspis, V, 91, Supp., 60
 ostreæformis, Supp., 60
diastrophæ, *Eurytoma*, Supp., 68
Dictyoptera, a division of Neuroptera, V, 14
Differential Locust, VII, 124, 173, VIII, 150, 153
 eggs of, VIII, 154
differentialis, *Caloptenus*, VII, 124, 173
Digger Wasp, II, 106, III, 8, VII, 174
dilatatus, *Velleius*, IV, 22
Dimera, a division of Homoptera, V, 13
dimidiatus, *Micropus*, VII, 22
Diminished Pezomachus, II, 52, VIII, 54
Dimorphism in butterflies, III, 165
 locusts, VIII, 115
 Prionus, II, 90
Dinarda, IV, 22
Dingy Cut-worm, I, 82
Diplosis tritici, II, 10
Diptera, classification of, V, 13
discoideus, *Bruchus*, III, 45
discolor, *Dendroica*, VI, 27
Diseases of Mulberry Silkworm, IV, 87
Disippus Butterfly, II, 125, III, 153, 168, 169
 Description of mature larva, III, 154
 Description of the egg, III, 154
 Parasites of, III, 157
 Winter quarters of, III, 155
 Microgaster, III, 158
disippus, *Limenitis*, III, 153, 168, 169, 171, Supp., 66
 Nymphalis, II, 125
dispar, *Hypogymna*, II, 10
distria, *Clisiocampa*, Supp., 55
distinctus, *Passalus*, IV, 141
Dixon, F. M., on Army Worm, VIII, 39
Dodge, C. R., on Rocky Mountain Locust, VII, 164, VIII, 173
Dolichonyx oryzivorus, VI, 27, VIII, 52

- domestica*, *Musca*, II, 10
 Domestication of insects, IV, 85
dominicana, *Coscinoptera*, VI, 127
 Dominican Case-bearer, VI, 127
 Dopf, J. D., on Canker Worms, II, 98
 on Rocky Mountain Locust, IX, 68
 Dorr, R. L., on Grape-vine Tomato-gall, V, 118
dorsatum, *Phalangium*, IV, 17
Doryphora concatenata, VIII, 2
 10-lineata, I, 101, 103, IV, 8, VI, 12, 18,
 VII, 1, 16, 18, VIII, 1, IX, 34, 43
 juncta, I, 103, 105, VII, 18, IX, 39, 43
 melanothorax, VIII, 10
 undecimlineata, VIII, 10
doryphoræ, *Lydella*, I, 111, IV, 6, Supp., 88
 Dotted-legged Plant-bug, IV, 19
 Douglass, J. B., on Army Worm, VIII, 39
 Downy Woodpecker destroying Canker Worms,
 IV, 28, VI, 28
Doxocopa, VI, 142
 Dragon Flies as enemies of Cicada, I, 26
Drasteria, VIII, 178
 Drop Worm, I, 147
Dryocampa rubicunda, III, 123, V, 137
 Larval changes of, V, 138
 Natural enemies of, V, 139
 Remedies for, V, 140
 senatoria, III, 123, IV, 41
 stigma, III, 123, IV, 41, V, 141
Dujardinii, *Hypopus*, VI, 53
 Dung-beetles, creaking noise made by, III, 14
 -carriers, II, 58, VI, 128
 Dunkley, B. F., on Rocky Mountain Locust, VIII,
 149
 Dunlap, M. L., on Rocky Mountain Locust, VII,
 155
 Dunn, William, on Rocky Mountain Locust, VII,
 152, IX, 118
duodecimpunctata, *Diabrotica*, II, 66
 Durand, J. B., on Rocky Mountain Locust, VIII,
 63
 Duties of State Entomologist, V, 27
 Dwarf Trogosita, III, 6
 Dye, A. A., on Rocky Mountain Locust, IX, 69, 117
 Dr. A. H., on Army Worm, VIII, 39
 Dyer, D. P., on Army Worm, VIII, 39
- E.**
- Ear-fly, II, 129
 Early, Sam. H. Y., on White Grub fungus, I, 158
 Earwigs, characteristics of, V, 16
 East India ants, III, 8
echinopus, *Tyroglyphus*, VII, 106
 Economic entomology, importance of, V, 18, VII
 (preface, p. 4).
Ecpantheria scribonia, IV, 141, 143
Ectobia germanica, II, 10
 Eddleston & Williams, manufacturers of fine ento-
 mological pins, V, 35
 Edwards, C. R., on Broad-necked Prionus, II, 88
egens, *Celaena*, Supp., 56
 Egg-burster of *Corydalis*, IX, 127
 Egg-guide, IX, 87
 Egg of Abbott's Pine Worm, IX, 31
 Apple Curculio, III, 31
 Archippus Butterfly, III, 144
 Egg of Beautiful Wood Nymph, VI, 89
 Broad-necked Prionus, V, 56
 Broad-winged Katydid, V, 123
 Chinch-Bug, II, 21, VII, 21
 Common May Beetle, V, 55
 Cotton Worm, II, 38
 Disippus Butterfly, III, 153
 Dominican Case-bearer, VI, 128
 Flat-headed Apple-tree Borer, VII, 73
 Gooseberry Span-worm, IX, 4
 Horned Passalus, V, 55
 Imported Currant-worm, IX, 10
 Mulberry Silkworm, movements of, IV,
 86
 Native Currant Worm, IX, 25
 Narrow-winged Katydid, V, 124
 Oblong-winged Katydid, V, 123
 Oeta compta, Supp., 58
 Pea-weevil, III, 47
 Strawberry Worm, IX, 28
 Yucca Borer, VIII, 174
 Eggs in Canes and Twigs, V, 119
 of the American Tent-caterpillar, III, 118, V
 56
 Army Worm Moth, VIII, 34, 183, 184
 When laid, VIII, 40, 182, 183
 Where laid, II, 48, VIII, 34, 182
 Belostoma grandis, IX, 128
 Buffalo Tree-hopper, V, 121
 Canker Worms, II, 94, VII, 82, 84, 86
 VIII, 13
 Chinch-Bug, VII, 21
 Differential Locust, VIII, 154
 Frosted Lightning-hopper, V, 122
 Grape Phylloxera, IV, 59, VI, 34, 38, 41,
 92, 98, VIII, 158
 Harlequin Cabbage Bug, IV, 37
 Hackberry Butterflies, VI, 139, 141, 148
 Jumping Sumach Beetle, VI, 120, 121
 Jumping Tree-hopper, V, 119
 Mantis carolina, I, 170
 Ecanthus latipennis, V, 119, Supp., 60
 Orchelimum, V, 123
 Orocharis saltator, Supp., 62
 Periodical Cicada, I, 25
 Rocky Mountain Locust, VII, 122
 Effects of burying at different
 depths, IX, 104
 Effects of exposure to air, IX, 104
 Effects of freezing and thawing
 on, IX, 99
 Effects of moisture on, IX, 101
 Experiments with, IX, 99
 How laid, IX, 86
 Snowy Tree-cricket, V, 120
 Tent-caterpillar of the Forest, II, 122
 Tortoise Beetles, II, 60, Supp., 53
 Unknown Tree-hopper, V, 122
 Wheat-head Army Worm, IX, 55
 Egg-Mass of Hellgrammite, IX, 120
 Rocky Mountain Locust, Philosophy
 of, IX, 87
 Egg-Parasite, the *Anthomyia*, IX, 92
egle, *Euchætes*, I, 139, III, 133, IV, 41
 Eight-spotted Forester, I, 136, II, 83, 86, VI, 94
Elachista cinereopunctella, VI, 138

- Elaphidion parallelum*, III, 6, IV, 54
villosum, III, 6
Elaphrus ruscarius, VIII, 52
 Elliott, F. R., on dying of Grape-vines, V, 59
 Elm Leaf-beetle, II, 10, VII, 5
 Scolytus, V, 107
 Elm-tree Louse, I, 123
 Elongate Ground-beetle, I, 115
elongatus, *Pasimachus*, I, 115, VIII, 52
 Elytra of Coleoptera, characteristics of, V, 10
 Emery, H. D., on Climbing Cut-worm, I, 77
 Rocky Mountain Locust, VII, 138
Emmenadia pectinata, VI, 125
Emphytus maculatus, IX, 27
Empidonax minimus, VI, 27
Empretia stimulea, V, 126
Endropia armataria, IX, 7
 Engelmann, Dr. G., on the Grape-vines of the U.
 S., IV, 60, VI, 70
 on Locusts, IX, 84
 Engelmann, Theod., on Grape-vine grafting, VI, 80
 on Raspberry Root-borer, VI, 112
 Entomological collecting instruments, V, 29
 Commission, argument in favor of
 creating, VII (preface, p. 5), VIII,
 134, 137
 pins, V, 34
 Entomologist, duties of a State, V, 27
 Entomology, definition of, V, 5
 Economic importance of, V, 18
 Importance of, as a study, V, 17
 Its advancement, V, 5
 Its relations to agriculture, V, 5
 Progress of economic, V, 19
 Entomophaga, a subsection of Hymenoptera, V, 10
 Entomophilous plants, V, 152
entomophagus, *Tyroglyphus*, VI, 52
 Entomotaxy, V, 34
Epargyrius tityrus, VIII, 173
ephemeraeformis, *Thyridopteryx*, I, 147
Ephestia zea, IX, 31
Ephialtes, I, 178
 notanda, IX, 98
Epicærus fallax, III, 58
 formidolosus, III, 58
 imbricatus, III, 58
 vadosus, III, 58
Epicauta cinerea, Supp., 54
 pensylvanica, Supp., 54
Epilachna borealis, IV, 18
 Epimenis, the Grape-vine, VI, 87
epimenis, *Psychomorpha*, III, 63, 64, VI, 87, 88, 90, 95
Epitrix, Supp., 53
Eragrostis poaeoides, VIII, 122
Erax, II, 122, 123
 bastardii, II, 124, IX, 98
 licitor, II, 124
 tabescens, II, 124
Eriosoma, VII, 97
 lanigera, I, 121, III, 95, IV, 69, VI, 63
 pyri, I, 118, III, 5, 95, 96, VI, 37, IX, 43,
 Supp., 59, 87
 rileyi, Supp., 87
 ulmi, I, 123, Supp., 87
eriosomatis, *Chrysopa*, I, 123
erippus, *Danaïd*, III, 143
Eromophila cornuta, IX, 91
 Erwin, J. L., on Fall Army-worm, III, 110
Erycinidae, VI, 138
Erynnis alcea, VIII, 182
 malvarum, VIII, 182
Erysiphe, V, 70
erythrocephalus, *Melanerpes*, VIII, 124
erythrocerus, *Bruchus*, III, 55, 56, Supp., 70
erythrophthalmus, *Coccyzus*, III, 121, VI, 27
Euchaetes egle, I, 139, III, 133, IV, 41
Euclea, VI, 140
 pænulata, V, 126
 querceti, V, 126
Eudemis, Supp., 57
 botrana, Supp., 57
Eudryas brevipennis, VI, 91
 grata, I, 136, II, 83, VI, 88, 89, 95
 unio, I, 136, II, 83, III, 63, VI, 90, 92, 95
Eufitchia ribearia, IX, 3
Eulophus, IV, 51
Eumenes fraterna, II, 103
Eunemoria gracilaria, Supp., 79
Eupelmides, Supp., 52
Eupelmus, VI, 162, Supp., 52
Euphanessa mendica, IX, 6
Euplexoptera, V, 16
Eupsalis minuta, VI, 113, 117
 Europe, American plants and insects acclimated
 in, IX, 43
 European Cattle Breeze-fly, II, 129
 Cock-chaffer, I, 157
 Meal Worm, IX, 43
 Oak Phylloxera, VI, 46, 64, VIII, 158
Euryomia inda, III, 6
 melancholica, III, 6, V, 154
Euryptychia, II, 134
 saligneana, II, 134, Supp., 57
Eurytoma, I, 52
 bolteri, I, 177, Supp., 68
 diastrophii, Supp., 68
Eurytomides, Supp., 68
Euschemon Rafflesia, VIII, 170
Euschistus punctipes, I, 113, IV, 19, 20, V, 12, Supp.,
 58
 variolaris, Supp., 58
Euura, IX, 23
 Evans, J. C., on Rocky Mountain Locust, IX, 70
 Everett, H., on Chinch Bug remedy, II, 29, VII, 35
 Evolution, III, 159, VIII, 170
exitiosa, *Ageria*, I, 47
 Exodus of Locusts, VIII, 104
exoleta, *Calocampa*, VIII, 23
Exorista flavicauda, II, 51, VIII, 53, Supp., 60, 88
 leucanica, II, 50, 51, III, 116, 129, IV, 108
 VIII, 53, Supp., 60
 var. *cecropica*, IV, 108, Supp., 60
 militaris, II, 50, III, 129, IV, 109
 Osten-Sackenii, II, 51
 phycitæ, IV, 40, Supp., 88
externum, *Calosoma*, VIII, 52
extranea, *Leucania*, II, 50
extrematis, *Cryptus*, IV, 110, 111, 123, Supp., 52
 Eyed Emperor, VI, 137
- F.**
- fabæ*, *Bruchus*, III, 52, 55, Supp., 69, 70, 71
fabricii, *Lytta*, I, 99

- Fall Army-worm, III, 109, 150, VI, 17, VIII, 23, 35, 37, 48
 How it differs from the true Army Worm, III, 112, VIII, 48
 Remedies for, III, 114
 Fall Canker Worm, VII, 83, VIII, 18
 Web Worm, II, 11, III, 130
 Natural history of, III, 130
 Remedies for, III, 132
 Fairchild, H. O., on Phylloxera, VI, 83
fallax, *Epicærus*, III, 58
falsarius, *Acoloithus*, II, 86
 False Chinch Bug, V, 111, VII, 46
 Indigo Gall-moth, II, 132
 Farris, M. W., on Rocky Mountain Locust, IX, 69
fasciatus, *Æcanthus*, Supp., 60
fascicularis, *Hemirhipus*, VI, 101
fastidiosa, *Chrysobothris*, VII, 71
 Fasting to avert locust injury, VIII, 96
 Faucon, Louis, on irrigation as remedy for Phylloxera, V, 72
 Faulkner, Dr. S. K., on Rocky Mountain Locust, VII, 140
femorata, *Chrysobothris*, I, 46, III, 6, VII, 71, Supp., 67
 Tiphia, VI, 124
femoratum, *Spectrum*, VI, 156
femoratus, *Micropus*, VII, 22
femur-rubrum, *Caloptenus*, VII, 126, VIII, 114, 115, 116, 117, 118, 153, Supp., 89, 90
 Ferguson, J. T., on Rocky Mountain Locust, IX, 73
 Ferris, Peter, on Tent Caterpillar, III, 125
 Fever Worm, IV, 142, 144
Fidia murina, Supp., 53
 viticida, I, 132, Supp., 53
 vitis, Supp., 33
 Fiery Ground-beetle, I, 89, 115, II, 46, 103
 Fifteen-spotted Ladybird, IV, 17
 Figure 8 minor, I, 86
 Filbert-gall, the Grape-vine, V, 116
 Fillery, Wm. H., on Rocky Mountain Locust, IX, 75
flimbriatus, *Stiretrus*, I, 114, IV, 20
 Fine, F. F., on Rocky Mountain Locust, IX, 71
 Fires for Codling Moth, IV, 27
 Fishburne, Dr. J. H., on poisoning by Colorado Potato-beetle, VII, 7
 Fisher, H. I., on Rocky Mountain Locust, IX, 85
 Fisher, J. C., on Periodical Cicada, I, 20
 Fitch, Col. H., on Rocky Mountain Locust, IX, 71
 Fitch, Dr. Asa, on Army Worm, II, 43, VIII, 25, 50
 on Bee-killer, II, 122
 on Curculio parasite, III, 24
 on Gooseberry Fruit-worm, I, 140
 on Pearl Wood Nymph, II, 84
 on Tent Caterpillar, III, 123, 127
fitchii, *Promachus*, Supp., 60
 Flat-headed Apple-tree Borer, I, 46, 47, VII, 71
 Enemies of, VII, 73
 Natural history of, VII, 72
 Remedies for, I, 47, VII, 76
 Flat-headed Borer, III, 6, VI, 107, 109
 Flea-beetle, IV, 35
 -like Negro-bug, II, 33, 34, VII, 48
 Fleas, characteristics of, V, 15
flavicauda, *Exorista*, II, 51, VIII, 53, Supp., 60, 88
flavicornis, *Anomalon*, III, 69
flavida, *Colaspis*, III, 41, 61, 82, IV, 34
flavifrons, *Scolia*, VI, 124
flavimanus, *Bruchus*, III, 56, Supp., 70
flavimedia, *Prodenia*, Supp., 56
 Flesh-fly, the common, VII, 180
 the Sarracenia, VII, 181
 Flower-beetles, III, 6
 -bug, the Insidious, VII, 41, 47
 Fly-catcher, Great-crested, destroying Locusts, VIII, 124
 -poison as remedy for Potato Bug, IV, 14
 Ford, S. H., apparatus of, for destroying Potato Bugs, I, 116
forcata, *Phylloxera*, VII, 118
 Forest Caterpillar, III, 129
 -flies, V, 13
 Tent-caterpillar, III, 121, 124, 128
 Forester, the Eight-spotted, VI, 94
Forficulidæ, Characteristics of, V, 16
formidulosus, *Epicærus*, III, 58
formosa, *Pepsis*, II, 106
 Foster, E. S., on Broad-necked Prionus, II, 88
 Foster, Suel, on remedy for Codling Moth, I, 65
 on Potato Beetle, I, 110
 Fox-glove leaves as remedy for Gooseberry Span-worm, IX, 7
fragariæ, *Analcis*, III, 42, 44, Supp., 71
 Anchylopera, I, 142, Supp., 57
 Tyloderma, Supp., 72
fraterna, *Eumenes*, II, 103
 Fraternal Potter-wasp, II, 103
 Fringe-wing, VI, 50
frontalis, *Termes*, II, 11
 Frosted Lightning-hopper, V, 122
frugiperda, *Laphygma*, II, 41
 Phalaena, VIII, 48
fugitiva, *Linneria*, V, 41, 141
fugitivus, *Banchus*, IV, 41
 Campoplex, I, 139
Fulgoridæ, V, 122
 Fuller, A. S., on Grape-vine grafting, VII, 109
 on Le Conte's Pine Worm, IX, 33
 on Seed-corn maggot, I, 154
fulvicosta, *Callimorpha*, III, 132, 134, VI, 92, Supp., 55
fulvivenosus, *Micropus*, VII, 22
fulvosa, *Laphygma*, III, 117, VIII, 49
 Fur-moth, II, 10
 Furnas, R. W., on Locust injury in Nebraska, VII, 151, 152
fusca, *Lachnosterna*, Supp., 53
fuscescens, *Turdus*, VI, 27
fuscipennis, *Mesochorus*, VII, 75

G.

 Gad-fly, V, 13
Galeruca californiensis, II, 9, 10, 95, VII, 5, 86
Gallicola or gall-inhabiting type of Phylloxera, V, 63, VI, 34, 66
gallæsolidaginis, *Gelechia*, I, 13, 173, II, 20, 132, 134, III, 158, Supp., 66, 83
 Gall-curculio of the grape, I, 131
Gallerea cereana, I, 166, II, 10
 Gall-gnat, V, 114
gallicola, *Phylloxera*, VII, 93
 Gall-louse, Grape-leaf, IV, 55, 66, 68, V, 63

- Gall-moth, False Indigo, II, 132
 The Misnamed, II, 134
 of the Golden-rod, I, 173
 -moths known to occur in the United States, II, 135
 -nut, V, 18
- Gall, the Grape-vine Apple, V, 114
 Filbert, V, 116
 Tomato, V, 117
 Trumpet, V, 118
Salicis-strobiloides, VI, 155
Vitis-coryloides, V, 116
pomum, V, 114
tomatos, V, 117
viticola, V, 118
- Galls, Aphidian, on Hickory, V, 154
 How produced, VI, 70
 made by moths, II, 132
- Garber, J. B., on irrigating Grape-vines, VI, 76
- Gardener's Chronicle, article from, on Colorado
 Potato-beetle, VII, 5
 article from, on Yucca fer-
 tilization, V, 159
- garrulus*, *Ampelis*, VII, 90
- Gattine*, a disease of Silkworms, IV, 91
- Gazera*, VIII, 178
- Gelechia gallæsolidaginis*, I, 13, 173, II, 20, 132, 134,
 III, 158, Supp., 66, 83
longifasciella, Supp., 83
- gelechice*, *Microgaster*, I, 178, Supp., 66
- geminatus*, *Paniscus*, I, 89
- Genitalia of male Army Worm, VIII, 30
- Geographical range of species, VII, 171, IX, 82
- Geometer of the Chick-weed, I, 179
- Geopinus incrassatus*, I, 77
- Georgetown Miner, article from, on Rocky Mount-
 ain Locust, IX, 62
- Geothlypis trichas*, VI, 27
- germanica*, *Ectobia*, II, 10
- gigantea*, *Buprestis*, IV, 141
- Gigantic Root-borers, III, 75
- Gillman, Henry, on Colorado Potato-beetle, VI, 12
- glaberrimum*, *Orchelimum*, Supp., 62
- Gladish, James E., on Rocky Mountain Locust,
 IX, 73
- glandulella*, *Holcocera*, IV, 144
- Glassy Cut-worm, I, 83
 Mesochorus, II, 52, VIII, 53
 -winged Soldier-bug, III, 137
- glomeratus*, *Microgaster*, III, 167
- Glover, T., on Boll Worm, III, 106, 107, 108
 on Paris green for Cotton Worm, III,
 19
- glycerium*, *Paphia*, II, 125, 127
- Glyphe viridascens*, II, 53, VIII, 53
- Goat-weed Butterfly, II, 125, V, 145
 Additional facts in its history, V, 145
 Its winter quarters, V, 148
 Larval changes, V, 146
 Natural enemies, V, 149
 New food-plant, V, 147
 The egg, V, 146
 The larva, V, 146
 Two broods each year, V, 148
- Göerius olens*, IV, 21
- Gold-banded Tachina-fly, V, 140
- Golden-crowned Thrush destroying Canker Worm,
 VI, 27
 Robin destroying Canker Worm, VI, 28
 -rod Gall-moth, I, 173, III, 158
 Tortoise-beetle, II, 60, 62
 -winged Woodpecker destroying Canker
 Worm, VI, 28
- Goniloba*, II, 125, V, 146
- Goodman, W. S., on Army Worm, VIII, 39
 on Rocky Mountain Locust, IX,
 74, 119
- Gooseberry Fruit-worm, I, 140, II, 9
 Worms, IX, 1
 Span Worm, IX, 3
 A native species, IX, 5
 How it spreads, IX, 5
 It prefers the Gooseberry to the
 Currant, IX, 6
 Its natural history, IX, 3
 Its past history, IX, 5
 Parasites, IX, 6
 Remedies, IX, 6
 The moth is closely imitated,
 IX, 6
- Gordii*, VIII, 124
- Gordius aquaticus*, IX, 98
- Gortyna nebris*, Supp., 56
nitela, I, 92, III, 105, VIII, 37, Supp., 56
- Gothic Dart, I, 81
- gracilaria*, *Eunemoria*, Supp., 79
- Grafting, VI, 79
- Grain Bruchus, III, 45, 50, 51, 54
 Plant-louse, II, 5, 10, 16
 Sylvanus, III, 6
 Weevil, II, 10, III, 60
- grallator*, *Cryptus*, VII, 75
Labena, VII, 75
- graminis*, *Bombyx*, II, 44
- granarius*, *Bruchus*, II, 11, 14, III, 50, 51
Sitophilus, II, 10, III, 60
- Grand-daddy-long-legs, IV, 17
- Grandfather-Gray-Beards, IV, 17
- grandis*, *Lebia*, III, 100, Supp., 52
Stizus, I, 27, Supp., 52
- granulata*, *Tettix*, VIII, 150
- Granulated Grouse-locust, VIII, 150
- Grape-berry Moth, III, 90
 -cane curculio, III, 60
 Gall curculio, I, 131
 Remedy for, I, 132
- Codling, I, 133
 Remedy for, I, 135
- Curculio, I, 128, III, 60
- Fidia, V, 108
- Gall curculio, I, 181
- Gall-louse, III, 90, 92, 96, IV, 66
- grower, a new friend to the, III, 137
- leaf-louse, III, 88, 94
 Trumpet-gall, V, 118
- Grape Phylloxera, III, 84, IV, 55, 67, V, 57, 63, VI,
 30, VII, 41, 90, VIII, 157, IX, 43
- Bibliographical, VI, 30
- Biological, VI, 33
- Completion of its natural history, VII, 90,
 VIII, 157
- Conclusions, IV, 70, VI, 65

Grape Phylloxera—Continued.

- Different forms which the insect assumes, VI, 33, VII, 93
 Direct remedies, VI, 55, VII, 105
 Early existence in America, VI, 82, 83
 First appearance in California, VI, 82
 False theories, VI, 60
 Gall-inhabiting type, VI, 34, 66, 67
 Grafting as a means of counteracting the work of, IV, 65, VII, 108
 Impregnated egg not necessarily hibernial, VI, 86
 Injury caused by it in America, VI, 58, VII, 99
 France, VII, 103
 Its spread in Europe, V, 63, VII, 104
 Male louse, V, 71
 Means of contagion from one vine to another, IV, 64, V, 69
 Mode of spreading, VI, 45
 Mortality of vines caused by it, V, 57
 Natural enemies, VI, 50, VII, 106
 New theories, V, 67
 Occurrence in Southern States, VIII, 164
 Other preventive measures, VI, 50
 Practical considerations, IV, 67, VI, 44, VIII, 163
 Probable reasons why its injuries are greater in Europe than with us, IV, 66
 Prophylactic means of coping with the disease, VI, 48
 Range of the insect in America, V, 62, VI, 57, VII, 101
 Ravages of, in California, VI, 82, VIII, 163
 Real cause of disease, VI, 85
 Remedies, IV, 68, V, 71, VI, 55, VII, 105
 Resolutions concerning destruction of, VIII, 165
 Résumé of its natural history, IV, 69
 Root-inhabiting type, IV, 58, VI, 38, 66
 Sexed individuals, VII, 86, 98, VIII, 158
 Specific identity of the root-inhabiting and leaf-inhabiting types, IV, 57, VII, 94
 Specific identity of the American and European insects, III, 86, IV, 57
 Susceptibility of different vines to the disease, IV, 60, V, 64, VI, 46, VII, 106
 The more manifest and external effects of the disease, VI, 44
 Transient nature of the galls, V, 63
 Type *gallicola* or gall-inhabiting, VI, 34, 66, 67
 Type *radicicola* or root-inhabiting, VI, 38, 66
 Where do the winged females lay their eggs? VII, 96
 Why the insect is more injurious in Europe than here, VI, 59
- Grape-root-borer, I, 124
 Remedies for, I, 128
- Grape-seed Curculio, I, 129
 -seed Maggot, II, 92
- Grape-vine Apple Gall, V, 114, 115
 , Blue Caterpillars of, I, 136, II, 79
 Colaspis, III, 44, 62, 81, IV, 34
 Epimenis, III, 63, 65, VI, 87
 Flea-beetle, III, 79
 Fidia, I, 132, 133
 Filbert Gall, V, 116
 , Hog-caterpillar of, II, 71
 , Insects injurious to, I, 124, II, 71
- Grape-vine Gall-louse (see Grape Phylloxera)
 Leaf-folder, III, 61
 Leaf-gall-louse (see Grape Phylloxera)
 Plume, I, 137, III, 65, 66, 67, 68, IV, 129
 Procris, II, 85, V, 134
 Root-borer, I, 124, III, 75
 Tomato Gall, V, 117
 Trumpet Gall, V, 118
- Gropholitha oculana*, III, 6
Graphiphora, I, 79
Graptæ, III, 103, V, 149
Graptodera, Supp., 53
Grasserie, a disease of Silkworms, IV, 91
 Grasshoppers, a division of Orthoptera, V, 14
grata, *Euclyptus*, I, 136, II, 83, VI, 88, 89, 95
grataria, *Hæmatopis*, I, 119
 Gray, Alfred, on Rocky Mountain Locust, VII, 148, 149
 Gray's Improved Sprinkler, VII, 15
 Greasy Cut-worm, I, 80
 Great-crested Fly-catcher destroying Locusts, VIII, 124
 Lebia, III, 100
 Leopard moth, IV, 141
 Green Apple-leaf Tyer, IV, 46
 -head Fly, II, 128
 larva of White-lined Morning-sphinx, VIII, 122
 striped Locust, VIII, 149
 Maple Worm, V, 137
 Larval changes, V, 138
 Natural enemies, V, 139
 Remedies, V, 141
 Gregg, Jacob, on Rocky Mountain Locust, IX, 73
grossulariæ, *Pempelia*, I, 140, II, 9, Supp., 57
grossulariata, *Abraxas*, IX, 5
grossulariella, *Phycis*, Supp., 57
 Grote, A. R., on poisonous properties of *Doryphora*, VIII, 10, 11
 Ground-beetle larva preying on *Curculio* larvæ, I, 59
 larvæ preying on locust eggs, IX, 97
 The Elongate, I, 115
 The Fiery, I, 89, 115
 The Murky, I, 115
 The Pennsylvania, I, 59
 The subangular, I, 58
 Ground-beetles destroying Canker Worms, II 103
 Locusts, IX, 98
gryllaria, *Astoma*, VII, 175, Supp., 63
Gryllidæ, stridulating apparatus of, VI, 154
Gryllus erythropus, VII, 126
 niger, II, 152
Gubernaculum ovi, IX, 87
Guerinii, *Attacus*, IV, 112, 113
Guiraca ludoviciana, V, 54
guttata, *Cassida*, II, 60, 63
 Coptocycla, Supp., 53
- II.**
- Habit, change of, III, 91
 Hackberry Butterflies, VI, 136
Hadena, I, 68, Supp., 56
 amputatrix, I, 87
 chenopodii, Supp., 73, 77
 subjuncta, I, 84

- Haeckel, Prof. E., on the unity of nature, III, 174
Hæmatopis grataria, I, 179
 Hair-worms, IX, 93
Halesidota Harrisii, III, 127
 tesselata, III, 127
 Half-winged Bugs, characteristics of, V, 12
 Hall, William, on Chinch Bug and Locust, VIII, 76
Haltica chalybea, I, 101, III, 79, 81, Supp., 53
 cucumeris, I, 101, II, 57, V, 112, Supp., 53
 nemorum, I, 101
 pubescens, I, 101
 rhois, VI, 122
 stolida, VI, 122
 striolata, III, 44
 Ham-beetle, the Red-legged, VI, 96
 Hammond, A. C., on Apple-leaf Skeletonizer, IV, 45
hammondi, *Acrobasis*, III, 7
 Pempelia, IV, 44, 48, Supp., 80
 Hammond's Knot-horn, IV, 45
 Hanan, B., on Oyster-shell Bark-louse, V, 74
 Hand-maid Moth, III, 124
 Hanway, James, on Rocky Mountain Locust, VIII, 102
 Hardin, Gov. C. H., proclamation by, VIII, 95
 Harlequin Cabbage-bug, IV, 35
 Harman, M. B. W., on Rocky Mountain Locust, IX, 74
Harmonia picta, Supp., 52
Harpactor cinctus, I, 114, VII, 41, Supp., 58
Harpalus? Larvæ feeding on locust eggs, IX, 97
Harpalus caliginosus, I, 115, VIII, 52
 pennsylvanicus, I, 59, VIII, 52, IX, 98
 Harris, Dr. T. W., on hibernation of Disippus Butterfly, III, 155
 on Oyster-shell Bark-louse, V, 79
 on Poplar Spinner, II, 19
 on Tent-caterpillar, III, 121
harrisii, *Aspidiotus*, I, 7, II, 9, Supp., 60
 Halesidota, III, 127
 Harris's Bark-louse, I, 7, II, 9
 Harvest-flies, II, 131
 -men, IV, 17
 mites, VI, 122
 Hateful Locust, VII, 188, 190
 Hawk Moth, II, 76, IV, 86, V, 12
 Hay-worm, the Clover, VI, 102
 Hagen, Dr. H. A., on the distinction of Cicadæ, I, 21
 Head Maggot, I, 161
 Heard, J. M., Boll-worm Moth Trap, III, 109
 Heart-worm of cabbage, II, 107
 Heavy rains destructive to the Chinch Bug, II, 24
 Hecker, Fred., on remedy for Potato-beetle, VIII, 4
 Hedge-hog Caterpillar, IV, 143, 144
Hedya scudderiana, Supp., 57
Heliconidæ, III, 103
Heliconius melpomene, III, 173
 thelxiope, III, 173
Heliophila, VIII, 22
Heliothis armigera, III, 45, 104, IV, 129
 Hellebore for Currant worms, IX, 7, 13, 14, 15
 Potato bugs, IV, 14
 White Pine Worm, IX, 32
 Hellgrammite, V, 142, IX, 125
 Characters of the young larva, IX, 127
 Hellgrammite—Continued.
 Eggs hitherto supposed to belong to it, IX, 128
 Its curious egg-mass, IX, 126
 The larva lives in rapid flowing streams, IX, 128
 Where and how the eggs are laid, IX, 127
Helminthophaga ruficapilla, VI, 27
 Helius, Abraham, on Rocky Mountain Locust, VIII, 91
Hemileuca californica, V, 128
 maia, V, 127
 nevadensis, V, 128
Hemiptera, classification of, V, 12
 Preparing of for cabinet, V, 34
Hemirhipus fascicularis, VI, 101
Hemiteles cressonii, I, 177, Supp., 65
 nemativorus, IX, 17
 (?) *thyridopterigis*, I, 150, Supp., 65
Hentzii, *Mygale*, II, 106
herbimacula, *Celaena*, I, 86
herilis, *Agrotis*, Supp., 55
 Herschell, C., on Locusts, VIII, 151
herse, *Apatura*, VI, 136, 140, 148
 Herse Butterfly, VI, 148
Hesperia, VIII, 175
Hesperides, VIII, 176
 Hessian Fly, II, 10, 16, 17, 19, III, 110, 111, IV, 67, V, 13, 25, VII, 22, 27, 36
 parasite, III, 120
 Heterocera, a section of Lepidoptera, V, 12
 Heteromera, a section of Coleoptera, V, 10
 Heteroptera, a section of Hemiptera, V, 12
 Hewitt, H. L., on Rocky Mountain Locust, IX, 69
hibisci, *Bruchus*, Supp., 70, 71
 Hickman, G. B., on Rocky Mountain Locust, IX, 69
 Hickory Bark-borer, V, 62, 103, 104
 Natural enemies of, V, 106
 Remedies for, V, 107
 Hickory Borer, VI, 101
hilarana, *Oochylis*, I, 175, II, 135
 Hill, John, on Rocky Mountain Locust, IX, 75
Hipparchia, VI, 143
Hippobosca, VII, 91
Hippoboscidae, V, 13
Hippodamia convergens, I, 112
 glacialis, IV, 18
 maculata, I, 112, II, 25, V, 149, VII, 39, Supp., 52
 13-punctata, I, 112
histrionica, *Murgantia*, IV, 35
 Strachia, IV, 35
 Hoag, C. R., on Flat-headed Borer, VII, 74
 Hoag, I. N., on silk culture, IV, 80
Hockeria perpulcra, II, 53
 Hoffmeister, A. W., on Colorado Potato-beetle, VII, 14
 on Army Worm, VIII, 29
 Hogan, John H., on Grape-root Borer, I, 125
 Hog-caterpillar of the vine, II, 71, 75
 Hogs as Apple-worm destroyers, I, 65
 Curculio destroyers, I, 59
Holcocera glandulella, IV, 144, 145
 Holmes, Wm. C., on Tile-horned Prionus, II, 90
holosericeum, *Trombidium*, VII, 175
 Holsinger, Frank, on destroying Locusts, VIII, 127
Holt County Sentinel, article from, on Rocky Mountain Locust, VIII, 69

- Homely Geopinus, I, 89
Homeosoma, Supp., 57
Homoptera, a section of Hemiptera, V, 12
Homopus, VII, 106
Honey-bee, insect enemies of, I, 166, VI, 101
Honey locust seed-weevil, III, 45
Honora, Supp., 57
Hooten's Curculio-catcher, III, 22, 23
Hopkins, B. F., on Rocky Mountain Locust, VII, 151
Hoplophora arctata, VI, 53, 81, VII, 106
contractilis, VI, 54
Hopps, Michael, on remedy for Chinch Bug, II, 30
hordei, *Isosoma*, II, 92
Horned Lark destroying Locust eggs, IX, 91
Horned Passalus, IV, 139
eggs of, V, 55
Horner, C. C., on machine for destroying Locusts, VIII, 129
Hornet, sting of the, I, 27
Horn-tails, V, 10
Hostetter, C. J., on Rocky Mountain Locust, IX, 71
House Pigeon destroying Ca
Howard, Sanford, on remedy for Canker Worm, II, 100
How to collect, preserve, and study insects, V, 29
counterwork noxious insects, V, 23
transmit insects, V, 44
Huggins, J., on Canker Worm, II, 101
Hull, Dr. E. S. on Canker Worm, II, 101
on Grape-vine Flea-beetle, III, 81
on scab in Apples, II, 7
Hull's Curculio-catcher, III, 19
Modification of, V, 25
Hurlburt, J., on Colorado Potato-beetle, IV, 9
Husmann, Geo., on grape-vine grafting, VII, 109, 110, 111
pruning, VI, 84
on importance of Phylloxera discoveries, IV, 55
Huttoni, *Bombyx*, IV, 85
Hyalophora cecropia, IV, 103
Hybernia, VIII, 17
Hydrocorisa, a division of Heteroptera, V, 12
Hylecætus americanus, III, 7
Hymenoptera, characters and classification of, V, 9
Hyperchiria io, V, 133
varia, V, 133
Hyperaspis normata, V, 100
Hyphantria cunea, Supp., 55
punctata, Supp., 55
textor, III, 130, 132, Supp., 55
Hypogymna dispar, II, 10
Hypopus, VI, 52, VII, 106
dujardinii, VI, 53
Hypsopigia, VI, 105
- I.**
- Ichneumon brevipennis*, IX, 55
leucaniæ, II, 53, VIII, 54
obsoletus, IX, 55
pullatus, III, 69
signatipes, III, 69
subcyaneus, III, 69
unifasciatus, III, 71
Ichneumonidæ, III, 27, 28, Supp., 65
Icterus Baltimore, VI, 27
Icy Ladybird, IV, 18
Idol, J. K. P., on Rocky Mountain Locust, IX, 74
idyja, *Apatura*, VI, 145
ignota, *Rhodites*, I, 13
Illinois Lace-wing, II, 26, VII, 39
Locust flights in, VIII, 151
illinoensis, *Chrysopa*, II, 26, VII, 39, 40
Imbricated snout-beetle, III, 58
imbricator, *Pemphigus*, I, 121
imbricatus, *Epicærus*, III, 58
imbricornis, *Prionus*, II, 89, III, 6, 75
immarginatus, *Micropus*, VII, 22
Importance of Entomology as a study, V, 17
Imported Apple-worm, I, 108
Cabbage Worm, II, 107
Imported Currant Worm, II, 13, VI, 43, 149, IX, 7
Descriptive, IX, 21
It furnishes an interesting instance of defunctionation of special parts, IX, 19
It presents a forcible example of Arrenotoky IX, 18
Its introduction and spread, IX, 8
Its natural history, IX, 9
Natural enemies, IX, 17
Preventive measures, IX, 13
Remedies, IX, 13
Imported *vs.* Native American Insects, II, 8, 106, 107, VII, 5
Imported Onion-fly, II, 9
Oyster-shell Bark-louse, II, 9
Improved Patent Insect Destroyer, IV, 15
impura, *Leucania*, VIII, 38
inequalis, *Cæloides*, I, 128, Supp., 54
incertus, *Melanotus*, III, 6
Incrassated Geopinus, I, 77
incrassatus, *Geopinus*, I, 77
inda, *Euryomia*, III, 6
indagator, *Perilitus*, IV, 43, Supp., 66
indagatrix, *Pimpla*, IV, 43
indecisa, *Ægiale*, VIII, 179
Indigo-bird destroying Canker Worm, VI, 27
Indian Territory, locusts in, VIII, 88, IX, 76, 78
indiginella, *Myelois*, IV, 38
inermis, *Agrotis*, I, 72, 74, II, 50, III, 15, 129, VIII, 37
Supp., 55
infesta, *Celaena*, Supp., 56
Inflating Chalcis fly, I, 176
Influence of food in determining sex, VIII, 19
Influence of wind in determining the course of locust swarms, IX, 81
Ingalls, Senator, Introduction of bill in Congress for destruction of injurious insects, V, 133
Innoxious insects, I, 172, II, 125, III, 140, V, 142, VI, 127, VIII, 169, IX, 125
inornata, *Amphipyra*, III, 75
Tiphia, VI, 123
inquisitor, *Cryptus*, I, 150
Insecta, number of joints in, V, 7
Insect enemies of the Honey-bee, I, 166
domestication, IV, 85.
What is an? V, 5
Insects, Breeding, V, 41
Cabinet and boxes for, V, 37
How to counterwork noxious, V, 23

- Insects, How to collect, preserve, and study, V, 29
transmit, V, 42, 44
Imported and Native American, II, 8
Infesting the Apple-tree, III, 5, 6
Grape-vine, I, 124, II, 71, III, 61, IV, 53, V, 114, VI, 30
Potato, I, 91
Sweet-potato, II, 56
Mounting, for cabinet, V, 34
Rearing, V, 41
Relation of to agriculture, V, 18
Relaxing dry, V, 41
Text-books on, V, 42
Transmitting, V, 44
insidiosus, *Anthocoris*, II, 27, 32, VI, 51, VII, 41, Supp., 58
Insidious Flower-bug, II, 27, 32, VI, 51, VII, 41 47
Instinct, Curious, III, 156
Philosophy of, V, 83
vs. reason, V, 83, 157
interrupta, *Acronycta*, II, 121, Supp., 73
interruptus, *Passalus*, IV, 141
Io Moth, V, 133
Food plants, V, 136
Larval changes, V, 135
Parasites, V, 136
Iowa, Rocky Mountain Locust in, VIII, 81, IX, 63
iris, *Apatura*, VI, 136
Irrigation as remedy for Chinch Bug, VII, 31
Phylloxera, IV, 69, VI, 55
Rocky Mountain Locust, VII, 182
irritans, *Leptus*, VII, 177
Irritating Harvest Mite, VII, 177
property of caterpillars, V, 131
various insects, VI, 70
Irvine, Clarke, on Rocky Mountain Locust, VIII, 105, IX, 72
isabella, *Arctia*, IV, 143, Supp., 55
Isabella Tiger Moth, IV, 143
Iske, Anthony, Machine for destroying Potato-beetles, VIII, 4
isocrates, *Thecla*, VIII, 177
Isosoma hordei, II, 92
vitis, II, 92, 93
italicus, *Caloptenus*, VII, 133, VIII, 140
Ithomia, III, 161, 165
Ithycerus noveboracensis, III, 6, 57
Ituna, III, 103
- J.**
iaculifera, *Agrotis*, I, 82, 83, Supp., 56
Jarring as remedy for Apple-worm, IV, 25
Jefferson City Tribune, article from, on Army Worm, VIII, 50
article from, on Rocky Mountain Locust, VIII, 108
Jewett, D. T., on growing European grape-vines, VI, 77, 78
Johnson, B. F., on Locusts, VIII, 152
Johnson, C. V., on Colorado Potato-beetle, VIII, 10
Johnson, Prof. J. W., on Paris Green, VII, 11
Johnson, J. W., Sprinkling machine invented by, VI, 20
Joint-worm Fly, II, 92
Jones, H. L., on Rocky Mountain Locust, VII, 150
- Jordan, C. W., on Rocky Mountain Locust, IX, 70
Journal of Agriculture, article from, on Fall Army Worm, III, 109, 110
article from, on Pickle Worm, II, 69
article from, on remedy for Peach Borer, I, 49
juglandis, *Acrobasis*, IV, 42, 43, Supp., 67, 80
Conotrachelus, Supp., 54
Phycita, Supp., 80
Jumping Sumach-beetle, VI, 118
Natural history of, VI, 120
Remedies for, VI, 121
Jumping Tree-cricket, I, 138, V, 119
juncta, *Doryphora*, I, 103, 105, VII, 18, IX, 39
juvenalis, *Nisoniades*, III, 155
- K.**
kansanus, *Brachinus*, IV, 21
Kansas acts to encourage the destruction of Locusts, IX, 112, 113
Bombardier-beetle, IV, 21
Kansas City *Journal of Commerce*, articles from, on Rocky Mountain Locust, VIII, 59, 107
Times, article from, on Rocky Mountain Locust ravages, VIII, 74
Kansas Farmer, article from, on Army Worm, III, 110
article from, on remedy for Locusts, VII, 184
Kansas, legislation in, regarding Locusts, IX, 112, 113
Locusts in, VIII, 76, IX, 65
Silk culture in, IV, 82
State relief work, VIII, 78
Katydid, Angular-winged, VI, 155
Oviposition of, VI, 156
Eggs of, VI, 155, 158
Earlier stages of, VI, 158, 161
Song of, VI, 159
Natural enemies of, VI, 162
Katydid, Broad-winged, VI, 167
Oviposition of, VI, 167
Katydid, Narrow-winged, V, 124, VI, 164
Oblong-winged, V, 123, VI, 169
Katydid, VI, 150
Characteristics of, VI, 154
Oviposition of, VI, 155, VIII, 37
Stridulation of, VI, 154
Kaucher, Wm., on Rocky Mountain Locust, IX, 72
Kayser, A., on poisonous qualities of *Doryphora*, VIII, 11
Kedzie, Prof. W. K., on influence of Paris Green on soil, VII, 12, VIII, 6
on the use of Paris Green, V, 53, VIII, 6
Kelsey, S. J., on remedy for Chinch Bug, VII, 42
on Rocky Mountain Locust, VII, 135, 164, 193
Kerosene as remedy for Canker Worm, VIII, 20
Flat-head Apple-tree Borer, VII, 78
Locusts, VIII, 130
Killing insects intended for cabinet, V, 32
Kimbertin, J., on Rocky Mountain Locust, IX, 68

- King, Bennet, on Rocky Mountain Locust, IX, 72
 King Bird destroying Canker Worm, VI, 27
 Kirkpatrick, J., on Army Worm, II, 43
 Klæger, W., insect pins manufactured by, V, 34
 Klippart, H. J., on Army Worm, II, 47
 Knot-horn, Hammond's, IV, 45
 Knotweed Geometer, I, 179
kollari, *Ægiale*, VIII, 179
 Krimminger, W. B., on Rocky Mountain Locust, VIII, 63
Kuwa jirami, IV, 100
- L.**
- Labena grallator*, VII, 75
Labium of Hymenoptera, V, 9
 Lace-wing Fly, IV, 45
 The Illinois, II, 25
 The Weeping, II, 26, VI, 51
 Lacewing larva, I, 57, II, 27
Lachnosterna fusca, Supp., 53
 quercina, I, 57, 67, II, 19, V, 55, VI, 123, VII, 27, Supp., 53
 Ladybird, II, 25, VI, 51
 The Convergent, I, 112
 The 15-spotted, IV, 17
 The 9-spotted, I, 112
 The Spotted, I, 112
 The 13-spotted, I, 112
 Ladybirds, I, 112, II, 27, V, 11, 27
Laërtias, Supp., 54
Lagoa crispata, V, 126
 opercularis, V, 126
Lalage, V, 140
Lancaster Farmer, article from, on Colorado Potato-beetle, IX, 35
 Lance Rustic, I, 80
 Land Bugs, a division of Heteroptera, V, 12
lanigera, *Eriosoma*, I, 121, III, 95, IV, 69, VI, 63
 Schizoneura, Supp., 59
 Langworthy, A. J., on Apple-leaf Bucculatrix, IV, 50
 on Army Worm, VIII, 42
Laphygma frugiperda, II, 41
lardarius, *Dermestes*, VI, 100
 Larder-beetle, VI, 100
 Large Black Bear, IV, 141
 Larva, the second stage of insect development, V, 7
 Larvæ, Stinging, V, 125
Lasioptera vitis, V, 117
lateralis, *Corimelæna*, II, 35
 Ophion, II, 53
laticollis, *Prionus*, II, 87, III, 6, 75
latipennis, *Æcanthus*, Supp., 60, 61
 Latreille, life of, saved by an insect, VI, 101
Latreillia, V, 140
 Law of priority, IV, 55, VII, 143, VIII, 179, IX, 7
 Lay, J. H., on Rocky Mountain Locust, IX, 69
 Leaf-beetle, the Elm, VII, 5
 -bug, the Ash-gray, VII, 47
 -crumpler, III, 7
 the Rascal, IV, 38, VII, 81
 -folder, the Lesser, IV, 47
 -hoppers, III, 6
 mistaken for Locusts, VIII, 150
 -roller of the Strawberry, I, 142
 Leaf-rollers, III, 6
 -tyer, the Green Apple, IV, 46
 Least Pewee destroying Canker Worm, VI, 27
 Le Baron, Dr. Wm., description of Locust Mite by, VII, 176
 on *Aphelinus mytilaspidis*, V, 87
 on Apple Worm, VI, 10
 on Pine-leaf Scale, V, 84
 on remedies for Canker Worm, VI, 24
Lebia atriventris, VIII, 3, Supp., 52
 grandis, III, 100, VIII, 3, Supp., 52
Lecanides, V, 85, 92
Lecanium aceris, V, 92
 LeConte, Dr. J. L., on the use of mineral poisons as insecticides, VII, 8
lecontei, *Callimorpha*, III, 134, VI, 92, Supp., 55
 LeConte's Pine Worm, IX, 32
 Descriptive, IX, 33
 Habits of, IX, 33
 Legged Maple Borer, VI, 107
 Legislation against injurious insects, VIII, 132
 to avoid locust injury, IX, 111
 to create national entomological commission, VIII, 133
Lema trilineata, I, 99, II, 58, III, 14
 Leopard Moth, the Great, IV, 141
Lepidoptera, characteristics of, V, 11
 preparing of, for cabinet, V, 38
Lepidosaphes, I, 9, V, 91
leporina, *Acronycta*, II, 121, Supp., 73
Leptalis, III, 161, 165
Leptinotarsa, VIII, 2
Leptus, VI, 52
 americanus, VI, 122
 irritans, VI, 122, VII, 177
lepusculina, *Acronycta*, II, 121, Supp., 73
 Lesser Apple Leaf-folder, IV, 47, 48
lesueuri, *Chrysobothris*, VII, 71
 Letterman, G. W., on Colorado Potato-beetle, IX, 35
Leucania albilinea, IX, 50, 55
 comma, VIII, 43
 extranea, II, 50, VIII, 34
 harveyi, IX, 50
 impura, VIII, 38
 lithargyria, VIII, 38, 43
 phragmatidicola, IX, 57
 turca, VIII, 43
 unipuncta, I, 109, II, 5, 11, 37, 55, V, 25, VIII, 22, 24, 29, 182, IX, 47, Supp., 76
Leucaniæ, II, 51, 52
leucaniæ, *Ichneumon*, II, 53, VIII, 54
 Exorista, II, 50, 51, 52, III, 116, 129, IV, 108, VIII, 53, Supp., 60
Leucopis, VI, 51
leucoptera, *Curvirostra*, VI, 27
leucopterus, *Micropus*, II, 15, VII, 19, Supp., 58
leucostigma, *Orgyia*, I, 144
 Lewelling, H., on Apple Curculio, III, 35
Libellulæ, II, 46
Libellula trimaculata, V, 14
lictor, *Erax*, II, 124
 Lightning-hopper, Frosted, V, 122
luna, *Attacus*, IV, 123, 138
Limacodes, III, 150
 scapha, V, 126

- Lime, air-slacked, as remedy for Potato beetle, IV, 14
as remedy for Locusts, VIII, 130
Carbolate of, as remedy for Potato-beetle, IV, 14
Limenitis, III, 168, 169
 arthemis, III, 171
 bredowii, III, 171
 disippus, III, 153, 171, V, 146, Supp., 66
 lorquini, III, 171
 misippus, III, 167, VI, 145
 proserpina, III, 171
 sibylla, III, 171
 ursula, III, 163, 167, 171
 weidemeyerii, III, 171
limenitidis, *Microgaster*, III, 158, Supp., 66
Limneria fugitiva, IV, 41, V, 133, 141
 lophyri, IX, 32, Supp., 65
Lincecum, Dr. Gideon, on Harlequin Cabbage-bug, IV, 36
Lincoln (Neb.) *Journal*, article from, on Rocky Mountain Locust, VIII, 108
linearis, *Coreus*, II, 113
 Phytocoris, II, 113
lineata, *Deilephila*, III, 140
lineatella, *Prodenia*, Supp., 56
lineola, *Tabanus*, II, 128
Lintner, J. A., on cabinet boxes for Lepidoptera, V, 38
 on larva of *Eudryas unio*, VI, 93
 on Hog-caterpillar of the vine, II, 72
 on sexual characters in butterflies, III, 103
 on stinging larvæ, V, 131
Liparis auriflua, V, 126
List of birds which feed on Canker-worm, VI, 27
Lithacodes fasciola, V, 126
lithargyria, *Leucania*, VIII, 38, 43
Lithophane, Supp., 75
Little, Jos. T., on White Pine Worm, IX, 30
Little-lined Plant-bug, II, 113
Loafman, Dr. J. M., on Rocky Mountain Locust, IX, 75
Lobesia botrana, Supp., 57
Lobophora, IV, 129
Locust Flights east of the Mississippi, IX, 81
 Flights in Illinois, VIII, 151
 Mite, VII, 175
 The Atlantic Migratory, VII, 169
 The Clumsy, VIII, 148
 The Colorado, VII, 188
 The Differential, VII, 124, 173
 The Green-striped, VIII, 149
 The Hateful, VII, 188, 190
 The Migratory, VII, 133
 The Red-legged, VII, 125, 188
 The Rocky Mountain, VII, 121, VIII, 57, IX, 57
 The Seventeen-year, VII, 27
 The Thirteen-year, VII, 27
 The Two-striped, VII, 173
Locusta erythropus, VII, 126
Locustidæ, VI, 124, 150
Locusts, V, 14, VI, 153
 as food for man, VIII, 143
Locusts not a divine visitation, VIII, 97
 vs. grasshoppers, VI, 150, 153
Lodi Potato Pest Poison, VIII, 7
Lomechusa, IV, 22
Long, Dr., on remedy for Canker Worm, II, 101
Long, Levi, on Rocky Mountain Locust, VIII, 69, 102, IX, 71
Long-horned Beetles, injurious to fruit trees, III, 6
 stridulating noise produced by, III, 14
 Boring Beetles, larval habits of, II, 91
longifasciella, *Gelechia*, Supp., 83
longispinus, *Dactylopius*, IV, 70, VI, 63
Long-tailed Ophion, IV, 107, 129, V, 136
lophyri, *Limneria*, IX, 32, Supp., 65
Lophyrus, IX, 12
 abbotii, IX, 29, 32, Supp., 65
 abietis, IX, 33
 americanus, IX, 34
 compar, IX, 34
 fabricii, IX, 34
 lecontei, IX, 31, 32
lorquini, *Limenitis*, III, 171
Loxopeza, Supp., 52
Loxotænia rosaceana, III, 6
Lucanus elaphus, V, 145
Lukens, W. E., on Strawberry Leaf-roller, I, 143
luna, *Actias*, IV, 123
 Attacus, IV, 74
Luna Silkworm, IV, 123
 larval changes of, IV, 124
lunatus, *Cratoparis*, III, 10
Lycænidæ, VI, 138
lycaon, *Apatura*, VI, 136, 146, 148
 Pholus, II, 76
 Sphinx, II, 76
lycarum, *Agrotis*, Supp., 77
Lycorea, III, 103
Lycosa, IX, 98
Lyctus opaculus, IV, 54
Lyda, IX, 10
Lydella doryphoræ, I, 111, IV, 6, IX, 40, Supp., 88
Lygæus leucopterus, VII, 21
Lygus robinæ, V, 154
Lyman, Jos. B., on Cotton Worm, II, 39, 40, Supp., 88
Lymexylon navale, V, 19
Lytta ænea, III, 6
 atrata, I, 98, Supp., 54
 cinerea, I, 97, Supp., 54
 fabricii, I, 99
 marginata, I, 98, Supp., 54
 murina, I, 98, Supp., 54
 vittata, I, 96

M.
Machines for catching Curculio, I, 60, III, 18, V, 25
 Colorado Potato-beetle, I, 117, VI, 14, VIII, 4
Macrobasis unicolor, Supp., 54
Macrocentrus delicatus, V, 50
Macroductylus subspinosus, V, 108
macrurum, *Ophion*, IV, 107
maculalis, *Desmia*, III, 61
maculata, *Hippodamia*, I, 112, II, 25, VII, 39, Supp., 52
 Vespa, Supp., 48
maculosa, *Dendroica*, VI, 27

- Madarus ampelopsidis*, I, 132
vitis, I, 131, Supp., 71
magnus, *Brachypeplus*, VIII, 148
Maia Moth, V, 127
 Its Eggs, V, 128
 Issuing of the moth, V, 132
 The larva, V, 129
 Larval changes, V, 129
 The pupa, V, 131
 The sting of the larva, V, 131
 Natural enemies, V, 132
maia, *Saturnia*, IV, 41
Maine Farmer, article from, on Canker Worm, II, 100
mali, *Acarus*, II, 6
mali, *Aphis*, II, 6, 10, III, 6
 Molobrus, III, 6
malivorana, *Tortrix*, IV, 47, Supp., 82
malus, *Acarus*, I, 16
malvarum, *Erynnis*, VIII, 182
Mamestra, I, 68, III, 123
 picta, II, 112
Mandibulæ of Hymenoptera, V, 9
Mandrake as remedy for Potato Bugs, IV, 15
Manitoba, Rocky Mountain Locust in, VII, 155, VIII, 89
Mankato Review, article from, on Rocky Mountain Locust, IX, 60
Mann, B. P., on Canker Worm, VI, 28
Mantis carolina, I, 169, III, 68, IX, 98
mantivora, *Sarcophaga*, VII, 180
Many-banded Robber, I, 114, VII, 41
Maple, J. W., on Rocky Mountain Locust, IX, 71
Maple Borer, the Legged, VI, 107
Maple Worm, the Green-striped, V, 137
 Enemies of, V, 139
 Larval changes of, V, 138
 Remedies for, V, 141
marginata, *Lytta*, I, 98, Supp., 54
marginatum, *Trochilium*, Supp., 72
Margined Blister-beetle, I, 98
mariae, *Chalcis*, IV, 109, 110, 123
Mark, Calvin A., on Rocky Mountain Locust, VIII, 102
Marsupialea, II, 12
Mary Chalcis-fly, IV, 109, 129
Maryland Yellow-throat, destroying Canker Worm, VI, 27
Mascicera, III, 150, IV, 129
maxillosus, *Brenthus*, VI, 116
Maxwell, J. H., on Rocky Mountain Locust, IX, 69
Maxwell, J. R., on the use of Paris green, VI, 19
May Apple for Potato Bugs, IV, 15
May Beetle, I, 156, II, 19, IV, 16, VII, 27
 Eggs of, V, 55
May-flies, III, 8, V, 143
Maynard, C. J., list of birds destroying Canker Worm, VI, 27
McAfee, H. H., on hand-picking as remedy for Potato Bugs, IV, 11
McCartney, Jno. P., on White Grub, I, 156
McNallie, Thomas, on Rocky Mountain Locust, IX, 72
Meador, W. R., on Rocky Mountain Locust, IX, 75
Meadow Grasshopper, V, 123
Meal-worm, II, 10, 11, VI, 118
Mealy Bug, V, 80
Meat-flies, II, 19, VII, 27
Meeker, N. C., on Rocky Mountain Locust, VIII, 84
Megarthus, IV, 22
Megathymus yuccæ, VIII, 169, 171, 179, IX, 129
Megilla, Supp., 52
melancholica, *Euryomia*, III, 6
Melanerpus erythrocephalus, VIII, 124
melanocephala, *Pimpla*, III, 129
Melanoplus, VII, 188
melanosus, *Micropus*, VII, 22
melanothorax, *Doryphora*, VIII, 10
Melanotus communis, III, 6
 incertus, III, 6
mellonella, *Galleria*, Supp., 57
melodia, *Melospiza*, VI, 27
Meloe, VI, 125
Melospiza melodia, VI, 27
melpomene, *Heliconius*, III, 173
melsheimerii, *Perophora*, V, 125
Membracididæ, a family of Homoptera, V, 122
Membrane-winged Flies, an Order of Insects, V, 9
Meracantha contracta, VI, 118
Merchant, J. W., on Oyster-shell Bark-louse, V, 77
Merchants' Exchange, (St. Louis) relief committee from, VIII, 93
merdigera, *Crioceris*, II, 58
Merdigerous habit in insects, II, 58, VI, 128
Mermis, V, 49
Meromyza americana, I, 159
Merritt's Patent Tree-protector, II, 99
Meske, Otto, on Army Worm, VIII, 44
Mesochorus fuscipennis, VII, 75
 vitreus, II, 52, VIII, 53
messoria, *Agrotis*, Supp., 77
Metagenetic Coccids, V, 84
Metzler, T., on grape culture, VI, 83
Microcentrus retinervis, VI, 155, 156
Microgaster, I, 89, II, 75, 102, III, 158, IV, 45, V, 133
 VIII, 54, Supp., 66
 acronyctæ, II, 120
 gelechicæ, I, 178, Supp., 66
 glomeratus, III, 167
 limititidos, III, 158, Supp., 66
 militaris, I, 89, II, 52, VIII, 53, Supp., 52
Micropteryx, III, 7
 pomivorella, III, 7
Micropus leucopterus, II, 15, VII, 19, Supp., 58
 var. *albivenosus*, VII, 22
 apterus, VII, 22
 basalis, VII, 22
 dimidiatus, VII, 22
 femoratus, VII, 22
 fulvivenosus, VII, 22
 immarginatus, VII, 22
 melanosus, VII, 22
 nigricornis, VII, 22
 rufipedis, VII, 22
Migration of butterflies, III, 152
 locusts, conditions of, VIII, 112
migratoria, *Ædipoda*, VII, 133, VIII, 145
migratorius, *Turdus*, VI, 27
Migratory Locust, VII, 133
milbertii, *Proctacanthus*, Supp., 88
militaris, *Exorista*, II, 50, III, 129, IV, 109
 Microgaster, I, 89, II, 52, VIII, 53, Supp., 52

- militaris*, *Senometopia*, II, 50
 Military Microgaster, II, 52, 103, VIII, 53
 Miller, J. B., on Strawberry Crown-borer, III, 42
 Miller, Samuel, on grape culture, VI, 76, 84
 on Locusts, VIII, 154
 Miller, Stephens, on Rocky Mountain Locust, IX, 92
 Miller, W. H., on Rocky Mountain Locust, VIII, 112
 Milliken, R. M., on Canker Worm Trap, VI, 25
 Milliken, R., on Rocky Mountain Locust, VII, 150, IX, 65
 Mimetic analogies, III, 163
 resemblance, V, 147
 Mimicry in animals, III, 159
 butterflies, III, 159
Mimus carolinensis, VI, 27, VIII, 124
minimus, *Empidonax*, VI, 27
 Pezomachus, II, 52, VIII, 54
ministra, *Datana*, III, 124, 127, 129, IV, 129
minuta, *Eupsalis*, VI, 113, 117
 Trichogramma, III, 158, Supp., 68
 Minnesota, act for the destruction of Locusts, IX, 113
 Legislation regarding Locusts, IX, 114
 Locust Commission, VIII, 83
 Locust history in, VIII, 81, IX, 60
 Locust invasion of 1876 in, IX, 60
mirabilis, *Antigaster*, VI, 162, Supp., 52
Mirror and Farmer, article from, on Locusts, VII, 172
misella, *Chrysobothris*, VII, 71
missippus, *Limenitis*, III, 167, VI, 145
 Misnamed Gall-moth, II, 134
 Mississippi, Locust flights east of the, IX, 81
 Missouri, act to encourage the destruction of Locusts, IX, 111
 Democrat, article from, on Army Worm, VIII, 50
 Legislation regarding Locusts, IX, 112
 Locusts in, VIII, 89, 90, IX, 66
 Silk-growing in, IV, 83
 Missouri Bee-killer, II, 121, 122, IV, 21
missouriensis, *Asilus*, II, 121, 122, IV, 21, Supp., 87
 Mitchell, Th., on Army Worm, VIII, 39
 Mite, the Locust, VII, 175
 The Mussel-shaped, VII, 106
 The Phylloxera, VII, 106
 The Silky, VII, 175
 Mites, V, 87
 transformation of, VI, 52, 53
modesta, *Arma*, V, 133
 Modest Soldier-bug, V, 133
 Modrel, M. L., on Rocky Mountain Locust, IX, 75, 76
molesta, *Myrmica*, II, 11, IX, 43
molitor, *Tenebrio*, II, 9, 10, VI, 118
 Mollusca, a branch of the Animal Kingdom, V, 6
Molobrus mali, III, 6
molochinus, *Quedius*, IV, 21
Molothrus pecoris, VI, 27
 Molts, mode of enumerating larval, III, 145
Monoleuca semifascia, V, 126
 Monomera, a division of Heteroptera, V, 13
 Montana, Locust history in, VIII, 87, IX, 59
 Montgomery, G. W., on Rocky Mountain Locust, IX, 70
mori, *Bombyx*, IV, 74, 75, 84, 138
 Sericaria, IV, 75
 Mormon Louse, II, 17
 Morris, Rev. Dr. J. G., on Periodical Cicada, IV, 31
 Mortality among Grapevines, V, 58
 Mosquito, V, 13
 Mosquito Hawks destroying Army Worm moths, II, 46
 Moths, a section of Lepidoptera, V, 12
 Motsinger, J. L., on Rocky Mountain Locust, IX, 73
 Mottled Tortoise-beetle, II, 60, 63
 Moulton, J. T., jr., on Chinch Bug, VII, 33
 Muench, Hon. F., on Grape mortality, V, 58
 Muhleman, J. R., on Chinch Bug, VII, 34
 on Fall Web-worm, III, 131
 on Mimicry in Insects, III, 160
 on Raspberry Root-borer, VI, 112
 Mulberry Silkworm, IV, 75, 114, 134, IX, 18
 Best food for, IV, 100
 Best varieties or races, IV, 91
 Choking the chrysalis, IV, 96
 Chrysalis of, IV, 87
 Cocoon, IV, 87
 Cocoonery, IV, 96
 Culture in California, IV, 79
 Kansas, IV, 82
 Missouri, IV, 83
 Egg, IV, 86
 Egg-laying, IV, 97
 Enemies and diseases, IV, 87
 How best to rear, IV, 92
 Its history in America, IV, 77
 Its past history, IV, 75
 Larva, IV, 86
 Moth, IV, 87
 Natural history, IV, 84
 Osage orange as food for, IV, 100
 Reeling, IV, 98
 Mulsant, E., insect pins used by, V, 35
munda, *Coccinella*, II, 25, VII, 39, Supp., 52
 Murgantia, IV, 35
 Mungor, H. A., on Tile-horned Prionus, II, 90
murænula, *Agrotis*, I, 78
murcimaculata, *Celaena*, Supp., 54
Murgantia histrionica, IV, 35
 munda, IV, 35
 Muriate of lime, Gould's, as a remedy for Canker Worm, II, 100
murina, *Fidia*, Supp., 53
 Lytta, I, 98, Supp., 54
 Murky Ground-beetle, I, 98, 115
 Murray, A., on Protective mimicry, III, 170
 Murtfeldt, Miss M. E., on Oviposition of Narrow-winged Katydid, VI, 165
Musca bifasciata, V, 140
 domestica, II, 10
 Muscardine, a disease of Silkworms, IV, 88
Muscidae, III, 150, VII, 175
 Mussel-shaped Bark-louse, I, 15
 Mite, VII, 106
muticus, *Scolytus*, V, 105, 107
Myiarchus crinitus, VIII, 124

- Myelois convolutella*, Supp., 57
 indiginella, IV, 38
 suavella, Supp., 79
Mygale, IX, 98
 Hentzii, II, 106
 Mygatt, E. G., on Remedies for Bark-lice, I, 18
Myiodioctes canadensis, VI, 27
mylitta, *Attacus*, IV, 138
Myobia, VI, 52
Myocoryna 10-lineata, VII, 16
 Myriapoda, a class of Articulate animals, V, 6
Myrmica molesta, II, 11, IX, 43
myron, *Darapsa*, II, 71
 Otus, II, 71
 Sphinx, II, 71
Mysia 15-punctata, IV, 18, Supp., 53
Mytilaspis conchiformis, V, 93
 linearis, V, 93
 pini, V, 98
 pinifoliae, V, 97
 pomicorticis, V, 73, 95, Supp., 85
 pomorum, V, 93, 94
- N.**
- nana*, *Trogosita*, III, 6
napi, *Ceutorhynchus*, III, 11
 Narrow-winged Katydid, V, 124, VI, 164
 Descriptions of adolescent stages, VI, 166
 Oviposition, VI, 165
 Nashville Warbler destroying Canker Worm, VI, 27
nasicornis, *Oryctes*, VI, 124
 National Agricultural Congress, Address before, VI, 17
 National Academy of Sciences, proposed Entomological Commission under, VII, V
 Native American Onion-fly, II, 9
 Native Currant Worm, IX, 23
 Descriptive, IX, 28
 Its habits, IX, 24
 Remedies, IX, 28
 Wherein it differs from the imported species, IX, 23
 Native home of the Colorado Potato-beetle, VIII, 8
 Rocky Mountain Locust, VIII, 109
 Natural selection, remarks on, III, 159, IV, 84, V, 83
Naturaliste Canadien, article from, on Polyphemus Silkworm, IV, 128
Nazara, IV, 20
 Neat Cucumber Moth, II, 68
 Nebraska Bee-killer, I, 168, II, 122, IV, 2
 , Locusts in, VIII, 79, IX, 64
nebris, *Gortyna*, Supp., 56
nebulella, *Acrobasis*, IV, 42, Supp., 80
nebulo, *Acrobasis*, IV, 38, 47, Supp., 79
 Phycita, III, 7, IV, 38, 41, Supp., 79
Necrobia ruficollis, VI, 101
 Necrophaga, a division of Coleoptera, V, 11
Necrophoridae, III, 14
 Needham, H. V., on Chinch Bug ravages, VII, 37
 Negro-bug, II, 35
 Flea-like, VII, 48
Nematus, Supp., 59
 affinis, IX, 7
 grossulariae, IX, 7
 grossulariatus, IX, 7
Nematus, ribesii, IX, 7
 salicis-pomum, IX, 20
 trimaculatus, IX, 7
 ventricosus, VI, 43, 149, IX, 7, 10, 21
 Nemocera, a section of Diptera, V, 13
Nemoræa, V, 140, Supp., 60
nemorum, *Hattica*, I, 101
 Phyllotreta, III, 83
Nemotois scabrosellus, V, 160
nenuphar, *Conotrachelus*, I, 50, III, 11, 28, 31, 127, Supp., 54, 65, 68
 Nerve-winged insects, V, 14
 Neuroptera, characteristics and classification of, V, 14
neustria, *Olisiocampa*, III, 119
 New England Farmer, article from, on Potato-bug in Massachusetts, IV, 8
 Grape-root Borer, II, 87
 York Tribune, article from, on Clover Worm, VI, 104
 article from, on Grape Phylloxera, VI, 85, VII, 94
 article from, on grape-vine grafting, VII, 109
 article from, on Potato-beetle, IX, 35
 article from, on remedy for Canker Worm, II, 101
 article from, on Rocky Mountain Locust, VII, 189, VIII, 156, IX, 64
 York Weevil, III, 6, 57
ni, *Plusia*, II, 112, Supp., 78
niger, *Gryllus*, VI, 152
nigricans, *Agrotis*, I, 81, 83, 87
nigricornis, *Micropus*, VII, 22
 Æcanthus, Supp., 61
nigripes, *Cassida*, II, 63, Supp., 53
 Nine-spotted Lady-bird, I, 112
Nisoniades juvenalis, III, 155, VIII, 177
 Nitchy, F. A., on Fall Army-worm, III, 109
nitela, *Gortyna*, I, 92, III, 105, VIII, 37, Supp., 56
nitidalis, *Phacellura*, II, 7, 64, 68
niveus, *Æcanthus*, I, 138, V, 120, Supp., 60, 61
Noctua clandestina, I, 79, Supp., 55
 unipuncta, II, 49, VIII, 34
Noctuidæ II, 45, III, 136
 Nomenclature, II, 71, III, 94, 133, 143, IV, 55, VI, 109, 150, 153, VII, 16, 89, 143, 187, VIII, 22, 179, IX, 50
 Need of popular terms that will not confuse, VI, 153, VII, 187
 Law of priority, IV, 55, VII, 143, VIII, 179, IX, 7
 Popular vs. scientific names, VII, 187
Nonagria, VIII, 38
 Northern Army-worm, VIII, 24
 Brenthian, VI, 113
 Northern Squash-beetle, IV, 18
Nothrus ovivorus, II, 102
novæboracensis, *Ithycerus*, III, 6, 57
novem-notata, *Coccinella*, I, 112
 Noxious Insects, I, 7, IV, 72, V, 46
 How to counterwork, V, 23
nucum, *Balaninus*, III, 11
nuncius, *Cryptus*, IV, 110, 111, 123, Supp., 52
Nycteribidæ, V, 14
Nymphalidæ, III, 167, VI, 138
Nymphalis disippus, II, 125

- Nysius angustatus*, V, 113, Supp., 85
destructor, V, 111, 113, VII, 190, Supp., 84, 85
raphanus, V, 111
thymi, V, 113, Supp., 85

O.

- Oak-feeding Tortricid, VI, 103
Oak Phylloxera, the American, VI, 64, VII, 118
Oak-pruner, I, 25
Oarisma poweshiek, VIII, 178
Oberea ocellata, I, 178, 179
perspicillata, VI, 111
obesa, *Amara*, Supp., 52
oblinita, *Acronycta*, III, 70, 71
oblineatus, *Capsus*, II, 113, VII, 27
oblonga, *Celæna*, III, 136, Supp., 75
oblongifolia, *Phylloptera*, II, 57, VI, 169
Oblong-winged Katydid, II, 57, V, 123, VI, 169
obscura, *Prodenia*, III, 117
Ohrysobothris, VII, 71
Laphygma, VIII, 49
obscurus, *Prionus*, I, 127
Tenebrio, II, 9, 11
obsoletus, *Bruchus*, III, 54, 56, Supp., 70, 71
occidentalis, *Acronycta*, II, 121, Supp., 73
Caloptenus, VIII, 116
ocellata, *Oberea*, I, 178, 179
octomaculata, *Alypia*, I, 136, II, 80, 82, VI, 88, 94, 95
oculana, *Grapholitha*, III, 6
Ecanthus bipunctatus, Supp., 61
californica, Supp., 61
fasciatus, Supp., 60
latipennis, Supp., 60, 61, 62
nigricornis, Supp., 61
niveus, I, 138, V, 120, Supp., 60, 61
varicornis, Supp., 61
Edipoda, VIII, 103, 128
atrox, VII, 124
carolina, VII, 175, 179
differentialis, V, 14
migratoria, VII, 133, VIII, 145
phænicoptera, VIII, 104
Æstrus, III, 150
ovis, I, 161
Æta compta, I, 151, Supp., 58
Eggs of, Supp., 58
Oil of cade as remedy for Grape Phylloxera, IV, 68
olens, *Goerius*, IV, 21
oleracea, *Pieris*, II, 105
olinalis, *Asopia*, VI, 103, 107
olivaceus, *Vireo*, VI, 27, VIII, 125
Omaha Bee, article from, on Rocky Mountain Locust, VIII, 80
Omaha Conference on the Locust subject, IX, 106
Oncideres cingulatus, III, 6
Onion Fly, I, 155
-maggot, V, 13
opaculus, *Lyctus*, IV, 54
Ophion, II, 130, IV, 123
bilineatus, III, 69
lateralis, II, 53
macrurum, IV, 107
purgatus, II, 53, VIII, 54
Ophion, the Long-tailed, IV, 129
Ophiusa xyliua, II, 40
Orchard Tent-caterpillar, III, 129

- Orchelimum*, VI, 155
glaberrimum, V, 123, Supp., 62
Orgyia leucostigma, I, 144
Oribatidæ, VI, 54
orientalis, *Blatta*, II, 10
orizivora, *Dolichonyx*, VIII, 52
ornata, *Strachia*, IV, 38
Orocharis saltator, I, 138, V, 119, Supp., 60, 62
Eggs of, Supp., 62
Stridulation of, Supp., 62
Ortalis arcuata, II, 9
Orthia, VIII, 178
Orthoptera, characteristics and classification of, V, 14
Orthosoma cylindricum, I, 124, 127, II, 87
Orthotylus discoidalis, V, 154
Orton, Edward, on Legged Maple-borer, VI, 109
ortonii, *Agrotis*
Ortyx virginiana, II, 28, VII, 41
Oryctes nasicornis, VI, 124
oryzivorus, *Dolichonyx*, VI, 27, VIII, 52
Oscinis vastator, I, 161
Osborne, Gov., Proclamation of regarding Locust ravages, VII, 149
Osten-Sackenii, *Exorista*, II, 51
ostreæformis, *Diaspis*, Supp., 60
Otiiorhynchus sulcatus, III, 11
Otus notus, II, 71
myron, II, 71
Ovipositor of *Æschna*, VIII, 36
Agriion, VIII, 36
Army Worm Moth, VIII, 32, 38, 39, 182
Canker-worm Moth, VIII, 37
Fall Army-worm, VIII, 37
Katydid, V, 154, VIII, 37
Plum Curculio, VIII, 36
Unarmed Rustic, VIII, 37
Yucca Moth, VIII, 37
ovis, *Æstrus*, I, 161
ovivorus, *Nothrus*, II, 102
Owlet Moth, II, 10, 45, 119
Oyster-shell Bark-louse, I, 7, II, 6, 10, 14, V, 26, 73, Supp., 86
Bibliographical and Descriptive, V, 91
Both single and double brooded, V, 79
Enemies and parasites, I, 16, V, 87
Food Plants, I, 15, V, 86
Formation of the scale, I, 12, V, 80
Its occurrence in Missouri, V, 74
Its range south, V, 77
Its spread westward, V, 79
Male Louse, V, 83
Mode of growth, I, 10
Mode of spreading, I, 15, V, 85
Remedies, I, 16, V, 90
Southern limits in Missouri, I, 8
True nature of the scale, V, 80
oxycoccana, *Teras*, Supp., 83
Tortrix, Supp., 83
Oxyptilus, Supp., 58

P.

- Pachymerus vulnerator*, IV, 28
Pachyrhynchus Schænerri, III, 57
Packard's "Guide to the Study of Insects," V, 44

- Pædisca affusana*, Supp., 57
Painesville (Ohio) *Telegraph*, article from, on Potato-beetle, VI, 12, 13
Painted Borer, III, 7
 Lady, III, 151
 Lady-bird, V, 101
 Mamestra, II, 113
 -winged Digger-wasp, V, 149
Paleacrita—A new genus for the Spring Canker-worm, VIII, 13, Supp., 58
Paleacrita vernata, VIII, 13
Pale-thighed Tortoise-beetle, II, 62
Palingenia, IX, 128
pallida, *Cassida*, II, 62
Palmer, R. B., on Oyster-shell Bark-louse, V, 77
Palson, W. D., on Chinch Bug, VII, 22
paludana, *Tortrix*, VI, 103
pandorus, *Daphne*, II, 76
Paniscus geminatus, I, 89
pampinatrix, *Chærocampa*, II, 71
paphia, *Antheræa*, IV, 138
Paphia, V, 149, VI, 138
 glycerium, II, 125, 127, V, 145
Papilio, II, 117, V, 146
 asterias, III, 169
 philenor, II, 116, III, 169, Supp., 54
 troilus, III, 169
 turnus, VI, 145
Papilionidæ, VI, 138
paradoxus, *Rhipiphorus*, VI, 125
Parallel Pruner, III, 6
parallelum, *Elaphidion*, III, 6, IV, 54
Parasa chloris, V, 126
Parasites, artificial propagation of, III, 29, IV, 40, V, 90
 artificial introduction of, VII (preface, p. 4)
 of Abbot's White Pine Worm, IX, 31
 Archippus Butterfly, III, 149
 Army Worm, I, 89, II, 50, VIII, 53
 Bag-worm, I, 150
 Buck Moth, V, 133
 Cabbage Worm, II, 109
 Cecropia Worm, IV, 107
 Codling Moth, IV, 28, V, 49
 Colorado Potato-beetle, I, 111, IX, 40
 Common Yellow Bear, III, 69
 Cottonwood Dagger, II, 120
 Cut-worms, I, 89
 Disippus Butterfly, III, 157
 Flat-headed Borer, VII, 74
 Goatweed Butterfly, V, 149
 Gooseberry Span-worm, IX, 6
 Grape Curculio, I, 129
 Green-striped Maple Worm, V, 139
 Hickory Bark-borer, V, 106
 Hog-caterpillar of the Vine, II, 73
 Imported Currant Worm, IX, 17
 Io Moth, V, 136
 Katydid, VI, 162
 Lasioptera vitis, V, 118
 Luna Silkworm, IV, 125
 Oyster-shell Bark-louse, V, 87
 Pine-leaf Scale-insect, V, 100
 Plum Curculio, III, 24
 Polyphemus Silkworm, IV, 129
 Promethea Silkworm, IV, 123
Parasites of Rascal Leaf-crumpler, IV, 40
 Rocky Mountain Locust, VII, 174, IX, 91
 Smeared Dagger, III, 71
 Solidago Gall Moth, I, 175
 Tawny Emperor Butterfly, VI, 142
 Walnut Case-bearer, IV, 43
 Wheat-head Army Worm, IX, 55
 White Grub, VI, 123
 White-lined Morning Sphinx, III, 142
 Yama-mai Silkworm, IV, 136
 Yucca Borer, VIII, 179
Parasitism, VI, 123
 Secondary, VI, 128
Paris Green, antidote for, IV, 13
 Contrivance for dusting, IV, 2
 Cost of application per acre, VIII, 3
 Experiments with, VIII, 6, 7
 for Canker Worm, VI, 26
 Colorado Potato-beetle, IV, 11, V, 53, VII, 8, VIII, 3
 Cotton Worm, VI, 17, VII, 9
 Curculio, III, 18
 Flat-headed Borer, VII, 78
 Locusts, VII, 183
 Influence of, on man through the soil or through the plant, VII, 13
 Influence of, on the plant, VII, 11
 the soil, VII, 11, VIII, 6
 Machines for spraying, VI, 20, VII, 15, VIII, 4, 5
 Patents on, VI, 20
Parker, Dr. S. J., on mortality among grape-vines, V, 59
Parrish, W. S., on Rocky Mountain Locust, IX, 73
Parthenogenesis, IV, 86, V, 85, VI, 35, IX, 18
Parula americana, VI, 27
Parus atricapillus, IV, 107, VI, 27
Pasimachus elongatus, I, 115, VIII, 52, IX, 98
 punctulatus, IX, 98
Passalus cornutus, IV, 139, 140, V, 7, 55
 distinctus, IV, 141
 interruptus, IV, 141
Passalus, the Horned, IV, 139
Patent remedies and nostrums, IV, 15
Patents on insecticides, VI, 21
Pauls, G., on Fall Army-worm, III, 111
 on Rocky Mountain Locust, VII, 173
Payne, M. S., on Rocky Mountain Locust, VIII, 68
Payne S. D., on Rocky Mountain Locust, VIII, 126
 on Enemies of Rocky Mountain Locust, IX, 96
Pea Bug, II, 11, III, 45
Pea-weevil, III, 44, 55, IX, 43
 The female deposits her eggs on the outside of the pod, III, 46
 Remedies, III, 48
Peach Borer, I, 47, II, 11, III, 76, 77, VI, 108
 Remedies for, I, 48
Peach-worm, the Blue-spangled, III, 132
Pear Blight, III, 58
Pear-tree Flea-louse, II, 10
Pearl Wood Nymph, II, 80, 83, 84, III, 63, VI, 90
Pébrine, a disease of Silkworms, IV, 88, 89, 90, 91
Peck, W. D., on Canker Worm, VII, 89
Pecks' Spray Machine, VIII, 4

- pecoris*, *Molothrus*, VI, 27
pectinatus, *Rhipiphorus*, VI, 125
Pelidnota punctata, III, 77, 78
pellionella, *Tinea*, II, 10
Pelopæus, V, 157
Pempelia, IV, 46
 grossulariæ, I, 140, II, 9, Supp., 57
 hammondi, IV, 44, 46, Supp., 80
Pemphigus, III, 96
 caryæcaulis, VII, 117
 caryævenæ, VII, 117
 imbricator, I, 121
 vagabundus, I, 112, 120, VII, 97
 vitifoliæ, I, 13, III, 85, 93, VI, 31, VII, 94, 117
 Pendleton, E. M., on Rocky Mountain Locust, IX, 85
 Pennsylvania Ground-beetle, I, 59, IX, 98
 Soldier-beetle, IV, 28
 -bug, I, 57
pennsylvanica, *Dendroica*, VI, 27
 Epicauta, Supp., 54
pennsylvanicus, *Chauliognathus*, I, 57, IV, 28, Supp., 53
 Harpalus, I, 59, VIII, 52
 Pentamera, a section of Coleoptera, V, 10
Pentarthron, Supp., 69
Pentarthrum, Supp., 69
Pentatoma rufipes, IV, 20
Penthina vitivorana, I, 133, Supp., 57
Pepsis formosa, II, 106
peregrinum, *Acridium*, VII, 133, VIII, 144, 145
Perilampus platygaster, II, 87
Perilitus indagator, IV, 43, Supp., 66
Perillus circumcinctus, IV, 19
 Periodical Cicada, I, 18, III, 6, IV, 30
 Chronological history, with predictions of the future appearance of broods, I, 30
 Enemies, I, 26
 Injury, to fruit trees, I, 29
 Natural history and transformation, I, 22
 Season of appearance and disappearance, I, 22
 17- and 13-year broods, I, 18
 Sting of, I, 26
 Two distinct forms, I, 20
periscelidactylus, *Pterophorus*, I, 137, III, 65, Supp., 58
Peritymbia vitisana, IV, 55, VI, 31, VII, 117
 Perkins, Geo. H., on Colorado Potato-beetle, VIII, 2
Perla, V, 143
pernyi, *Antheræa*, IV, 137
 Attacus, IV, 74, 137, 138
 Perny's Silkworm, IV, 137
Perophora Melsheimerii, V, 125
perpulchra, *Hockeria*, II, 53
persimilata, *Acidalia*, VI, 138
Perthostoma, IX, 129
 Peterson, J. M., on Rocky Mountain Locust, IX, 73
Pezomachus minimus, II, 52, VIII, 54
Pezotettix, VIII, 115
Phacellura nitidalis, II, 7, 64, 68
Phalæna frugiperda, VIII, 48
 punctella, Supp., 58
 vernata, VII, 80
Phalænidæ, description of a new genus of, VIII, 12
phalanga, *Catocala*, III, 166
Phalangium dorsatum, IV, 17
Phaneroptera curvicauda, V, 124, VI, 164
 Phares, Dr. D. L., on Cotton Worm, II, 38, 40, VI, 24
 Phelps, Wilson, on Remedy for Chinch Bug, II, 29
Philampelus achemon, II, 74
 satellitæ, II, 76
philenor, *Papilio*, II, 116, III, 169, Supp., 54
 Philenor Swallow-tail, II, 116, III, 169
Philonthus apicalis, IV, 21
Phobetron hyalinum, V, 126
 pithicium, V, 126
phæniceus, *Agelaius*, VI, 27
phænicoptera, *Ædipoda*, VIII, 104
Pholus crantor, II, 74
 lycaon, II, 76
Phoxopteris comptana, Supp., 57
Phryganeidæ, V, 10
Phtheir, VI, 63
Phycis grossulariella, Supp., 57
Phycita juglandis, Supp., 80
 nebulo, III, 7, IV, 38, 41, Supp., 79, 80
 var. *nebulella*, IV, 42, Supp., 80
phycitæ, *Exorista*, IV, 40, Supp., 88
 Tachina, IV, 40, Supp., 88
Phygadeuon brevis, IV, 28
Phylloptera oblongifolia, II, 57, V, 123, VI, 169
Phyllotreta nemorum, III, 83
 striolata, III, 83
Phylloxera acanthohermes, VII, 119
 balbianii, VII, 91, 97, 99, 119
 caryæcaulis, VII, 97, 99, 117
 caryæfoliæ, IV, 66, V, 70, VI, 45, VII, 117
 caryæ-fallax, VII, 118
 caryæ-globuli, VII, 117
 caryæ-gummosa, VII, 118
 caryæ-ren, VII, 118
 caryæ-semen, VII, 117
 caryæ-septa, VII, 118
 castaneæ, VII, 118
 coccinea, VII, 119
 conica, VII, 118
 corticalis, VII, 119
 depressa, VII, 118
 forcata, VII, 118
 gallicola, VI, 30, 33, VII, 93
 lichtensteinii, VII, 119
 quercus, IV, 66, VI, 30, 41, 43, 68, VII, 91, 119, VIII, 158
 radicicola, VI, 33, 36, 37, VII, 93
 rileyi, IV, 66, VI, 42, 43, 64, 86, VII, 91, 117, 118, VIII, 158
 scutifera, VII, 119
 spinosa, VII, 118
 vastatrix, III, 85, IV, 55, V, 57, VI, 30, 63, 66, 86, 87, VII, 91, 117, VIII, 157
 vitifoliæ, II, 27, III, 84, IV, 55
 Position of the genus in the system, III, 96
 Synopsis of the American species of, VII, 117
 Phylloxera, American Oak, VI, 64, VII, 99, 118
 European Oak, VI, 46, 64
 Grape, III, 84, IV, 55, 67, V, 57, 63, VI, 41, 90, VIII, 157, IX, 43
 Phylloxera, Mite, VII, 106
phylloxeræ, *Thrips*, VI, 50
 Tyroglyphus, VI, 52, 53, 81

- Physonota quinquepunctata*, II, 59, Supp., 53
 unipunctata, Supp., 53
Phytocoris linearis, II, 113
Phytophaga, a division of Coleoptera, V, 11
 Terebrantine Hymenoptera, V, 10
Phytophagic varieties and species, I, 154, III, 127
Pickle Worm, II, 7, 64, 67, 70
picta, *Coccinella*, V, 101, Supp., 52
 Harmonia, Supp., 52
 Mamestra, II, 112
pictus, *Clytus*, III, 7, VI, 101
Pieris, III, 161
 brassicæ, III, 167
 oleracea, II, 105
 protodice, II, 104, IX, 57
 rapæ, II, 10, 107, III, 167, V, 24, 26, VII, 5
 vernalis, IX, 57
Piesma cinerea, II, 32, VII, 47
Pimpla, I, 178, III, 129, IV, 44
 annulipes, IV, 43, V, 49
 conquisitor, IV, 43
 indagatrix, IV, 43
 melanocephala, III, 129
Pine-leaf Scale, III, 92, V, 84, 97
 Confined to Pines proper, V, 100
 Its natural history, V, 98
 Natural enemies, V, 100
 Remedies, V, 101
 Two-brooded, V, 99
Pine Worm, Le Conte's, IX, 32
pinifoliæ, *Aspidiotus*, III, 92
 Mytilaspis, V, 97
pini, *Mytilaspis*, V, 98
Pioneer Press and Tribune, article from, on Rocky Mountain Locust, IX, 61
Piophila casei, II, 10
Pipiza radicum, I, 121, VI, 52
Pirene, I, 176
pisi, *Bruchus*, II, 11, III, 44, Supp., 53
pisorum, *Bruchus*, Supp., 53, 71
Pissodes strobi, III, 60
Placid Soldier-bug, IX, 17
Plant-lice, III, 87, IV, 35, VI, 33
 Number of annual broods of, II, 19, VII, 27
Platte City Landmark, article from, on Rocky Mountain Locust, VIII, 74
Platygaster, II, 103
platygaster, *Perilampus*, II, 87
Platyphyllum concavum, V, 123, 124, VI, 167
Platypsyllus castorinus, V, 16
Platyptilia, Supp., 84
 zetterstedtii, Supp., 84
Platysamia cecropia, IV, 103
Pleasant Hill Review, article from on Rocky Mountain Locust, VIII, 65
Plectrophanes lapponicus, IX, 91
plicata *Chlamys*, VI, 128
plorabunda, *Chrysopa*, II, 26, VI, 51, VII, 40
Plum Curculio, I, 50, II, 6, 48, III, 5, 6, 11, 13, 26, 30, 37, 38, 57, 58, 127, V, 26, 47, 106, 121, VI, 9, VII, 29, VIII, 36
 Artificial Remedies, I, 60
 Hooten's Curculio-catcher, III, 23
 Hull's Curculio-catcher, III, 19
 Jarring by machinery, III, 18
 Plum Curculio—Continued.
 Keeping it in check by the offer of premiums, III, 17
 Natural Remedies, I, 56
 Nocturnal rather than diurnal, III, 14
 Paris green as a remedy, III, 18
 Porizon Curculio Parasite, III, 28
 Remedies, I, 60, III, 41
 Sigalphus Curculio Parasite, III, 24
 Single-brooded and hibernates as a beetle, III, 11
 The Ransom Chip-trap process, III, 15
 Ward's Curculio-catcher, III, 20
Plum Gouger—Its character, distribution, II, 11, III, 39
 Habits and natural history, III, 40
 Its time of appearance, III, 40
 Often mistaken for the Plum Curculio, III, 40
 Remedies, III, 41
Plum Moth, III, 6, 25, 26, V, 51
Plum-weevil, III, 30, 31
Plume, Grape-vine, I, 137, IV, 129
Plusia brassicæ, II, 110, 111, Supp., 77, 78
 ni, II, 112, Supp., 78
 precactionis, II, 112
Plutella cruciferarum, II, 10, IV, 36
Podisus, Supp., 58
 placidus, IX, 17
Pœcilopectera compta, I, 152
 pruinosa, V, 122
Poisonous qualities of Colorado Potato-beetle, VIII, 10
polistiformis, *Egeria*, I, 127
Polistes rubiginosus, V, 54
Pollen carried in thunder-showers, V, 86
Pollyxerus cagurus, VII, 106
Polson, W. D., on Rocky Mountain Locust, IX, 74
Polygramma, VII, 18
 10-lineata, VII, 16
Polymorphism, VI, 43
polyphemus, *Attacus*, III, 170, IV, 13, 74, 85, 110, 121, 125, 138
 Telea, IV, 125
Polyphemus Moth, II, 19, VII, 27
 Issuing of, from cocoon IV, 127
Polyphemus Silkworm, IV, 125
 Food-plants, IV, 126
 Larval changes, IV, 126
 Natural enemies, IV, 126
 Natural history, IV, 125
 Parasites, IV, 129
 Value of silk, IV, 129
Polysphincta bicarinata, III, 71
pometaria, *Anisopteryx*, II, 97, VI, 29, VII, 80, 83, 86, VIII, 13, Supp., 56
pomicorticis, *Mytilaspis*, V, 73, 95, Supp., 85
pomifoliella, *Bucculatrix*, IV, 49, 51
pomivorella, *Micropteryx*, III, 7
pomonella, *Carpocapsa*, I, 62, 108, II, 10, III, 6, 101, IV, 27
 Trypeta, I, 108, III, 6, 91
pomorum, *Anthonomus*, III, 11
 Mytilaspis, V, 93, 94
Poplar Dagger, II, 119
 Spinner, II, 19, VII, 27

- Popular names, confusion from improper use of, VII, 187
- populi*, *Acronycta*, II, 119, 120, Supp., 72, 74
- populnea*, *Saperda*, IV, 22
- Porizon conotracheli*, III, 28, Supp., 64
- Porizon Curculio Parasite, III, 28
- Potash, Bichromate of, for Potato-bugs, IV, 14
- Potato-beetle (see Colorado Potato-beetle).
- Potato Bug (see Colorado Potato-beetle).
- Pest poison, VIII, 7, IX, 45
- Stalk-borer, I, 92
- Stalk-weevil, I, 93, III, 60
- worm, I, 95, V, 125
- Potherb Butterfly, II, 105
- Potts, R. B., on Rocky Mountain Locust, IX, 92
- Powers, Rev. Grant, on Northern Army-worm, II, 42
- Poweshiek*, *Oarisma*, VIII, 178
- Practical Entomologist*, article from, on Climbing Cut-worms, I, 71
- article from, on Harlequin Cabbage Bug, IV, 36
- Prairie Farmer*, article from, on Ailanthi-culture, IV, 114
- article from, on Canker Worm trap, VI, 25
- articles from, on Chinch Bug, II, 23, 30
- article from, on Climbing Cut-worms, I, 69
- article from, on Colorado Potato-beetle, I, 110
- article from, on Dark-sided Cut-worm, I, 75
- article from, on Food of Periodical Cicada, I, 29
- article from, on Hickory Bark-borer, V, 105
- article from, on Katydid, VI, 154
- article from, on Lesser Leaf-folder, IV, 48
- article from, on Locust Mite, VII, 177
- article from, on Remedy for Potato-beetle, IV, 15, VIII, 3
- article from, on Rocky Mountain Locust, VII, 135, 138
- Prairie Warbler destroying Canker Worm, VI, 27
- Pratt, S. M., on enemies of Rocky Mountain Locust, IX, 93
- Prayers to avert insect injury, VIII, 96
- precationis*, *Plusia*, II, 112
- Predictions verified, VII, 3, VIII, 58, 163, 184, IX, 57
- Preying Mantis, IX, 98
- Prionus imbricornis*, II, 89, III, 6, 75
- laticollis*, II, 87, III, 6, 75, V, 56
- obscurus*, I, 127
- Priority, law of, IV, 55, VII, 143, VIII, 179, IX, 7
- Pristiphora*, II, 8
- grossulariæ*, IX, 23, 26
- Pritchett, H. Carr, on Rocky Mountain Locust, IX, 75
- Processionary caterpillar, V, 126
- Proclamation of Governor Hardin relating to Locusts, VIII, 95
- Procris americana*, II, 85
- vitis*, II, 86
- Procris, the Grape-vine, V, 134
- Proctacanthus milbertii*, Supp., 88
- Proctotrupidæ*, V, 118
- Prodenia autumnalis*, III, 109, 116, 151, IV, 129, VIII, 48
- var. *fulvosa*, VIII, 49
- obscura*, VIII, 49
- commelinæ*, I, 88, III, 13, Supp., 56
- flavimedia*, Supp., 56
- lineatella*, Supp., 56
- Progress of Economic Entomology, V, 19
- Promachus*, Supp., 87
- apivora*, IX, 98
- bastardii*, II, 122, IV, 21, Supp., 60
- fitchii*, Supp., 60
- vertebrata*, II, 123
- promethea*, *Attacus*, IV, 74, 110, 121, 138
- Callosamia*, IV, 121, Supp., 55
- Promethea Silkworm, IV, 121
- Foodplants, IV, 123
- Larval changes, IV, 121
- Natural enemies, IV, 123
- Value of the cocoon, IV, 121
- Pronuba*, V, 150
- yuccasella*, V, 151, 160, VI, 131, VIII, 171, Supp., 58
- Chrysalis of, VI, 131
- Generic characters of, V, 150
- How the female fertilizes the plant, V, 154
- Larva of, V, 155
- Range of, V, 159
- proserpina*, *Apatura*, VI, 145
- Limnitis*, III, 171
- Protective imitation, III, 142
- protodice*, *Pieris*, II, 104
- Provancher, Abbé, on parasite of Cabbage Worm, II, 110
- on Polyphemus Silkworm, IV, 128
- pruinosa*, *Cicada*, I, 27
- Pruner, III, 6
- prunicida*, *Anthonomus*, III, 39, Supp., 54
- prunivora*, *Semasia*, I, 65, III, 6, 25
- Pseudohazis eglanterina*, V, 126
- Pseudo-Neuroptera, a division of Neuroptera, V, 14
- Pseudo-Tetramera, a section of Coleoptera, V, 10
- Pseudo-Trimera, a section of Coleoptera, V, 10
- Pseudopontia*, VIII, 170
- psi*, *Acronycta*, II, 121, Supp., 73
- Psoci* as museum pests, V, 41
- Psyche*, article from, on Rocky Mountain Locust, VIII, 109
- Psychomorpha epimenis*, III, 63, 64, VI, 87, 88, 90, 95
- Psylla pyri*, II, 10, 33
- Pterognostic variation, IX, 22
- Pterophorus*, II, 86
- cardui*, Supp., 83
- carduidactylus*, I, 180, III, 67
- periscelidactylus*, I, 137, III, 65, Supp., 58
- Ptinidæ*, IV, 53
- pubescens*, *Haltica*, I, 101
- pulchellus*, *Sphenophorus*, III, 60
- pulicaria*, *Corimelæna*, II, 33, VII, 48

- Pulicidae*, V, 15
pullatus, *Ichneumon*, III, 69
 Pullen, B., on Flea-like Negro-bug, II, 34
 on remedy for Peach Borer, I, 48, 34
punctata, *Hyphantria*, Supp., 55
 Pelidnota, III, 77, 78
punctella, *Phataena*, Supp., 58
punctipes, *Euschistus*, I, 113, IV, 19, 20, Supp., 58
 Pupation of Butterflies, III, 146, IV, 55, VI, 138,
 VIII, 179, Supp., 55
purgatus, *Ophion*, II, 53, VIII, 54
 Purged Ophion, II, 53, VIII, 54
 Purinton, J. A., on Rocky Mountain Locust, IX, 75
 Purple Emperor Butterfly, VI, 136
 Purple-finch unjustly accused of doing injury, I, 72
 destroying Canker Worms, VI, 27
 Purple Grackle destroying Canker Worm, VI, 28
purpureus, *Carpodacus*, VI, 27
pusilla, *Rhizopertha*, II, 14
pustulella, *Tinea*, Supp., 58
pustulosus, *Sphenophorus*, Supp., 54
 Putnam, J. D., on Rocky Mountain Locust, VII,
 141
Pyrameis atalanta, III, 167
 Pyramidal Grape-vine Worm, III, 72
pyramidea, *Amphipyra*, III, 73, 74
pyramidoides, *Amphipyra*, III, 72, 74, Supp., 75
Pyranga rubra, VI, 27
pyri, *Anthonomus*, III, 11
 Eriosoma, I, 118, III, 5, 95, 96, VI, 37, Supp., 59,
 87
 Psylla, II, 10, 33
Pyrrharctia, Supp., 55
- Q.**
- quadrigibbus*, *Anthonomus*, III, 29, 35
quadri-impressa, *Chrysobothris*, VII, 71
quadrispinosus, *Scolytus*, V, 105, 107, Supp., 54
 Quail destroying Chinch Bugs, VII, 41
 Locust eggs, IX, 91
 unjustly accused of being injurious, I, 72
Quedius molochinus, IV, 21
quercina, *Lachnosterna*, I, 67, 157, II, 19, VI, 123,
 VII, 27, Supp., 53
quercus-aciculata, *Cynips*, Supp., 59
 -frondosa (Gall), III, 25
 -inanis, *Cynips*, I, 14
 -spongifica, *Cynips*, I, 14
quercus, *Phylloxera*, IV, 66, VI, 30, 41, 43, 68, VII, 91,
 119, VIII, 158
 Quick, T. J., on Rocky Mountain Locust, IX, 71
 Quince Curculio, III, 35
 How it differs from the others, III, 35
 Its transformations and habits, III, 37
 Remedies, III, 38
quindecim-punctata, *Mysia*, IV, 18, Supp., 53
quinque-maculata, *Sphinx*, I, 95
quinquepunctata, *Physonota*, II, 59, Supp., 53
Quiscalus versicolor, VIII, 124
- R.**
- Radiata, a branch of the animal kingdom, V, 6
Radicicola or Root-inhabiting type of *Phylloxera*,
 VI, 36, 37, 38, 66, VII, 93
radicum, *Anthomyia*, IX, 92
 Pipiza, I, 121, VI, 52
 Rhodites, I, 13
rafflesia, *Euschemon*, VIII, 170
 Ragan, Z. F., on Rocky Mountain Locust, VIII,
 70, 105
 Ralls, Wm. C., on Rocky Mountain Locust, IX, 117
 Randolph, T. C., on Clover-hay Worm, VI, 103
 Ransom's Chip-trap for Curculio, III, 15
rapae, *Pieris*, II, 10, 107, III, 167, VII, 5
 Rapacious Soldier-bug, I, 114
 Rape Butterfly, II, 10, 107, V, 24, VII, 5
Raphigaster, IV, 20
 Raptatoria, a section of Orthoptera, V, 14
raptatorius, *Reduvius*, I, 114, Supp., 58
 Rascal Leaf-crumpler, IV, 38, 42, 44, VII, 81
 Natural enemies of, IV, 40
 Remedies for, IV, 40
 Raspberry Geometer, I, 139
 Root-borer, VI, 111
 Rathvon, S. S., on American Bean-weevil, III, 53
 on Colorado Potato-beetle, IX, 35
 on Periodical Cicada, I, 20, 22, IV, 31
 Ravenel, H. W., on Grape Phylloxera, VI, 83, VII,
 102, VIII, 164, 165
 Raymond, H. C., on Rocky Mountain Locust, IX,
 118
 Read, M. C., on Grape-vine Plume, III, 67
 Rearhorse, I, 169
 Reason vs. instinct, V, 83, 15.
 Reavis, D. B., on Rocky Mountain Locust, IX, 73
rectus, *Balaninus*, IV, 144
 Red Currant Borer, II, 10
 Red-eyed Vireo destroying Canker Worm, VI, 27
 Locusts, VIII, 124
 Woodpecker destroying Locusts, VIII,
 124
 -legged Ham-beetle, VI, 96
 Locust, VII, 125, 188, VIII, 150
 -shouldered Sinoxylon, II, 53, V, 54
 -tailed Tachina-fly, II, 50, III, 129, VI, 96, VII,
 179, VIII, 53
 -start destroying Canker Worm, VI, 27
Reduvius, II, 32
 raptatorius, I, 114, Supp., 58
 Red Weevil, II, 16, IX, 17
 Red-winged Blackbird destroying Canker Worm,
 VI, 27
 Chinch Bug,
 VII, 41
 Reed, E. B., experiments with various substances
 for Potato-bugs, IV, 14
regalis, *Citheronia*, III, 151, IV, 129, V, 141
 Relation of Insects to Agriculture, V, 5, 18
 Relaxing Insects, V, 41
 Remedies, V, 25
 for Abbot's White Pine Worm, IX, 32
 Ailanthus Worm, I, 152
 American Meromyza, I, 161
 Apple Curculio, III, 34
 -leaf Bucculatrix, IV, 50
 Skeletonizer, IV, 45
 -root Plant-louse, I, 123
 -tree Tent-caterpillar, III, 120
 -twig Borer, IV, 53
 Army Worm, II, 53, VIII, 54
 Bag-worm, I, 151
 Bee-killer, I, 168
 Moth, I, 167

Remedies for Blister-beetles, I, 99

Blue Caterpillars of the Vine, II, 84

Boll Worm, III, 108

Cabbage Plusia, II, 111

Worms, II, 109

Canker Worms, II, 98, VI, 24, VII, 85, VIII, 19

Chinch Bug, II, 28, VII, 31

Clover-hay Worm, VI, 105

Codling Moth, I, 65, IV, 23, V, 46, VI, 9

Colorado Potato-beetle, I, 116, III, 99, IV, 11, V, 53, VI, 13, VII, 8, VIII, 3, IX, 45

Corn Worm, III, 108

Cotton Worm, II, 41, VI, 17

Cut-worms, I, 90

Fall Army Worm, III, 114

Web-worm, III, 132

Flat-headed Borer, I, 47, VII, 76

Flea-like Negro-bug, II, 35

Gooseberry Fruit-worm, I, 141

Span-worm, IX, 6

Grain Bruchus, III, 51

Grape-cane Gall-curculio, I, 132

-leaf Folder, III, 62

Phylloxera, III, 89, IV, 68, V

71, VI, 55, VII, 105

-root Borer, III, 77

-vine Colaspis, III, 84

Fidia, I, 133

Flea-beetle, III, 80

Fruit-worm, I, 135

Plume, I, 138, III, 68

Green-striped Maple Worm, V, 141

Harlequin Cabbage-bug, IV, 38

Hickory Bark-borer, V, 107

Imported Currant Worm, IX, 13

Jumping Sumach Beetle, VI, 121

Legged Maple Borer, VI, 109

Lesser Apple Leaf-folder, IV, 49

Native Currant Worm, IX, 26

New Grape-root Borer, I, 128, II, 88

Oyster-shell Bark-louse, I, 16, V, 90

Peach Borer, I, 48

Pea-weevil, III, 48

Periodical Cicada, I, 30

Pickle Worm, II, 70

Pine-leaf Scale-insect, V, 101

Plum Curculio, I, 60, III, 15, V, 25

Plum Gouger, III, 41

Potato Stalk-borer, I, 92

-weevil, I, 95

Worm, I, 96

Pyramidal Grape-vine Worm, III, 73

Quince Curculio, III, 38

Rascal Leaf-crumpler, IV, 40

Red-legged Ham-beetle, VI, 100

-shouldered Sinoxylon, IV, 54

Rocky Mountain Locust, VII, 181,

VIII, 125, IX, 99, 108

Rose Chafer, V, 110

Round-headed Apple-tree Borer, I, 45

Seed-corn Maggot, I, 155

Sheep Bot-fly, I, 165

Smeared Dagger, III, 70

Strawberry Crown-borer, III, 43

Remedies for Strawberry Leaf-roller, I, 143

Worm, IX, 28

Striped Cucumber-beetle, II, 66

Tarnished Plant-bug, II, 115

Tent-caterpillar of the Forest, III, 128

Tile-horned Prionus, II, 90

Tobacco Worm, V, 56

Tortoise-beetles, II, 60

Tree-cricket, I, 139

Wheat-head Army Worm, IX, 54

White Grub, I, 157

White-marked Tussock Moth, I, 147

Zebra Caterpillar, II, 113

Remington, M. C., on Clover Hay Worm, VI, 104

renigera, *Celaena*, I, 86, Supp., 56*repanda*, *Cicindela*, VIII, 52*repentis*, *Agrotis*, Supp., 77

Report of Committee on Entomology, read before the Mo. State Horticultural Society, II, 5

Retarded development, V, 130, 132

retinervis, *Microcentrus*, VI, 155*Rhipheus*, VIII, 170*Rhipiphorus paradoxus*, VI, 125*pectinatus* var. *ventralis*, VI, 125*Rhizophis vastatrix*, VI, 31*Rhizopertha pusilla*, II, 14*Rhodites ignota*, I, 13*radicum*, I, 13*Rhodobæus*, Supp., 54*rhois*, *Blepharida*, I, 100, II, 58, VI, 118

Rhopalocera a section of Lepidoptera, V, 12

Rhynchites bacchus, III, 11*betuleti*, III, 11*conicus*, III, 11*Rhyparochromus devastator*, VII, 22*Rhyssa*, VIII, 38*ribesii*, *Aphis*, VI, 46*ribis*, *Aphis*, II, 10

Rice Bunting destroying Army Worm, VIII, 52

Richmond *Conservator*, article from, on Rocky Mountain Locust, VIII, 75Richmond *Whig*, article from, on Apple grape-vine gall, V, 115*ricini*, *Samia*, IV, 112

Riehl, Wm., on Army Worm, VIII, 39

rileyana, *Tortrix*, I, 153, Supp., 81*rileyi*, *Aleiodes*, III, 71*Eriosoma*, Supp., 87*Phylloxera*, IV, 66, VI, 42, 43, 64, 86, VII, 91, 117, 118, VIII, 158

Ring-banded Soldier-bug, IV, 18

Ring-legged Pimpla, V, 49

Roberts, A., on Rocky Mountain Locust, IX, 117

Robin destroying Canker Worm, VI, 27

robinæ, *Spermophagus*, III, 45*roborana*, *Spilonota*, Supp., 57

Robords, Chas. J., on Rocky Mountain Locust, IX, 69

Robson, J. W., on birds destroying Canker Worms, VI, 27

on Rocky Mountain Locust, IX, 66, 91

on Wheat-head Army Worm, IX, 51

Rocky Mountain Locust, VII, 121, VIII, 22, 57, IX,

Rocky Mountain Locust—Continued.

- Account of Damage done in Missouri, VIII, 89
- Additional Natural Enemies, IX, 91
- Animals which destroy the Eggs, IX, 91
- Area in which Eggs were laid in 1876, IX, 116
- Artificial Means of Destroying the Eggs, VIII, 125
- Bill to provide for investigation of, VIII, 133
- Bounties for catching and destroying Locusts, VIII, 138
- Changes that followed the Locusts, VIII, 121
- Chronological history, VII, 132
- Conditions of Migration, VII, 112
 - which prevent the permanent Settlement of the Species in Missouri, VIII, 113
- Contrast between Spring and Fall, during locust injury, VIII, 119
- Definition of the Species, VIII, 114
- Departing swarms do not return, VIII, 124
- Descriptive, VII, 126
- Destination of departing Swarms, VIII, 106, IX, 77
- Destitution in Missouri in 1875, VIII, 91
- Destruction of the unfledged young, VIII, 126, IX, 108
- Does the Female lay more than one egg-mass? IX, 85
- Direction of flight, IX, 81
- Direction in which young Locusts travel, VIII, 101
 - taken by winged Locusts, VIII, 105, IX, 81
- Easily confounded with the Red-Legged Locust, VII, 125
- Eastern line reached in 1876, IX, 80
- Egg-mass, philosophy of, IX, 87
- Eggs, condition of, in winter, IX, 116
 - description of, IX, 87
 - how laid, IX, 86
 - where laid by preference, VII, 123
 - experiments with, IX, 99
- Enemies and parasites, VII, 174, VIII, 124, IX, 91
- Exodus of the swarms in 1875, VIII, 104
- Experience in the Spring of 1875, VIII, 118
- Experiments with the Eggs, and conclusions drawn therefrom, IX, 99, 106
- Food-plants, VII, 158, VIII, 121
- General outlook in the Spring of 1875, VIII, 60
- Governor's Proclamation, VIII, 95
- Green variety of, VIII, 117.
- Habits of the unfledged Locusts, VIII, 100
- Hatching of Locusts, IX, 89
- How the young Locust escapes from the Egg, IX, 88
- How to avert Locust Injuries, VIII, 131
- Influence of burying the eggs at different depths, IX, 104
 - exposure to air on the eggs, IX, 104
 - freezing and thawing on the eggs, IX, 99
 - moisture on the eggs, IX, 101
 - wind in determining the course of Locust swarms, IX, 81

Rocky Mountain Locust—Continued.

- Injury from other, non-migratory Locusts, VII, 171
 - to fruit and fruit trees, VIII, 121
- Invasion of 1873, VII, 141
 - 1874, VII, 143
 - 1876, IX, 59
- Legislation, both national and local, VIII, 132
- Lessons of the year 1875, VIII, 142
- Locusts as food for Man, VIII, 143
- Measurements of *Caloptenus spretus*, VII, 130
- Migratory instinct and great destructive Power belong to but one species west of the Mississippi, VII, 124
- Native home, VII, 161, VIII, 109
- Natural enemies, VII, 174, VIII, 124, IX, 91
- Natural history, VII, 121, VIII, 97
- Not a divine Visitation, VIII, 97
- Not led by "Kings" and "Queens," VIII, 103
- Omaha Conference, IX, 106
- Outlook in Missouri in 1875, VIII, 61
- Predictions for 1875, VII, 166
- Previous experience in the Spring of 1867, VIII, 57
- Prospects in 1877, IX, 121
- Rate at which the young travel, VIII, 102
- Rate at which the insects spread, IX, 80
- Ravages, VII, 156
 - of migratory Locusts in the Atlantic States, VII, 167
- Reports of Correspondents, IX, 69, 117
- Source of Locust swarms of 1876, IX, 79
- Suggestions, VIII, 140
- Time of appearance, VII, 160
- Time of leaving of the winged insects, VIII, 104, 125
- Unnecessary alarm caused by native Locusts, VIII, 148
- Wind, influence of, on flight, IX, 87
- Roe, J. E., on Rocky Mountain Locust, IX, 119
- Rogers, Dan F., on Chinch Bug, II, 23
- Rogers, J. R., on Apple-tree Bark-louse, V, 78
- Root-borer of the Grape-vine, I, 124, II, 87, III, 75
 - Raspberry, VI, 111
 - Squash, II, 64
- Root-borers, III, 6
- Root-louse of the Grape-vine. (See Grape Phylloxera.)
- Root Plant-louse of the apple tree, I, 118, III, 5, IV, 68, 69
 - Syrphus fly, I, 121
- Rope and tin band for Canker Worm, VI, 26, 27
- rosaceana*, *Loxotenia*, III, 6
- rosæ*, *Selandria*, II, 19, VII, 27
- rosea*, *Uroplata*, III, 6
- Rose-breasted Grosbeck destroying Potato-beetle, V, 54
- Rose-bug, III, 6
- Rose-bush Saw-fly, IX, 19
- Rose Chafer, V, 108
 - Hispa, III, 6
- Rose Leaf-roller, III, 6
- Rosy Dryocampa, V, 139
- Round-headed Apple-tree Borer, I, 45, II, 19, IV, 124, VII, 27

Round-headed Apple-tree Borer—Continued.

- Food plants, I, 43
 Natural history, I, 43
 Remedies, I, 45
 Rove-beetle, larva of, IV, 21
 Rove-beetles, habits of some, VI, 162
 Royall's Paris Green mixture, VI, 21
rubi, *Egeria*, VI, 113, Supp., 72
 Selandria, I, 52
rubicunda, *Anisota*, V, 140
 Dryocampa, III, 123
rubivora, *Aplodes*, I, 139, Supp., 79
rubivoraria, *Synchlora*, Supp., 79
rubra, *Pyranga*, VI, 27
ruficapilla, *Helminthophaga*, VI, 27
ruficollis, *Necrobia*, VI, 101
rufimanus, *Bruchus*, III, 56, Supp., 70
rufipedis, *Micropus*, VII, 22
rufipes, *Corynetes*, VI, 101
 Pentatoma, IV, 20
rufus, *Sigalphus*, III, 27, Supp., 68
 Rummaging Ground-beetle, II, 103, III, 129
Ruptor ovi, structure in many insect embryos, for
 bursting the egg-shell, IX, 127
Rural Carolinian, article from, on Locusts, VII,
 173
Rural New Yorker, article from, on Apple-leaf Buc-
 culatrix, IV, 50
 article from, on trapping Cur-
 culio, III, 16
Rural World, Colman's, article from, on Apple-tree
 Bark-louse, V, 77
Rural World, article from, on Fall Army Worm,
 III, 109
 article from, on Grape Phylloxera,
 IV, 55, VI, 84
ruscarius, *Elaphrus*, VIII, 52
 Rust-red Social Wasp, V, 54
 Rutbottom, W. F., on Rocky Mountain Locust,
 IX, 117
ruticilla, *Setophaga*, VI, 27

S.

- Sacramento Union*, article from, on Silk Industry,
 IV, 80
salicis-strobiloides, Gall, VI, 155
saligneana, *Euryptychia*, II, 134, Supp., 57
saltator, *Orocharis*, I, 138, V, 119, Supp., 60, 62
Saltatoria, a section of Orthoptera, V, 14
Samia cecropia, IV, 103, Supp., 55
 columbia, IV, 107, 111, 128, Supp., 55
 cynthia, IV, 112
 guerinii, IV, 112
 ricini, IV, 112
samiæ, *Cryptus*, IV, 110, 111, Supp., 52
 Sanborn, F. G., on frame for insect net, V, 31
sanguinea, *Cycloneda*, Supp., 52
Saperda bivittata, I, 42, II, 19, III, 6, VII, 27, Supp.,
 53
 candida, Supp., 53
 discoidea, V, 106
 populnea, IV, 22
Sarcophaga, II, 110
 caenaria, VII, 180, IX, 95, Supp., 60
 var. *mantivora*, VII, 180
 sarraceniae, VII, 180, 181, IX, 95, Supp.,
 60
Sarracenia Flesh-fly, VII, 181
sarraceniae, *Sarcophaga*, VII, 180, 181, IX, 95, Supp.,
 60
 Satellite Sphinx, II, 76
satellitica, *Philampelus*, II, 76
Saturnia maia, IV, 41, V, 127
 io, V, 133
saucia, *Agrotis*, I, 74, Supp., 55
 Saunders, Wm., Experiments with poisons for Po-
 tato-beetle, IV, 14
 on Grape-seed Maggot, II, 93
 on Imported Currant worm, IX,
 12, 14
 on Pea-weevil, III, 49
 Saunders, W., on *Oidium Tuckeri* in America, V,
 69, 70
 Saw-flies, II, 8, V, 9, 10, VIII, 38
 Sawyer's Canker Worm trap, VI, 26
scabiei, *Acarus*, VI, 61
scabrosellus, *Nemotois*, V, 160
scabrum, *Trombidium*, VII, 175
 Scale of Bark-louse, True Nature of, V, 80
scandens, *Agrotis*, I, 76, 78, III, 6, Supp., 55
 Scarlet Mite, VII, 175
 Tanager destroying Canker Worm, VI, 27
Scenopinus, V, 8
Schizoneura, Supp., 87
 lanigera, Supp., 59
Schæmherri, *Pachyrhynchus*, III, 57
Sciara, VIII, 23, 24, Supp., 59
Scientific American, article from, on Colorado Po-
 tato-beetle, IX, 35
Scolia, VII, 174
 bicincta, VI, 124
 flavifrons, VI, 124
Scolytidae, III, 6
scolytivorus, *Bracon*, V, 106, Supp., 67
Scolytus, III, 6, V, 106
 caryæ, V, 103, Supp., 54
 destructor, V, 104
 muticus, V, 105, 107
 4-spinosus, V, 105, 107, Supp., 54
scribonia, *Ecpantheria*, IV, 141, 143
scrutator, *Calosoma*, II, 103, III, 129, VIII, 52
scudderiana, *Hedya*, Supp., 57
 Scudder, S. H., on Protective resemblance in But-
 terflies, III, 166
 on Rocky Mountain Locust, VIII,
 109
 on Southern Cabbage Butterfly, II,
 104
sculptilis *Sphenophorus*, Supp., 54
scutellaris, *Anthonomus*, Supp., 54
Scutellera, a family of Heteroptera, II, 32, 33, IV,
 19, VII, 48
Scymnus, II, 25, 27, VI, 51, VII, 39
 cervicalis, I, 122, V, 100
 consobrinus, V, 100
 terminatus, V, 100
 Seabrook, W. B., on Cotton Moth, II, 40
Sedalia Press, article from, on White Grub, I, 158
 Seed corn maggot, I, 154
 Curculio of the Grape, I, 129
Seiurus aurocapillus, VI, 27
Selandria, V, 26
 cerasi, II, 18, VII, 27

- Selandria rosæ*, II, 19, VII, 27, IX, 19
rubi, I, 52
selene, *Attacus*, IV, 125
Semasia prunivora, I, 65, III, 6, 25, V, 51
Semiotellus clisiocampæ, III, 120
semisculpta, *Chrysobothris*, VII, 71
senatoria, *Dryocampa*, III, 123, IV, 41
Senometopia bicincta, V, 140
militaris, II, 50
septemdecim, *Cicada*, I, 18, 19, 20, II, 19, III, 6, VII
27, Supp., 58, 59
septentrionis, *Brenthus*, VI, 116, 117
Sericaria mori, IV, 75
sericeum, *Trombidium*, VII, 175, Supp., 63
sericeus, *Asilus*, II, 123
Spermophagus robiniae, III, 45
serratus, *Bruchus*, III, 56, Supp., 70
sesostris, *Ampelogypter*, Supp., 71
Baridius, III, 60, Supp., 71
Setophaga ruticilla, VI, 27
Seventeen-year Locust, II, 19, III, 6, VII, 27
Seventeen and thirteen year broods of the Period-
ical Cicada, I, 18
Sex, law of, V, 85
Sex not affected by food, VIII, 19
Sexed Phylloxera, VIII, 158
Shad-fly, V, 143
Shane, J. B., on Rocky Mountain Locust, IX, 118
Shattuck, J. C., on Rocky Mountain Locust, VII,
178
Shaw, G. W., on Soldier Bugs, V, 51
Sheep Bot-fly, I, 161
-ticks, V, 13
Shepherd, S., on Hickory Bark-borer, V, 105
Shimer, Dr. H., on Chinch Bug, II, 20, 24, 26, 30,
VII, 39, 40
Short-winged Ichneumon, IX, 55
Shulz, G. E., on Rocky Mountain Locust, IX, 70
sibylla, *Limenitis*, III, 171
Sigalphus Curculio Parasite, III, 25
Sigalphus curculionis, III, 25, 27, Supp., 67
var. *rufus*, III, 27, Supp., 63
signatipes, *Ichneumon*, III, 69
signifer, *Cassida*, II, 63
Silk-growing in California, IV, 79
Kansas, IV, 82
Missouri, IV, 83
Silkworm, The *Ailanthus*, IV, 112
American, IV, 104
Cecropia, IV, 103
Luna, IV, 123
Mulberry, IV, 75
Perny, IV, 137
Polyphemus, IV, 125
Promethea, IV, 121
Tusseh, IV, 138
Yama-mai, IV, 130
Silkworms, IV, 72
Silky *Asilus*, II, 123
Silky Mite, VII, 175, IX, 91
Silpha americana, VI, 100
Simmons & Tillson, on Grape root Borer, I, 125
Simpson, T. W., on Rocky Mountain Locust, IX, 75
Sinea diadema, Supp., 58
Sinoxylon basillare, IV, 53, 54, V, 54
Sinoxylon, the Red-shouldered, IV, 52, 53, 54, V, 54
siro, *Tyroglyphus*, VI, 52
Sitophilus granarius, II, 10, III, 60
Skimmed Milk as remedy for Gooseberry Span-
worm, IX, 6
Skunk destroying Locust eggs, IX, 91
Slug Worm of the Pear, II, 19
Rose, II, 19, VII, 27
Slug-worms, V, 26
Small White Bristly Cut-worm, I, 86
Smeared Dagger, III, 70
Smiley, W., on Rocky Mountain Locust, IX, 70
Smith, H. J., on Green-striped Maple-worm, V, 137
Smith, Jos., on Snake Worm, VIII, 24
Smith, J. F., on Rocky Mountain Locust, IX, 68
Smith, J. H., on Rocky Mountain Locust, VIII, 62
Smith, S. I., on oviposition of *Conocephalus*, VI,
155
Smith, S. S., on Army Worm, VIII, 39
Smith, T., on seeding of Yuccas, V, 159
Smith, W. A., on Rocky Mountain Locust, IX, 70
Smith, W. R., on Rocky Mountain Locust, VIII, 85
smithii, *Cryptus*, IV, 111
Snake-worms, VIII, 23, 24
Snapping-beetles, III, 6
Snidow, W. L., on Rocky Mountain Locust, IX, 71
Snout-beetle, II, 92, III, 5, 10, 37, VI, 116
The Imbricated, III, 57
Snow, F. H., on False Chinch Bug, V, 111
on Rocky Mountain Locust, VIII, 77,
114, IX, 93
Snowy Tree-cricket, V, 120
Snyder, C., on Rocky Mountain Locust, VII, 194
socialis, *Spizella*, VI, 27
Soda as remedy for Apple-tree Bark-lice, I, 17
Soldier-bug, The Glassy-winged, III, 137
The Spined, I, 77, 89, 113, II, 32, 34, IV,
19, V, 51, 133, IX, 17
Soldier-bugs as enemies of Cicada, I, 26
Codling Moth, V, 51
solidaginis, *Trypeta*, I, 13, 173
Solidago Gall-maker, II, 134
Gall-moth, I, 173, II, 20, 132
Song-sparrow destroying Canker Worm, VI, 27
soror, *Chrysobothris*, VII, 71
Sorsby, B. A., on attracting Boll-worm Moth by
sweets, III, 108
Sounds from insects sometimes inaudible, VI, 152
Southern Cabbage Butterfly, II, 104
Cotton Army-worm, II, 49, VIII, 34
Grass-worm, II, 41
Southern Farmer, article from, on Paris Green for
Cotton Worm, VI, 19
Span-worms, II, 110
Spanish blister-beetle, V, 18
Spathius trifasciatus, V, 106, Supp., 67
Species, definition of, VI, 143, VII, 115, 179
geographical range of, IX, 82
speciosus, *Stizus*, I, 27, Supp., 52
Speckled cut-worm, I, 84
Spectrum femoratum, VI, 156, VII, 181
Spermophagus robiniae, III, 45
Sphecius, Supp., 52
Sphenophorus pulchellus, III, 60
pustulosus, Supp., 54
sculptilis, Supp., 54
13-punctatus, III, 60, Supp., 54

- Sphenophorus truncatus*, III, 59
zeæ, III, 59, Supp., 54
- Sphinges, V, 12
- Sphingidæ*, III, 123, IV, 86
- Sphinx*, II, 71, 74, 76
carolina, I, 96, IV, 129
crantor, II, 74
lycaon, II, 76
myron, II, 71
5-maculata, I, 95, V, 125
- Sphinx moth, II, 78, V, 56, VI, 162
- Spiderwort Owlet moth, III, 113
- Spilonota roborana*, Supp., 57
- Spilosoma virginica*, III, 68
- Spined Soldier-bug, I, 77, 89, 113, II, 32, 34, IV, 19, V, 51, 133, IX, 17
- spinosa*, *Arma*, I, 77, 89, 113, II, 32, IV, 19, Supp., 58
Phylloxera, VII, 118
- spiralis*, *Trichina*, IV, 70
- Spizella socialis*, VI, 27
- Spotted Ladybird, I, 112, II, 25, 27, 36, V, 149, VII, 39
Pelidnota, III, 77
- Spray Machine, Peck's, VIII, 4
- spretis*, *Acridium*, VII, 128
- spretum*, *Acridium*, VII, 128
- spretus*, *Caloptenus*, VII, 121, 128, VIII, 57, 109, 114, Supp., 89, 90
- Spring in Europe and America, VI, 151
- Spring Canker-worm, VII, 80, VIII, 17, 18
Canker Worm Moth, VIII, 37
- Sprinkler for the use of Paris Green Water, VI, 20, VII, 15, VIII, 5
- Squash bug, I, 113, II, 31, VII, 46
Borer, II, 64
- St. Joseph Herald*, article from, on Colorado Potato-beetle, III, 97
article from, on Rocky Mountain Locust, VIII, 67, 69, 73, 75
- St. Louis Globe-Democrat*, articles from, on Rocky Mountain Locust, VIII, 63, 71, 92, 155
Republican, articles from, on Rocky Mountain Locust, VIII, 69, 73, 75, 148, IX, 73
- Stag-beetle, V, 145
- Stainton, H. T., on *Pronuba yaccasella*, V, 160
- Stalk-borer, III, 105, VIII, 37
of the Potato, I, 92
weevil of the Potato, I, 93
- Staphylinidæ*, VI, 162, VIII, 20, 24
- State University, cabinet for, VII, (preface, p. 5)
- Steel-blue Flea-beetle, I, 101, III, 79
- Stelle, J. P., on Periodical Cicada, IV, 32
- Stenocorus villosus*, I, 25
- Stenopogon consanguineus*, IX, 98
- Stenoptycha*, Supp., 57
- Stevenson, Hugh, on Rocky Mountain Locust, IX, 70
- stigma*, *Dryocampa*, III, 123, IV, 41, V, 141
- Sting of the Periodical Cicada, I, 26
- Stinging larvæ, V, 125
- Stiretrus fimbriatus*, I, 114, II, 34, IV, 20
- stirpicola*, *Crabro*, IX, 95, Supp., 89
- Stizus grandis*, I, 27, Supp., 52
speciosus, I, 27, Supp., 52
- Stomoxys*, V, 13
- Stone, W. B., on False Chinch Bug, V, 111
- Strachia histrionica*, IV, 35
ornata, IV, 38
- Strawberry Crown-borer, III, 42
Leaf-roller, I, 142
Worm, IX, 27
Descriptive, IX, 28
Remedies, IX, 28
- Strepsiptera*, V, 15
- striata*, *Dendroica*, VI, 27
- Stridulation of Acrididæ, VI, 153
Burying beetles, III, 14
Gryllidæ, VI, 154
Horned Passalus, IV, 139
Katydidæ, III, 154
Locusts, III, 153
Cecanthus latipennis, Supp., 60
Orocharis saltator, Supp., 62
Plum Curculio, III, 14
Three-lined Leaf-beetle, III, 14
- Stringer, J. E., on Rocky Mountain Locust, IX, 71
- striolata*, *Haltica*, III, 44
Phyllotreta, III, 83
- Striped Blister-beetle, I, 96, 115
Bug, II, 64, 66
- Striped Chrysops, II, 129
Cucumber-beetle, II, 64, 65, III, 6
Flea-beetle, III, 44
Squirrel destroying Locust eggs, IX, 91
- strobi*, *Pissodes*, III, 60
- Strobiloides* Gall on Willow, VI, 155
- Strong, W. C., on grafting grape-vine, VII, 114
- Strongylium tenuicolle*, VI, 118
- Stroop, L. J., on Archippus Butterfly, III, 151
- Structure, adaptation of, to habit, VI, 154
- Struggle for existence, VIII, 122
- Stylopidæ*, V, 15
- Stylops*, V, 15, VI, 125
- suavella*, *Myelois*, IV, 41, Supp., 79
- Subangular Ground-beetle, I, 58
- subangulata*, *Aspidoglossa*, I, 58
- subcadens*, *Celæna*, Supp., 56
- subcostalis*, *Xylina*, III, 136, Supp., 75
- subcyaneus*, *Ichneumon*, III, 69
- subgothica*, *Agrotis*, I, 81, 83, III, 151, Supp., 55, 56
- Subimage, Chrysopa issues from cocoon as, I, 57, II, 26
- Subjoined Hadenæ, I, 84
- subjuncta*, *Hadenæ*, I, 84
- Submersion as remedy for Chinch Bug, VII, 31
Phylloxera, IV, 69, VI, 55
Rocky Mountain Locust, VII, 182
- suffusa*, *Agrotis*, Supp., 55
- suilla*, *Colaspis*, III, 82
- Suits due to insects, VI, 96
- sulcatus*, *Otiorynchus*, III, 11
- Sulphide of Potassium as remedy for Gooseberry Span-worm, IX, 7
- Sulphur as remedy for *Phylloxera*, IV, 69
- Sulphuretted hydrogen as remedy for *Phylloxera*, IV, 69
- Sumach-beetle, The Jumping, VI, 118
- Summer Yellow Bird destroying Canker Worm, VI, 28
- surinamensis*, *Sylvanus*, III, 6

- suturalis*, *Anthonomus*, III, 60
 Swarming of butterflies, III, 151
 Sweetened water for Codling Moth, IV, 27
Sylvanus surinamensis, III, 6
sylvatica, *Olisiocampa*, II, 7, 37, III, 121, IV, 41, Supp., 55
symphoricarpi, *Tortrix*, I, 154, Supp., 82
Synchlora, Supp., 79
 albolineata, Supp., 79
 rubivorana, Supp., 79
Synemon thesesa, VIII, 178
 Synopsis of the American species of the genus
 Phylloxera, VII, 117
Syrphidæ, V, 13
Syrphus-fly, VI, 51
 of Root-louse, I, 121
 Larva, V, 149, VI, 51
Systæchus, Supp., 60
- T.**
- Tabanidæ*, VI, 123
Tabanus atratus, II, 128, 129, 130
 bovinus, II, 129
 cinctus, II, 128
 costalis, II, 128
 lineola, II, 128
tabescens, *Erax*, II, 124
tabida, *Chrysopa*, VII, 106
Tachina, II, 103, VIII, 107
 anonyma, IV, 129, V, 133, 139, VII, 178, VIII, 179, IX, 54
 archippivora, III, 116, 150, V, 149
 auricincta, V, 140
 bicincta, V, 140
 bifasciata, V, 140
 phycitæ, IV, 40, Supp., 88
Tachina-fly, II, 109, 110, 120, III, 62, 129, 142, 149, 157, 161, IV, 123, V, 133
 The Anonymous, VII, 178
 of Army-worm, IV, 109, VIII, 50
 of Cecropia Worm, IV, 108
 The Red-tailed, VI, 96, VII, 179
Tachinidæ, V, 13
 Talbot, R. H., on Rocky Mountain Locust, IX, 68
tapetzella, *Tinea*, II, 10
Tarantula, II, 106
tarda, *Tiphia*, VI, 126
 Tarnished Plant Bug, II, 113, 114, IV, 20
 Tawny Emperor, VI, 140
 Earlier states, VI, 141, 148
 Food-plants, VI, 141
 Parasite, VI, 142
 Taylor, A. S., Locust History in America, VII, 133
Telea polyphemus, IV, 125
Telephorus bilineatus, IV, 29, 30
telifera, *Agrotis*, I, 80, Supp., 55
 Ten-lined Spearman, I, 103
Tenebrio molitor, II, 9, 10, VI, 118, IX, 43
 obscurus, II, 9, 11, IX, 43
 Tenebrionid Larva, VI, 118
 Tent Caterpillar, II, 100, III, 130, 132
 Tent Caterpillar of the Apple, II, 7, III, 117
 Development, III, 119
 Eggs, III, 118
 Food-plants, III, 120
 Remedies, III, 120
 Tent Caterpillar of the Forest, II, 7, 37, III, 120, 121, 134, V, 128, VIII, 23
 Is it ever destructive? III, 127
 Larval habits of, III, 124
 Natural history of, III, 121
 Remedies for, III, 128
 Summary, III, 129
Tenthredinidæ, V, 10, VI, 70
tenuicollis, *Strongylium*, VI, 118
Teras oxycoccana, Supp., 83
Terebrantia, a section of Hymenoptera, V, 9
Termes flavipes, IX, 43
 frontalis, II, 11
tesselata, *Halesidota*, III, 127
Tetracha virginica, I, 115
Tettigonia, I, 171, III, 38
Tettix granulata, VIII, 150
texana, *Cassida*, Supp., 54
 Texas, Locust History in, VIII, 88, IX, 76
 Text Books on Entomology, V, 44
textor, *Hyphantria*, III, 130, 132, Supp., 55
Thecla, VI, 140, VIII, 177
thelxiope, *Heliconius*, III, 173
Theognis albicinctus, V, 154
 phyllopus, V, 154
thesesa, *Synemon*, VIII, 178
Thersilochus, III, 28, Supp., 65
 Thick-legged Buprestian, VII, 72
thuiella, *Bucculatrix*, IV, 51
 Thirteen-spotted Lady-bird, I, 112
 Thirteen-year Locust, II, 19, VII, 27
 Thistle Plume, I, 180, II, 112
 Thomas, Prof. Cyrus, Controversy on habits of
 Army Worm, II, 47
 Description of Red-legged
 Locust, VII, 126
 on Army Worm, VIII, 43, 45
 on Colorado Potato-beetle, VIII, 8
 on Rocky Mountain Locust, VII, 141, VIII, 115, 153
 Thompson, E. A., on remedy for Peach Borer, I, 49
 Thompson, Wm., on Rocky Mountain Locust, IX, 118
 Thornburg, J. M., on Rocky Mountain Locust, IX, 72
 Thornton, Dr. C. W., on Army Worm, III, 111
 Thrasher, J., on Rocky Mountain Locust, IX, 74
 Three-banded Spathius, V, 106
 Three-lined Leaf-beetle, I, 99, II, 58, III, 14
Threnodes, VIII, 170
Thripidæ, V, 16
Thrips, II, 6, III, 29, V, 16, 118, VI, 50
 phylloxera, VI, 50
Thymele, VIII, 175
thymi, *Nysius*, V, 113, Supp., 85
Thyreus abbotii, II, 78
thyridopterigis, *Hemiteles*, I, 150, Supp., 65
Thyridopteryx ephemeraeformis, I, 147
Thysanoptera, V, 16
tibiale, *Trochilium*, VI, 113, Supp., 72
 Tiger-beetles preying on Locusts, IV, 98
 Tiger-moth, IV, 88
 The Isabella, IV, 143

- Tilden, Josiah, on Rocky Mountain Locust, IX, 72
 Tile-horned Prionus, II, 89, 90
 Dimorphous male form, II, 90
 Food-plants, II, 89, 90
 Remedy, II, 91
 Titts, R. H., on parasite of Flat-headed Borer, VII, 74
tinctorium, *Trombidium*, VII, 175
 Tinea, the Cabbage, IV, 36
Tinea pellionella, II, 10
 pustulella, Supp., 58
 tapetzella, II, 10
 vestianella, II, 10
Tineidæ, II, 133
 On a new Genus in, V, 150
Tingis pyri, II, 33
Tiphia femorata, VI, 124
 inornata, VI, 123, 126
 tarda, VI, 126
 transversa, VI, 126
Tipula, I, 180
Tipulidæ, II, 132
tipuliformis, *Ægeria*, II, 10
 Titmouse, Blackcapped, destroying Codling Moth, IV, 28
tityrus, *Epargyrius*, VIII, 173
 Tityrus Skipper, II, 125
 Tobacco Worm, counterworking the, V, 56
Tolmerus, IX, 98
 Tomato-gall, Grape-vine, V, 117
 Tomato-worm, I, 95, IV, 17
 Tortoise-beetle, the Black-legged, II, 63
 Golden, II, 62
 Mottled, II, 63
 Pale-thighed, II, 62
 Tortoise-beetles, I, 100, II, 56, 58, 59, 61
Tortrix botrana, Supp., 57
 cinderella, IV, 47, Supp., 82
 malivorana, IV, 47, Supp., 82
 oxycoccana, Supp., 82
 paludana, VI, 103
 rileyana, I, 153, 154, Supp., 81
 symphoricarpi, I, 154, Supp., 82
 vaccinivorana, Supp., 82
 Townley, John, on Climbing Cut-worms, I, 71
trabeata, *Cassida*, II, 63
 Trabue, A. E., on Army Worm, II, 44, VIII, 27
Tragocephala viridifasciata, VIII, 149
 Transformation of insects, remarks on, III, 146
transversa, *Tiphia*, VI, 126
transversus, *Bruchus*, Supp., 70
 Traps for Canker Worm, VI, 25, 26, VIII, 20, 21
 Codling Moth, I, 66, IV, 23, V, 46
 Plum Curculio, III, 15
 Treadwell, C. C., on parasite of Rocky Mountain Locust, IX, 93
 on Rocky Mountain Locust, IX, 68
tredecim, *Cicada*, I, 19, II, 19, III, 6, VII, 27, Supp., 58, 59
tredecimpunctata, *Hippodamia*, I, 112
tredecimpunctatus, *Sphenophorus*, III, 60, Supp., 54
 Tree-cricket, I, 138
 The Jumping, V, 119
 The Snowy, V, 120
 Tree-hopper, the Buffalo, V, 121
 Tree-hoppers, III, 6
 Trivoltin Silkworms, IV, 85
trichas, *Geothlypis*, VI, 126
Trichina spiralis, IV, 70
Trichobaris, Supp., 54
Trichodactylus, VII, 106
Trichogramma (?) *minuta*, III, 158, Supp., 68
Trichogrammidæ, VI, 142
Trichoptera, V, 16
tricosa, *Agrotis*, Supp., 55
tridens, *Acronycta*, II, 121, Supp., 73
trifasciatus, *Spathius*, V, 106, Supp., 67
trilineata, *Lema*, I, 99, II, 58, III, 14, IV, 8
 Trim Ladybird, II, 25, 27, VII, 39
 Trimble, Dr., on Quince Curculio, III, 36
 Trimeria, a division of Heteroptera, V, 13
trinotatus, *Baridius*, I, 93, III, 60, Supp., 54
Triphlebs, Supp., 58
tristis, *Coreus*, I, 113, II, 31, VII, 46, Supp., 58
tritici, *Diplosis*, II, 10
Trochilium, VI, 27
 acerni, VI, 108
 marginatum, VI, 113, Supp., 72
 tibiale, VI, 113, Supp., 72
Trogosita nana, III, 6, V, 51
troilus, *Papilio*, III, 169
Trombidium holosericeum, VII, 175
 scabrum, VII, 175
 sericeum, VII, 175, IX, 91, Supp., 63
 tinctorium, VII, 175
 True, Dr. U. T., on Locusts, VII, 168
 Trough remedy for Canker Worm, VI, 26
 Truland, N. B., on Rocky Mountain Locust, VII, 150
 Trumpet-gall, Grape-leaf, V, 118
truncatus, *Sphenophorus*, III, 59
Trupanea apivora, I, 168, II, 122, Supp., 60
 vertebrata, II, 123, Supp., 87
Trypeta pomonella, I, 108, III, 6, 91
 solidaginis, I, 13, 173
Trypoxylon, VI, 162
tuckeri, *Oidium*, VI, 30, 63, 79, 85
turbatella, *Dakruma*, Supp., 57
turca, *Leucania*, VIII, 43
Turdus fuscescens, VI, 27
 migratorius, VI, 27
 Turnip Flea-beetle, III, 83
turnus, *Papilio*, VI, 145
 Tusseh Silk-worm, IV, 138
 Tussock Moths, II, 15
 Tuttle, W. F., on Rocky Mountain Locust, IX, 69
 Twelve-spotted Diabrotica, II, 66
 Twice-stabbed Lady-bird, I, 16, V, 100
 Twig-borer, III, 6
 Twig-girdler, III, 6
 Two-lined Soldier-beetle, IV, 29
 Two of our common Butterflies, III, 142
 Two-striped Locust, VII, 124, 173, VIII, 150
 Potato-beetle, II, 61
 Saperda, I, 43, III, 6
Tyloderma fragariæ, Supp., 72
Tyroglyphus echinopus, VII, 106
 entomophagus, VI, 52
 phylloxera, VI, 52, 53, 81
 siro, VI, 52

U.

- ulmi*, *Eriosoma*, I, 123, Supp., 87
 Unadorned Tiphia, VI, 123
 Unarmed Rustic, I, 72, III, 114
undecimlineata, *Doryphora*, VIII, 10
 Uni-banded Ichneumon-fly, III, 77
unicolor, *Corimelæna*, II, 35
 Macrobasis, Supp., 54
unifasciatus, *Ichneumon*, III, 71
unio, *Eudryas*, I, 136, II, 83, III, 63, VI, 90, 92, 95
unipuncta, *Leucania*, I, 109, II, 5, 11, 37, 55, VIII, 22,
 24, 29, Supp., 76
 Noctua, II, 41
unipunctata, *Physonota*, Supp., 53
Urania, VIII, 170
Uroceridæ, V, 10
Uroplata rosea, III, 6
Uropoda americana, IX, 41
 vegetans, IX, 40
Ursula butterfly, III, 163
ursula, *Limenitis*, III, 163, 167, 168, 169, 171
urticæ, *Vanessa*, III, 167
 Urticating larvæ, V, 126
 Useful Labena, VII, 75
Utica (N. Y.) *Herald*, article from, on Paris Green,
 VII, 9

V.

- vadosus*, *Epicærus*, III, 58
vagabundus, *Pemphigus*, I, 112, 120, VII, 97
Valley Farmer, article from, on injury done by
 Cicadas, I, 29
 Van Deman, H. E., on Rocky Mountain Locust,
 IX, 65
 on Rose-Chafer, V, 108
Vanessa antiopa, V, 148
 atalanta, V, 148
 urticæ, III, 167
 Vaporers Moths, II, 15
 Variation in locusts, VIII, 155
 number of antennal joints, II, 89
 wing-venation, IX, 22
varius, *Anthribus*, III, 10
varicornis, *Bruchus*, III, 55, 56, Supp., 69, 70, 71
 Æcanthus, Supp., 61
 Variegated Cut-worm, I, 72, II, 50
variolarius, *Euschistus*, Supp., 58
vastator, *Oscinis*, I, 161
vastatrix, *Phylloxera*, III, 85, IV, 55, VI, 30, 63, 66,
 86, 87, VII, 91, 117, VIII, 157
 Rhizophis, VI, 31
Velleius dilatatus, IV, 22
ventricosus, *Nematus*, VI, 43, 149, IX, 10
Ver du Coeur, II, 107
 Vermes, a section of Segmented Animals, V, 6
vernata, *Anisopteryx*, I, 109, II, 94, VI, 28, VII, 80, 86
 Paleacrita, VIII, 13
 Phalæna, VII, 80
versicolor, *Quiscalus*, VIII, 124
 Vertebrata, a subkingdom of the Animal King-
 dom, V, 6
vertebrata, *Trupanea*, II, 123, Supp., 87
Vespa crabro, IV, 22
 maculata, Supp., 48
 vulgaris, VI, 125
vestalis, *Callimorpha*, III, 133

- vestianella*, *Tinea*, II, 10
villosum, *Elaphidion*, III, 6
 Vinegar for Codling Moth, IV, 27
 Vine Root-borer, I, 124
violaceus, *Corynetes*, VI, 101
Vireo olivaceus, VI, 27, VIII, 124
Vireo, Red-eyed, VIII, 120
virens, *Contopus*, VI, 27
virginiana, *Ortyx*, II, 28, VII, 41
 Virginia Tiger-beetle, I, 115
virginica, *Spilosoma*, III, 68
 Tetracha, I, 115
viridascens, *Glyphe*, II, 53, VIII, 53
viridifasciata, *Tragocephala*, VIII, 149
viridis, *Callochloa*, III, 150
 Caloptenus, VIII, 117
viticida, *Fidia*, I, 132, Supp., 53
viticola, *Botrytis*, VI, 36
vitifolia, *Daktylosphaera*, I, 13
 Pemphigus, I, 13, III, 83, 93, VI, 31
 Phylloxera, II, 27, III, 84, IV, 55
vitis, *Aphis*, I, 13
 Fidia, Supp., 53
 Isosoma, II, 92, 93
 Lasioptera, V, 117
 Madarus, I, 131, Supp., 71
 Procris, II, 86
vitis-coryloides (Gall), V, 116
 -pomum (Gall), V, 114
 -tomatos (Gall), V, 117
 -viticola (Gall), V, 118
vitisana, *Peritymbia*, IV, 55, VI, 31
vitisella, *Carpocapsa*, I, 133
vitivorana, *Penthina*, I, 133, Supp., 57
vitreus, *Mesochorus*, II, 52, VIII, 53
vitripennis, *Capsus*, III, 139
 Campyloneura, III, 137
vittata, *Diabrotica*, I, 100, II, 62, 64, III, 6
 Lytta, I, 96
vittatus, *Chrysops*, II, 129
 Viviparous nature of *Æstrus ovis*, I, 164
 Vorhees, H., on the use of Paris Green, VI, 14
vulgaris, *Vespa*, VI, 125
vulnerator, *Pachymerus*, IV, 28

W.

- Walking-leaves, III, 159
 Walking-stick, III, 159, V, 14, VI, 156, VII, 181
 Wallace, A. R., on Evolution, III, 172
 Wallace, T. D., on Rocky Mountain Locust, IX, 74
 Walnut Case-bearer, IV, 42, V, 49
 Case of, IV, 42
 Natural enemies of, IV, 43
 Walnut Tortrix, I, 153
 Walsh, B. D., controversy on habits of Army
 Worm, II, 47
 description and habits of larva of
 Black Breeze-fly, II,
 130, 131
 of Corn Sphephenorus,
 III, 59
 of Native Currant
 Worm, IX, 26
 of Pickle Worm, II, 67
 of Red-tailed Tachina-
 fly, II, 51

- Walsh, B. D., experiments with *Curculio*-larvæ, I, 55
 on Colorado Potato-beetle, I, 102
 on Hellgrammite larvæ, V, 144
 on Native Currant Worm, IX, 25
 on Oviposition of Katydid, VI, 154
 on Rocky Mountain Locust, VII, 162
 on Thirteen-year Cicada, Supp., 58
- Walshia amorphella*, II, 132, 133
- Ward's *Curculio*-catcher, III, 20
- Warblers destroying Canker Worm, VI, 27
- Warder, R. H., on mould infesting Cicadas, I, 26
- Warrensburg (Mo.) *News*, article from, on Rocky Mountain Locust, VIII, 108
- Water Bugs, a division of Heteroptera, V, 12
- Water-moth, V, 16
- Wax-worm, I, 166
- Weeping Lacewing, II, 26, VI, 51, VII, 40
- weidemeyerii*, *Limenitis*, III, 171
- Western Farmer*, article from, on Birds destroying Canker Worm, VI, 27
- Western Rural*, articles from, on Colorado Potato-beetle, IV, 6, VIII, 8
 article from, on Peach Borer, I, 48
 article from, on Rocky Mountain Locust, VII, 135
- Western Striped Cut-worm, I, 81
- Wetherell, W. H., on Rocky Mountain Locust, IX, 74
- Wheat Cut-worm, I, 87, III, 112
- Wheat-Head Army-worm, IX, 50
 Descriptive, IX, 55
 Habits and natural history, IX, 52
 Natural enemies, IX, 54
 Remedies, IX, 55
- Wheat Midge, II, 10, 13, 16, 70, V, 13, IX, 17
- Wheeler, Wm., on Rocky Mountain Locust, VII, 151
- White, E. M., on counterworking the Tobacco Worm, V, 56
- White, J. D., on Rocky Mountain Locust, IX, 72.
- White, J. K., on Rocky Mountain Locust, VIII, 62, IX, 68
- White, J. W. C., on Rocky Mountain Locust, VII, 152
- White Ant, II, 11, V, 15, IX, 43
 Bark-louse, V, 74
 Grub, I, 88, 156, II, 16, 19, III, 31, 78, IV, 17, 35, VII, 27
 fungus, I, 158, VI, 123, 125
 parasite, VI, 123
 Currant Worms, IX, 14
 Hellebore as remedy for Currant Worms, IX, 14
 -lined Morning Sphinx, III, 140, 141, VIII, 123
 Green Larva of, VIII, 122
 Black Larva of, VIII, 122
 -marked Tussock-moth, I, 144, VI, 29
 -Pine weevil, III, 60
 Worm, Abbot's, IX, 29
 Le Conte's, IX, 32
 -winged Crossbill destroying Canker Worm, VI, 27
- Whitely, Jos., on Rocky Mountain Locust, IX, 71
- Whitescarver, C. S., on Rocky Mountain Locust, IX, 70.
- Whitman, A., on Enemies of Rocky Mountain Locust, IX, 96
- Whittemore, O. A., on Rocky Mountain Locust, VII, 154
- W-marked Cut-worm, I, 79
- Wielandy, J. F., non-publication of report, VII (preface, p. 5)
- Wier, D. B., on remedy for Round-headed Apple-tree Borer, I, 46
- Wier, Mrs. H., on trapping *Curculio*, III, 16
- Wier's Apple-worm Trap, IV, 33, V, 46, VI, 10
- wilcoxi*, *Calosoma*, VIII, 52
- Williams, Prof. A. D., on Rocky Mountain Locust, IX, 65
- Willow-apple Saw-fly, IX, 20
- Wilson, R. P. C., on Rocky Mountain Locust, IX, 75
- Wilson, T. W., on Rocky Mountain Locust, IX, 75
- Wilson's Thrush destroying Canker Worm, VI, 27
- Wings, development of, VI, 40
- Winnipeg Standard*, article from, on Rocky Mountain Locust, IX, 78
- Winter-egg of *Phylloxera*, VI, 42
- Wire Worm, II, 16
- Wise, Jno. C., on Rocky Mountain Locust, VIII, 82, IX, 93
- Witherton, McNeil, on remedy for Canker Worm, II, 101
- Wombat, II, 12
- Wood Nymph, The Beautiful, II, 83, 84, III, 64, VI, 88, 91, 95
 The Pearl, II, 80, 83, 84, III, 63, VI, 90
- Woodpecker, Red-headed, destroying locust, VIII, 124
- Wood Pewee destroying Canker Worm, VI, 27
- Wooly Aphis, III, 95, IV, 100, IX, 43
 Apple-tree louse, I, 118
 -bear, IV, 88
 Elm-tree louse, I, 123
 Plant-louse, I, 119
- Workman, R. A., on Rocky Mountain Locust, VIII, 68, IX, 70
- Wratten, G. L., on Grape *Phylloxera* in California, VI, 82
- Wyoming, Locust History in, VIII, 88, IX, 59
- X.**
- Xabea bipunctata*, Supp., 61
- Xerophylla*, III, 94
caryæ-semen, VII, 117
- Xiphidium*, VI, 155
- Xyela*, IX, 20
- Xyleutes*, VIII, 175
- Xylina*, III, 135
bethunei, III, 136, Supp., 75
capax, III, 136, Supp., 75
cinerea, III, 134, V, 125, Supp., 74
cinerosa, III, 136, Supp., 75
conformis, III, 136, Supp., 75
subcostalis, III, 136, Supp., 75
- xylina*, *Anomis*, II, 37, 40, VI, 17, VIII, 23, Supp., 56
- Ophiura*, II, 40

Y.

- yama-mai*, *Antheraea*, IV, 130
Attacus, IV, 74, 130, 138
 Yama-mai Silkworm, IV, 130
 Acclimatization in Europe and America, IV, 130
 Culture in Japan, IV, 134
 Larval changes, IV, 132
 Parasite, IV, 136
 Value of the cocoon, IV, 133
 Yellow Bear, III, 141
 Yellow-billed Cockoo destroying Canker Worm,
 VI, 28
 Locusts, VIII, 124
 -headed Cutworm, I, 87
 -jacket, VI, 125
 -tail Moth, V, 126
 -tailed Tachina Fly, II, 51, VIII, 53
 Warbler destroying Canker Worm, VI, 27
 Youmans, Prof. E. L., on Evolution, III, 174
 Young, Waller, on Rocky Mountain Locust, VIII, 64
ypsilon, *Agrotis*, Supp., 55
 Yucca, insects affecting, V, 154
 Yucca Borer, VIII, 169, IX, 129
 Affinities, VIII, 176

Yucca Borer—Continued.

- Biological, VIII, 171
 Bibliographical, VIII, 173
 Descriptive, VIII, 174
 Enemies, VIII, 179
 It is single-brooded, IX, 129
 It thrives in the latitude of St. Louis, IX, 129
 Yucca Moth, V, 153, VI, 131, VIII, 37
 Chrysalis, VI, 131
 How the female fertilizes the plant, V, 154
 Larva, V, 155
 Range, V, 159
 Oviposition, VIII, 37
yuccæ, *Castnia*, VIII, 173
 Eudamus, VIII, 173
 Megathymus, V, 129, VIII, 169, 171, 179
yuccasella, *Pronuba*, V, 160, VI, 131, VIII, 171,
 Supp., 58

Z.

- zeæ*, *Anthomyia*, I, 154, Supp., 89
 Sphenophorus, III, 59, Supp., 54
 Zebra Caterpillar, II, 112
zetterstedtii, *Platyptilia*, Supp., 84
 Zimb, V, 13
Zophodia convolutella, Supp., 57

INDEX TO PLANTS AND FOOD-PLANTS.

A.

Abies canadensis, I, 24
Acer saccharinum, IV, 108
 Adam's Needle, V, 158
Æstivalis Grape, V, 65, 66, VII, 103
æstivalis, *Vitis*, III, 89, 90, IV, 60, 61, 63, VI, 36, 47, 48
 72, 74, 75, 80, VII, 103
Agawam Grape, V, 65
Ailanthus, I, 149, 151, III, 130, IV, 75, 82, 112, 118, 120,
 VII, 160
 Tree, A good word for the, IV, 120
Ailanthus glandulosa, IV, 113, 120
alba, *Carya*, V, 105, VII, 117, 118
Morus, IV, 100
album, *Chenopodium*, II, 113, VI, 12
 Alder, II, 121, III, 80
Alnus serrulata, III, 80
aloifolia, *Yucca*, V, 153, VIII, 171, 181, IX, 129
 Alton Large Nutmeg Melon, II, 69
 Alvey Grape, IV, 63, V, 65, VI, 47, VII, 102
amara, *Carya*, V, 104, VII, 118
Amaranthus, VII, 159
blitum, VIII, 61, 119, 121, 122, IX, 122
retroflexus, V, 52, VI, 12
 Amber Grape, VI, 47, 48
Ambrosia, Supp., 54
trifida Supp., 56
americana, *Prunus*, I, 15
 American Grape-vines in Europe, IV, 62, V, 65,
 VII, 116, VIII, 167
 Grape-vines, classification of, IV, 60, VI,
 70
 Ivy, II, 74
americanus, *Ceanothus*, II, 35, VII, 48
Amorpha canescens, II, 90
fruticosa, II, 132, V, 136
Ampelopsis, II, 76, VI, 90
quinquefolia, I, 132, II, 74
Anacharis canadensis, II, 11, IX, 43
Andropogon, VI, 155
Angræcum sesquipedale, V, 153
angustifolia, *Yucca*, V, 157, 159, VIII, 169
Apocynum, III, 144, VIII, 119
 Apple, I, 7, 29, 42, 46, 47, 53, 62, 63, 70, 71, 77, 78, 80, 89,
 108, 118, 119, 126, 128, 144, 146, 150, II, 6, 9,
 15, 88, 89, 90, 91, 95, 96, 114, III, 5, 11, 13, 25,
 30, 32, 33, 34, 38, 57, 58, 92, 102, 114, 118, 120,
 124, 125, 126, 127, 132, 135, 141, IV, 22, 25, 29,
 39, 44, 46, 47, 50, 52, 104, 126, 132, V, 51, 74,
 86, 93, 109, 114, 120, 122, 129, VI, 9, 127, 158,
 VII, 47, 72, 73, 146, 159, 169, VIII, 19, Supp.,
 79

Apple, Ben Davis, IV, 45
 Benoni, IV, 52, V, 87
 Dwarf, I, 69, 70
 Early Harvest, V, 86
 Limber Twig, V, 87
 Lowell, V, 87
 Maiden's Blush, V, 87
 Northern Spy, V, 87
 of Peru, IV, 10
 Rambo, V, 86
 Rawles Janet, III, 34
 Red Astrachan, V, 86
 Red Romanite, V, 86
 Rome Beauty, I, 71
 Soulard, V, 87
 Summer Rose, V, 86
 Sweet June, V, 75
 Tallman's Sweet, III, 35
 Wild Crab, IV, 42, V, 87, Supp., 80
 Willow Twig, V, 87
 Winesap, IV, 45
 Yellow Bellflower, IV, 45
 Apricot, I, 30, III, 11, IV, 29, V, 86, VII, 159
 Aramon Grape, VII, 111
 Arbor vitæ, I, 150, IV, 123, IX, 30
arietinum, *Cicer*, III, 105
Aristida oligostachya, VIII, 122
Aristolochia, II, 116, 118
Aristolochia serpentina, II, 116
sipho, II, 116
tomentosa, II, 116
arizonica, *Vitis*, IV, 60, VI, 73, 76
aromatica, *Rhus*, II, 58, VI, 121, IX, 6
Artemisia, II, 135, V, 35
campestris, I, 175
Asclepias, II, 58, III, 144, 167, V, 152, VIII, 61, 92, 119
cornuti, III, 144
curassavica, III, 144
phytolaccoides, III, 144
purpurascens, III, 144
tuberosa, III, 144
 Ash, III, 126, IV, 123, VII, 72, 160
 Black, VII, 160
 Mountain, I, 43, V, 86, VII, 72
 Ash-leaved Spirea, V, 86
 Asparagus, II, 113, III, 70
 Aster, I, 92, II, 113, 114, III, 105
 Astragalus, Supp., 71
Atropa belladonna, IX, 4
aurantiaca, *Maclura*, IV, 100
aureum, *Ribes*, IX, 2
austriaca, *Pinus*, I, 24, V, 100
 Austrian Pine, V, 100, IX, 30, 32, 33

aviculare, *Polygonum*, Supp., 47

Azalea, IV, 126

B.

Balm of Gilead, II, 89, V, 136, VII, 160

Balsam, II, 114, V, 136

Fir, VII, 172, IX, 30

Banyan, III, 5

Baptisia, V, 136

Barberry, IV, 104, 123

Barley, I, 160, II, 23, III, 112, VII, 27, VIII, 143

Bass-wood, III, 126, IV, 126

batatus, *Ipomea*, II, 56

Bayberry, IV, 123

Bean, I, 79, II, 14, III, 51, 52, 68, 131, VII, 146, Supp., 69, 70

Early Snap, I, 98

English, I, 98

String, III, 105

Windsor, I, 98, III, 51

Bear grass, V, 158

Beech, I, 121, IV, 124, 132, VII, 72

Beet, II, 113, V, 111, 114, VII, 146, 159, VIII, 23, 143

Belladonna, IV, 10

belladonna, *Atropa*, IX, 4

Bellflower, IV, 45

Ben Davis Apple, IV, 45

Benoni Apple, IV, 52, V, 87

Bermuda Sweet-potato, II, 61

Bhotan Pine, V, 100

Bignonia radicans, III, 64, VI, 87

Bind-weed, II, 10

Birch, II, 121, IV, 29, 123, 124, 126, Supp., 73

Bitternut Hickory, V, 104, 105, VII, 97

Bitter-sweet, II, 62

Black Ash, VII, 160

Blackberry, I, 70, 139, IV, 104, V, 108, 120, VI, 111, 127, VII, 159

Dorchester, VI, 113

Black Cherry, Wild, V, 132, 136

Currant, III, 105, IX, 2

July Grape, VII, 102

-gum Elm, V, 78

Hamburg Grape, VI, 76

Henbane, VI, 12

-knot, III, 25

Locust, IV, 82, 142, V, 136, VII, 160

Oak, I, 14, III, 127, V, 132, VI, 115

Prince Grape, VI, 77, 78

Spanish Alabama Grape VII, 107

Sweet Grape, VI, 79

Thorn, III, 36

Walnut, III, 125, 127, 131, IV, 82, VII, 160

blitum, *Amaranthus*, VIII, 61, 119, 121, 122, IX, 123

Blueberry, IV, 126

Blue Dyer Grape, VI, 68

Grass, II, 44, 54, 55, VIII, 27, 49, 122, 182, IX, 47, Supp., 56

Box Elder, VII, 72, 159

Brazilian Sweet-potato, II, 61

Broom-corn, V, 40, VII, 146, 159

Buckwheat, I, 79, III, 109, 141, VII, 43, 146, 159

Burdock, II, 10

Bush Grape, IV, 60, VI, 73

Butter-bean, VII, 146

-cup, II, 10

-nut, III, 68, 127, Supp., 54

C.

Cabbage, I, 79, 83, 84, 156, II, 104, 106, 107, 110, 112, 113, 114, 115, III, 11, IV, 10, 11, 35, 36, V, 111, VI, 158, VII, 47, 159, Supp., 78

californica, *Vitis*, IV, 60, VI, 73

campestris, *Artemisia*, I, 175

Canada Thistle, II, 10

canadense, *Erigeron*, II, 11, VI, 63, IX, 43

canadensis, *Abies*, I, 24

Anacharis, II, 11, IX, 43

Cercis, I, 132, III, 72, 136, Supp., 75

candicans, *Vitis*, IV, 60, VI, 73, 76

canescens, *Amorpha*, II, 90

capitatum, *Croton*, II, 125, V, 147

Carex, VI, 138

carolinense, *Solanum*, I, 103, 107, 108, II, 105, IV, 10, VIII, 122

Carrot, VII, 146, 159, VIII, 143

Carya alba, V, 105, VII, 117, 118

amara, V, 104, VII, 118

glabra, VII, 99, 117

olivæformis, V, 105

porcina, V, 105

Castor Bean, IV, 112, VII, 43, 146, 159

Catalpa, I, 71, 150

Catawba Grape, IV, 62, 63, V, 59, 62, 65, 72, 73, VI, 47, 75, 79, 80, 81, 83, VII, 101, 102, 110, 116

Cayenne Pepper, IV, 10

Ceanothus americanus, II, 35, VII, 48

Cedar, I, 128, II, 91, IV, 51

Red, I, 150, IV, 51, VII, 159, VIII, 119, IX, 30

Celery, VII, 146

Celtis crassifolia, VI, 137

mississippiensis, VI, 137

occidentalis, VI, 137

Cembra Pine, V, 100

Cephalanthus, IV, 123

Cerasus serotina, III, 120

Cercis canadensis, I, 132, III, 72, 136, Supp., 75

Challenge Grape, IV, 63, VI, 47

Chasselas Grape, VII, 102

Cheat, II, 10, VIII, 40, 50

Chenopodium, Supp., 78

album, II, 113, VI, 12

hybridum, VI, 12

Cherry, I, 53, 77, 150, II, 35, 96, 114, III, 11, 32, 57, 58, 120, 127, 163, IV, 39, 41, 104, V, 86, 108, 109, VI, 158, VII, 48, 72, 159, VIII, 19, Supp., 79

Black, V, 132, 136

Mahaleb, VI, 49

Mazzard, VI, 49

Morello, I, 53, VI, 49

Sour, VII, 159

Sweet, VII, 159

Wild, III, 120, IV, 123

Chess, II, 10, 55, VIII, 49

Chestnut, I, 24, II, 121, IV, 132

Horse, I, 146

Spanish, IV, 56

Chickasaw Plum, I, 53

Chickory, IV, 112

Chick-pea, III, 105

Chickweed, I, 179

Chinese Yam, VIII, 143

Choke Cherry, IV, 126, V, 136

Chufa, VIII, 143

cicadina, *Massospora*, Supp., 59
Cichorium sativa, I, 79
Cicer arietinum, III, 105
 Cinquefoil, III, 82
Cirsium lanceolatum, I, 180, VI, 12
 Clara Grape, VI, 79
 Clinton Grape, I, 13, 14, 30, 130, 131, II, 86, 92, III, 87, 89, 91, 92, IV, 62, 63, 64, V, 63, 64, 65, 66, 109, VI, 36, 47, 48, 67, 68, 73, 78, 79, 80, 81, 83, 84, VII, 102, 110, 111, 117
 Clover, II, 113, III, 11, 83, IV, 143, V, 136, VI, 103, VII, 146, 168, VIII, 49, Supp., 78
 Cluster Tomato, V, 118
coccifera, *Quercus*, VII, 96
 Cocklebur, I, 92, III, 60, 105
 Coffee-pea, III, 105
coloratum, *Epilobium*, VI, 90
 Columbia Plum, I, 53
communis, *Ricinus*, IV, 112
 Concord Grape, I, 125, 130, 131, 132, 133, II, 86, III, 72, 87, 92, IV, 62, 63, 65, V, 59, 63, 64, 65, VI, 47, 48, 75, 79, 80, 81, 83, 84, 95, VII, 102, 111, 115, 117
Convolvulus, II, 62, III, 68, IV, 136
copallina, *Rhus*, II, 58, VI, 119
cordata, *Salix*, IX, 20
cordifolia, *Vitis*, III, 87, 88, 89, 90, 91, IV, 60, V, 66, 118, VI, 36, 73, 74, 75, 76, VII, 96
Cordyceps ravenelii, Supp., 53
Coreopsis, II, 35
 Corn, I, 79, 80, 81, 87, 126, 154, 155, 157, 158, II, 16, 23, 27, 28, 42, 43, 44, 54, 55, 89, III, 59, 68, 104, 107, 108, 109, 111, 112, VII, 31, 38, 146, 159, 172, VIII, 25, 27, 49, 123, 143, Supp., 56, 62
 Indian, I, 92, III, 105, V, 40, 123, 136, VII, 169
 Cornucopia Grape, IV, 63, VI, 47
cornuti, *Asclepias*, III, 144
cornutum, *Solanum*, IV, 10
 Corsican Pine, V, 100
cotinus, *Rhus*, VI, 119
 Cotton, I, 150, II, 38, 41, III, 68, 70, 104, V, 136, VI, 17, Supp., 56
 Cottonwood, I, 24, 178, II, 119, VII, 160, IX, 127
 Cow-cockle, II, 10
 Crab-apple, I, 43, 65, III, 25, 30, 31, 40, IV, 39, 42, 52, V, 86, 87
 Transcendent, I, 15
 Crab-grass, III, 111
 Cranberry, Supp., 82
crassifolia, *Celtis*, VI, 137
Crataegus, IV, 42, Supp., 80
 tomentosa, III, 36
 Creeper, I, 132
 Trumpet, III, 64, VI, 87
 Virginia, II, 78, 86, III, 77, VI, 88
Crepis, Supp., 78
 Creveling Grape, I, 72, IV, 63, V, 65, VI, 47, 80
Croton capitatum, II, 125, V, 147
 monanthogynum, V, 147
 Croton Grape, V, 59
 Cucumber, II, 64, 66, 67, 68, 69, 70, III, 111
 Cucurbitaceous vines, II, 64, 66, VII, 159
 Cunningham Grape, IV, 62, 63, V, 64, 65, VI, 47, 48, 78, 81, VII, 96, 111, 115, 117, VIII, 167
curassavica, *Asclepias*, III, 144
 Currant, I, 15, 70, 79, 140, II, 9, 96, III, 68, IV, 104, V, 51, 86, 136, VI, 12, 46, 111, VIII, 121, IX, 1, 2, 3, 5, 7, 25

Currant, Black, III, 105, IX, 2
 Fetid, IX, 2
 Golden, IX, 2
 Missouri, IX, 2
 Red, II, 8, 9, VI, 12, IX, 2
 -flowered, IX, 2
 White, IX, 2
Cynoglossum officinale, V, 101
 Cynthiana Grape, IV, 61, 63, 64, V, 64, 65, VI, 47, 74, VII, 111, 115
Cyperus esculentus, VIII, 143
 Cypress vine, I, 80

D.

Dahlia, I, 92, II, 114, III, 105
 Damson, V, 119
 Dandelion, IV, 143, Supp., 78
Datura, I, 107, IV, 10
 stramonium, V, 56
decandra, *Phytolacca*, VIII, 122
decapetalus, *Helianthus*, IV, 142
 Delaware Grape, I, 130, II, 92, III, 87, IV, 63, V, 59, 63, 65, 66, 111, VI, 47, 73, 80, 81, 83, 84, VII, 106, 110, 116
Delphinium, VII, 185
 Devereux Grape, V, 65
 Diana Grape, IV, 63, V, 59, 65, VI, 47, VII, 96
Dipsacus, IV, 112
discolor, *Solanum*, IV, 10
 Dogbane, III, 144, VIII, 119
 Dog-fennel, II, 10, IV, 15
 Dogwood, V, 86, 93, 136
 Doolittle Raspberry, VI, 111
 Dorchester Blackberry, VI, 113
 Dracut Grape, IV, 63
 Amber Grape, VI, 47, 48
Dryas octopetala, I, 143
 Duane's Early Plum, IV, 142
 Duc de Malacoff Grape, IV, 64
 Duchesse Pear, III, 36
 Dutchman's Pipe, II, 116
 Dwarf Apple, I, 69, 70
 Pear, I, 128
 Sumach, VI, 119

E.

Early Goodrich Potato, I, 110, III, 101, V, 51
 Harvest Apple, V, 86
 Rose Potato, IV, 11
 Snap Bean, I, 98
Echinosperrum strictum, V, 52
 Egg Plant, I, 103, 108, IV, 10
 Elder, III, 100, 101, IV, 104, V, 40
 Elderberry, IV, 104
eleagnifolium, *Solanum*, Supp., 54
 Elm, I, 123, 146, II, 95, 96, 98, 121, III, 73, 127, IV, 104, 126, 129, V, 86, 93, 94, 104, 136, VI, 29, VII, 83, 85, 160, VIII, 19, IX, 127
 White, I, 123
 Elsimboro Grape, IV, 64
Empusa muscae, IV, 88
 Endive, Wild, I, 79, 83
Engelmannia, V, 147
 English Gooseberry, IV, 56, V, 70
 Hawthorn, V, 70
Entomophilæ, V, 152
Epilobium coloratum, VI, 90

Eragrostis poaeoides, VIII, 122
Erigeron canadense, II, 11, VI, 63, IX, 43
Erysiphe, V, 70
esculentus, *Cyperus*, VIII, 143
Eucalyptus globulus, VI, 55
Eumelan Grape, V, 65, VII, 101
Eupatorium perfoliatum, VI, 12
 serotinum, I, 152
European Grape, IV, 55, 63, VI, 47, 77
 Larch, VII, 172
 Rulander Grape, VI, 77, 78
excelsa, *Pinus*, V, 100

F.

falcata, *Quercus*, VIII, 182
False Indigo, II, 132, V, 136
fasciculata, *Vernonia*, I, 153
Fetid Currant, IX, 2
Field Garlic, II, 10
Figwort family, VII, 14
filamentosa, *Yucca*, V, 158, 159, VIII, 171
Fir, IX, 29
 Balsam, VII, 172, IX, 30
flaccida, *Yucca*, V, 159
Flax, II, 42, VII, 42, 159, VIII, 25
floridum, *Ribes*, IX, 2
Foment Grape, V, 66
Fox Grape, VI, 75, 84
Fox-tail Grass, II, 10
Fragrant Sumach, VI, 121, IX, 6
French Hazel, VI, 121
Frost Grape, III, 87, 89, 90, IV, 60, VI, 73
fruticosa, *Amorpha*, II, 132, V, 136
Fungus infesting Cicada, I, 26
 White Grub, I, 158, VI, 123, 125
Furze, III, 51

G.

Garden Gooseberry, IX, 2
Garlic, Field, II, 10
Garrigues Grape, VI, 84
Geranium, III, 68
glabra, *Carya*, VII, 99, 117
 Photinia, IV, 132
 Rhus, II, 58, VI, 119
Gladiolus, III, 105
glandulosa, *Ailanthus*, IV, 113, 120
glauca, *Yucca*, V, 153
globulus, *Eucalyptus*, VI, 55
gloriosa, *Yucca*, V, 159, VIII, 171
Goat-weed, II, 125, V, 147
Gæthe Grape, IV, 63, V, 62, 64, 65, 66, VI, 47, 80, 81,
 VII, 96, 111, 116
Golden Chasselas Grape, VI, 77
 Clinton Grape, III, 87, VI, 47, 48, 68, VII, 102
 Currant, IX, 2
 -rod, I, 98, 152, 173, II, 134, IV, 28, Supp., 57
 Willow, III, 168
Gooseberry, I, 140, II, 8, 9, III, 58, 68, VI, 46, VIII,
 121, IX, 1, 2, 3, 5, 6, 8, 26
Gooseberry, English, IV, 56, V, 70
 Garden, IX, 2
 Houghton's Seedling, I, 140
 Showy, IX, 2
Goosefoot, II, 113, VI, 12
Gourd, II, 66

Grape, Agawam, V, 65
 Alvey, IV, 63, V, 65, VI, 47, VII, 102
 Amber, VI, 47, 48
 Aramon, VII, 111
 Black Hamburg, IV, 56, VI, 76
 July, VII, 102
 Prince, VI, 77, 78
 Spanish Alabama, VII, 107
 Sweet, VI, 79
 Blue Dyer, VI, 68
 Bullet, IV, 62, VI, 75
 Bush, IV, 60, VI, 73
 Catawba, IV, 62, 63, V, 59, 62, 65, 72, 73, VI,
 47, 75, 79, 80, 81, 83, VII, 101, 102, 110, 116
 Challenge, IV, 63, VI, 47
 Chasselas, IV, 56, VII, 102
 Clara, VI, 79
 Clinton, I, 13, 14, 30, 130, 131, II, 86, 92, III,
 87, 89, 91, 92, IV, 62, 63, 64, V, 63, 64, 65, 66,
 109, VI, 36, 47, 48, 67, 68, 73, 78, 79, 80, 81, 83,
 84, VII, 102, 110, 111, 117
 Concord, I, 125, 130, 131, 132, 133, II, 86, III,
 72, 87, 92, IV, 62, 63, 65, V, 59, 63, 64, 65, VI,
 47, 48, 75, 79, 80, 81, 83, 84, 95, VII, 102, 111,
 115, 117
 Cornucopia, IV, 63, VI, 47
 Creveling, I, 72, IV, 63, V, 65, VI, 47, 80
 Croton, V, 58
 Cunningham, IV, 62, 63, V, 64, 65, VI, 47, 48,
 78, 81, VII, 96, 111, 115, 117, VIII, 167
 Cynthiana, IV, 61, 63, 64, V, 64, 65, VI, 47, 74,
 VII, 111, 115
 Delaware, I, 130, II, 92, III, 83, 87, IV, 63, V,
 59, 63, 65, 66, 111, VI, 47, 73, 80, 81, 83, 84,
 VII, 106, 110, 116
 Devereux, V, 65
 Diana, IV, 63, V, 59, 65, VI, 47, VII, 96
 Dracut, VI, 47, 48
 Amber, IV, 63
 Duc de Malacoff, IV, 64
 Elsinboro, IV, 64
 Eumelan, V, 65, VII, 101
 European, IV, 55, 63, VI, 47, 77
 Rulander, VI, 77, 78
 Foment, V, 66
 Fox, VI, 75, 84
 Frost, III, 87, 89, 90, IV, 60, VI, 73
 Garrigues, VI, 84
 Gæthe, IV, 63, V, 62, 64, 65, 66, VI, 47, 80, 81,
 VII, 96, 111, 116
 Golden Chasselas, VI, 77
 Clinton, III, 87, VI, 47, 48, 68, VII,
 102
 Hamburg, V, 59
 Hartford, III, 72, IV, 63, VI, 47
 Prolific, I, 125, 130, 131, IV, 62, 64,
 65, V, 65, VI, 75, 81, VII, 116
 Herbemont, IV, 61, 62, 63, 64, V, 63, 64, 65, VI,
 36, 47, 48, 74, 78, 80, 81, VII, 102, 109, 111,
 115, 117, VIII, 167
 Hermann, V, 65, VII, 111
 Huntington, III, 87
 Iona, III, 72, V, 59, 62, 65, VI, 48, 79, 80, 81, 83,
 VII, 96, 116
 Isabella, I, 130, II, 81, IV, 62, 63, 64, V, 59, VI,
 48, 75, 81, VII, 96, 101, 102

Grape, Israella, III, 72
 Ives, IV, 63, V, 65, VI, 48, 81, 84, VII, 96, 111, 116
 Seedling, I, 133, V, 64, VI, 79, 80
 Jacques, V, 66, VII, 107, 117, VIII, 167
 Lenoir, IV, 64, V, 66, VII, 108, 117
 Lindley, V, 65
 Long, V, 66
 Longworth's Ohio, VII, 107
 Louisiana, IV, 63, V, 65, VI, 47
 Madam Pince, IV, 64
 Madeira, V, 63
 Malaga, VI, 78
 Malvasia, VI, 77
 Marion, VI, 47, 48, 68
 Martha, IV, 62, 63, VI, 48, VII, 96, 111
 Massasoit, V, 65
 Maxatawney, IV, 63, 64, V, 65, VI, 48, 80, 81, VII, 96, 116
 Merrimac, V, 65
 Muscadine, IV, 60, 62, 63, VI, 72
 Muscat, VII, 102
 Hamburg, IV, 64
 Mustang, IV, 60, 62, 64, VI, 73, 76, VIII, 167
 North Carolina, III, 72, IV, 63, V, 64, VI, 48, VII, 96, 111
 Northern Fox, III, 87, 90, IV, 60, 63, VI, 47, 72, 75, VII, 106
 Muscadine, VI, 48, VII, 102
 Norton's, V, 65, VI, 78, 84
 Catawba, VIII, 96
 Virginia, I, 132, IV, 62, 63, 64, VI, 47, 80, 81, VII, 109, 111, 115
 Ohio, VII, 107
 Othello, IV, 63, VI, 47
 Pauline, IV, 64, VII, 102
 Post Oak, VI, 79
 Rebecca, V, 59, VI, 48
 Rent, V, 64, VI, 81, VII, 115
 River Bank, IV, 60, 63, V, 116, 117, VI, 36, 47
 Riverside, VI, 72
 Rogers, I, 130, V, 66
 Hybrid, I, 130, V, 117, VI, 80
 No. 4, I, 130, II, 92
 Rulander, V, 65, VI, 47, 80, VII, 111
 European, VI, 77, 78
 Salem, V, 65, 66, VI, 48, 79
 Sand, IV, 60, VI, 73
 Scegety, V, 66
 Scuppernong, III, 77, IV, 62, 64, VI, 50, 76, VII, 106
 Segar Box, VII, 107
 Sonora, IV, 64
 Southern Fox, III, 77, IV, 60, 62, 63, VI, 48, 72, 75
 St. Augustine, V, 63
 Sugar, IV, 61
 Summer, III, 89, 90, IV, 60, 63, VI, 47, 72
 Taylor, I, 30, II, 86, III, 87, IV, 63, 64, V, 63, 65, 66, VI, 36, 47, 48, 68, 80, 95, VII, 102, 115
 Bullet, VI, 73
 Telegraph, V, 65, VI, 47, VII, 106
 Tinto, VI, 36
 Tokay, V, 66
 Venango, VI, 84

Grape, Virginia Seedling, VI, 61, 74
 Walter, V, 59, 62, VI, 80
 Warren, VII, 102
 Weehawken, V, 59
 White Riesling, VII, 102
 Scotch Cluster, VI, 77
 Wilder, IV, 63, V, 62, 63, VI, 48, 81, VII, 101, 102, 111, 116
 Winter, IV, 60, VI, 73
 York Madeira, IV, 64, V, 64
 Grape Mildew, IX, 43
 -vine, I, 30, 70, 72, 78, 80, 124, 128, 129, 131, 132, 133, 136, 137, 138, 180, II, 71, 74, 76, 78, 81, 82, 83, 85, 86, 87, 88, 89, 91, 92, III, 11, 61, 63, 65, 68, 70, 72, 75, 77, 79, 81, 84, 111, 130, 137, 141, IV, 52, 54, 55, V, 54, 57, 108, 112, 114, 116, 117, 118, 120, VI, 30, 87, 88, 91, 95, 111, VII, 47, 91, 146, 159, 172, VIII, 121, 123, 157, Supp., 75
 disease, IV, 55, VI, 58, 65
 -vines, American, Classification of, VI, 70
 in Europe, IV, 62, V, 65, VII, 116, VIII, 167
 of the United States, The true, IV, 60, VI, 70
 rooting of, V, 65
 Varieties to graft, VI, 81
 use as stock in grafting, VII, 115

Green Citron Melon, II, 69
 Greengage Plum, I, 140
grossulariæ, *Ribes*, IX, 2
 Ground-cherry, I, 107, IV, 10
 Gympson-weed, I, 107

H.

Hackberry, V, 119, VI, 137
 Hackmatack, VII, 169
Halesia, IV, 123
Hamamelis, III, 120
 Hamburg Grape, V, 59
 Hard Maple, III, 126, Supp., 55
 Harrison Potato, I, 110
 Hartford Grape, III, 72, IV, 63, VI, 47
 Prolific Grape, I, 125, 130, 131, IV, 62, 64, 65, V, 65, VI, 75, 81, VII, 116
 Haw, I, 108, III, 25, 32, 35, 38, 92, V, 51
 Hawthorn, I, 43, III, 36
 English, V, 70
 Hazel, III, 11, 37, 75, IV, 104, 126, V, 132
 French, VI, 121
 Heart Cherry, VIII, 121
 -shaped Willow, IX, 20
 Hedge-mustard, VI, 12
Helianthus, VII, 159, Supp., 53
 decapetalus, IV, 142
 petiolaris, V, 52
 tuberosus, VIII, 143
 Hellebore, White, IX, 13
 Hemp, III, 105, VII, 42, 146
 Henbane, IV, 10
 Herbemont Grape, IV, 61, 62, 63, 64, V, 63, 64, 65, VI, 36, 47, 48, 74, 78, 80, 81, VII, 102, 109, 111, 115, 117, VIII, 167
 Herds-grass, VII, 168

Hermann Grape, V, 65, VII, 111
Hibiscus militaris, VI, 92
 Hickory, I, 126, 153, 154, III, 37, 124, 126, 127, 131, 135,
 IV, 42, 43, 52, 54, 104, 124, 126, 140, V, 104,
 119, VI, 64, 101, VII, 160, Supp., 81
 Bitternut, V, 104, 105, VII, 97
 Pecan, V, 105
 Pignut, V, 105
 Shagbark, II, 33, IV, 54
 Shellbark, IV, 66, V, 103, 105
 Honey-locust, I, 92, 150, III, 45, IV, 104, 126, VII, 159
 -suckle, II, 113
 Hop-plantain, III, 131
 -vine, V, 136
 Horse Chestnut, I, 146
 Gentian, III, 134
 -nettle, I, 103, 104, 107, 108, IV, 10
 Weed, II, 11, IX, 43
 Houghton's Seedling Gooseberry, I, 140
 Hound's-tongue, V, 101
humilis, *Salix*, V, 132
 Hungarian Grass, VII, 27, 28, VIII, 29, 39
 Huntington Grape, III, 87
hybridum, *Chenopodium*, VI, 12
hydropiper, *Polygonum*, III, 70, VI, 12
Hyoscyamus, IV, 10
 niger, VI, 12

I.

incana, *Quercus*, IV, 114
 Indian Corn. (See Corn.)
 Indigo, V, 136
 False, II, 132, V, 136
infectoria, *Quercus*, V, 18
 Iona Grape, III, 72, V, 59, 62, 65, VI, 48, 79, 80, 81, 83,
 VII, 96, 116
Ipomea, III, 45
 batatas, II, 56
 Irish Potato. (See Potato.)
 Ironweed, V, 136
 Isabella Grape, I, 130, II, 81, III, 72, IV, 62, 63, 64, V,
 59, VI, 48, 75, 81, VII, 96, 101, 102
 Israella Grape, III, 72
 Ives Grape, IV, 63, V, 65, VI, 48, 81, 84, VII, 96, 111,
 116
 Seedling Grape, I, 133, V, 64, VI, 79, 80
 Ivy, Poison, VI, 121

J.

Jamestown Weed, I, 107, II, 10, V, 56
 Japan Varnish tree, IV, 120
 Jaques Grape, V, 66, VII, 107, 117, VIII, 167
 Jerusalem Artichoke, VIII, 143
jujuba, *Rhamnus*, IV, 138
 Jujube, V, 18
 June-berry, I, 43
Juniperus virginiana, I, 24

K.

King of the Earlies Potato, IV, 11
 Knotweed, Supp., 47
 Kolrabi, VII, 159
 Kunogi, IV, 130, 134, 136

L.

labrusca, *Vitis*, III, 87, 89, 90, IV, 60, 63, V, 60, 65, 66,
 118, VI, 36, 47, 48, 71, 72, 74, 75, 76, VII, 103

Laburnum, IV, 118
 Lamb's quarter, II, 10, 113, VI, 12
lanceolatum, *Cirsium*, I, 180, VI, 12
 Larch, European, VII, 172
laricio, *Pinus*, V, 100
Lathyrus, III, 52
 Laurel, V, 33
 -cherry, V, 33
 -leaved oak, IV, 134
 Lawrence Pear, III, 36
 Lenoir Grape, IV, 64, V, 66, VII, 108, 117
 Lettuce, IV, 100, 112, VI, 158
 Lilac, III, 68, IV, 104, 123, VII, 159
 Persian, I, 15, V, 70, 86, 127
 Lima Bean, VII, 172
 Limber Twig Apple, V, 87
lincecumii, *Vitis*, VI, 74
 Linden, I, 150, V, 93, VII, 72
Lindera, II, 121, IV, 123
 Lindley Grape, V, 65
Liquidambar, IV, 123
Liriodendron, IV, 123
 tulipifera, Supp., 55
 Live-oak, IV, 129
 Locust, I, 24
 Black, IV, 82, 142, V, 136, VII, 160
 Honey, I, 98, 150, III, 45, IV, 104, 126, VII, 159
 Lombardy Poplar, I, 150, 157, II, 89, VII, 160
 Long Grape, V, 66
 Longworth's Ohio Grape, VII, 107
 Louisiana Grape, IV, 63, V, 65, VI, 47
 Lowell Apple, V, 87
 Lucern, III, 83, 105
Lycopersicum, IV, 10

M.

Maclura aurantiaca, IV, 100
 Madam Pince Grape, IV, 64
 Madeira Grape, V, 63
Madia sativa, VI, 55
 Mahaleb Cherry, VI, 49
 Maiden's Blush Apple, V, 87
 Malaga Grape, VI, 78
 Mallow, VI, 89
Malva, VI, 89
 sylvestris, VIII, 182
 Malvasia Grape, VI, 77
 Mandrake, IV, 15
 Mangel wurzel, VIII, 143
 Maple, I, 47, 146, 150, II, 121, IV, 104, 123, 126, VI, 107,
 111, VII, 160
 Hard, III, 126, Supp., 55
 Silver, I, 150, V, 137
 Soft, I, 47, 150, IV, 42, V, 120, 137, VI, 108,
 Supp., 55, 62, 74
 Mare's Tail, II, 11
 Marigold, II, 114
 Marion Grape, VI, 47, 48, 68
 Martha Grape, IV, 62, 63, VI, 48, VII, 96, 111
Maruta, VIII, 100
Massospora cicadina, Supp., 59
 Massasoit Grape, V, 65
 Maxatawney Grape, IV, 63, 64, V, 65, VI, 48, 80, 81,
 VII, 96, 116
 May Apple, IV, 15
 -weed, II, 10

Mazzard Cherry, VI, 49
 Meadow-sweet, III, 51
media, *Stellaria*, I, 179
 Medlar, V, 86
 Neapolitan, IV, 132
 Melon, II, 64, 66, 69, 70
 Alton Large Nutmeg, II, 69
 Green Citron, II, 69
 Merrimac Grape, V, 65
 Mignonette, II, 113
militaris, *Hibiscus*, VI, 92
 Milkweed, I, 139, II, 58, III, 133, 144, 168, VIII, 61, 92, 119
 Miner Plum, I, 53
mississippiensis, *Celtis*, VI, 137
 Missouri Currant, IX, 2
missouriensis, *Solidago*, I, 174
mitis, *Pinus*, V, 100
monanthogynum, *Croton*, V, 147
monilifera, *Populus*, II, 119
monoica, *Strombocarpa*, I, 65
monticola, *Vitis*, VI, 57, 74
 Morello Cherry, I, 53, VI, 49
moretti, *Morus*, IV, 100
 Morning Glory, I, 100, II, 62
Morus alba, IV, 100
 moretti, IV, 100
 multicaulis, IV, 80, 100
 rubra, IV, 100
 Mountain Ash, I, 43, V, 86, VII, 72
 Mulberry, IV, 74, 75, 76, 79, 82, 100
 Red, IV, 100
 White, I, 72, 73
 Mullein, II, 10, V, 35, VII, 14
multicaulis, *Morus*, IV, 80, 100
 Muscardine Grape, IV, 60, 62, 63, VI, 72
muscae, *Empusa*, IV, 88
 Muscardine, IV, 88, 89, 91, 144
 Muscat Grape, VII, 102
 Hamburg Grape, IV, 64
 Mustang Grape, IV, 60, 62, 64, VI, 73, 76, VIII, 167
mustangensis, *Vitis*, IV, 62, VI, 76
 Mustard, IV, 36, V, 112
 Hedge, VI, 12
Myrica, IV, 123

N.

Nansemond Sweet-potato, II, 61
 Neapolitan Medlar, IV, 132
 Neck-weed, II, 35, VII, 48
 Nectarine, III, 40
nemoralis, *Solidago*, I, 173
 Nettle, II, 105, IV, 10, VIII, 122
 Horse, I, 103, 104, 107, 108, IV, 10
 New Jersey Tea-plant, II, 35, VII, 48
Nicandra, IV, 10
Nicotiana, IV, 10
niger, *Hyoscyamus*, VI, 12
 Nightshade, IV, 10, VI, 12
nigrum, *Ribes*, IX, 2
 Solanum, VI, 12
 North Carolina Grape, III, 72, IV, 63, V, 64, VI, 48, VII, 96, 111
 Northern Fox Grape, III, 87, 90, IV, 60, 63, VI, 47, 72, 75, VII, 106
 Muscadine Grape, VI, 48, VII, 102
 Spy Apple, V, 87

Norton's Grape, V, 65, VI, 78, 84, VII, 96
 Virginia Grape, I, 132, IV, 62, 63, 64, VI, 47, 80, 81, VII, 109, 111, 115
 Norway Spruce, I, 150, VII, 172, VIII, 119, IX, 30
novaeboracensis, *Vernonia*, VIII, 119

O.

Oak, I, 47, 126, 128, 139, 146, II, 91, III, 73, 94, 124, 125, 126, 127, 131, 134, 138, IV, 45, 52, 114, 126, 137, 140, V, 18, 126, 127, 132, 139, VI, 103, 113, 127, 128, 158, 166, VII, 72, VIII, 23, IX, 52
 Black, I, 14, V, 132, VI, 115
 Chermes, VII, 96
 Laurel-leaved, IV, 134
 Live, IV, 129
 Pin, I, 157
 Post, I, 157, IV, 42, 66, 134, V, 132, VI, 64, 115, VII, 97
 Red, I, 14, V, 132, VI, 115
 White, VI, 64, 115
 Oat, I, 88, II, 16, 44, 54, III, 111, 112, 115, VI, 12, VII, 38, 146, VIII, 27, 49, 119
occidentalis, *Celtis*, VI, 137
 Thuja, I, 24
octopetala, *Dryas*, I, 143
officinale, *Sisymbrium*, VI, 12
 Ohio Grape, VII, 107
Oidium tuckeri, V, 57, 70, VI, 30, 63, 79, 85, IX, 43
 Proof of its occurrence in America, V, 70
oligostachya, *Aristida*, VIII, 122
olivæformis, *Carya*, V, 105
 Onion, II, 9, VII, 159, 169, VIII, 49
 Osage Orange, I, 126, 150, II, 89, III, 131, IV, 75, 79, 100, VII, 159, IX, 95, Supp., 89
 Othello Grape, IV, 63, VI, 67
 Ox-eye Daisy, II, 10

P.

Panicum sanguinale, VIII, 122
 Parsnip, VII, 146, VIII, 119, 143
 Pauline Grape, IV, 64, VII, 102
 Pea, II, 14, 42, III, 44, 50, 68, 105, 107, VIII, 25, 119
 Chick, III, 105
 Coffee, III, 105
 Peach, I, 47, 50, 77, II, 15, III, 15, 27, 30, 34, 38, 40, 57, 103, 105, 114, 120, 127, 132, 134, IV, 22, 29, 40, 52, 82, V, 108, 120, 127, 129, VI, 112, VII, 72, 146, 159, VIII, 19, Supp., 75
 -blow Potato, I, 79, 98, 99, III, 101
 Rot, I, 52
 Peanut, VII, 146
 Pear, I, 15, 43, 64, 69, 70, 77, 128, 146, 150, II, 33, 38, 114, III, 11, 33, 36, 38, 57, 78, 120, 131, IV, 40, 52, 104, V, 54, 86, 93, 122, VII, 72, 146, 159, VIII, 24
 blight, III, 58, VIII, 24
 Duchess, III, 36
 Dwarf, I, 128
 Lawrence, III, 35
 Seckel, III, 35
 Standard, I, 128
 White Doyenne, I, 15
 Pecan, V, 105, VI, 101
peltatum, *Podophyllum*, IV, 15
peregrina, *Veronica*, II, 35, VII, 48

- perfoliatum*, *Eupatorium*, VI, 12
Triosteum, III, 134
Peronospora, V, 70, VI, 85
 Persian Cantelope, II, 69
 Lilac, I, 15, V, 70, 86, 127
 Persimmon, V, 69, 109
petiolaris, *Helianthus*, V, 52
 Petunia, IV, 10
Phaseolus, III, 52, 53
Photinia glabra, IV, 132
Physalis, I, 107, IV, 10
Phytolacca decandra, VIII, 122
phytolaccoides, *Asclepias*, III, 144
 Pig-nut Hickory, V, 105
 -weed, V, 52, VI, 12
 Pimpernel, IV, 118
 Pine, I, 24, 127, II, 15, 91, V, 100, IX, 29, 32
 Austrian, V, 100, IX, 30, 32, 33
 Bhotan, V, 100
 Cembra, V, 100
 Corsican, V, 100
 Pitch, V, 100, IX, 32
 Pyrenaian, V, 100
 Scotch, III, 92, V, 100, IX, 30, 32, 33
 White, III, 92, V, 97, 100, 102, IX, 13, 29, 30, 32
 Yellow, V, 100
 Pin Oak, I, 157
Pinus austriaca, I, 24, V, 100
 cembra, V, 100
 excelsa, V, 100
 laricio, V, 100
 mitis, V, 100
 pumilio, V, 100
 pyrenaica, V, 100
 resinosa, V, 100
 strobis, I, 24
 sylvestris, I, 24, V, 100
 Pitch Pine, V, 100, IX, 32
Plantago, IV, 142
 Plantain, II, 10, III, 68, IV, 142, 143
 Plum, I, 15, 65, 140, 146, 150, II, 15, 96, III, 11, 25, 27,
 32, 34, 40, 41, 57, 103, 120, 127, 153, 163, IV, 23,
 29, 39, 41, 104, 118, 123, 124, 126, V, 86, 93, 109
 VI, 127, 141, VII, 72, 159, IX, 2
 Chickasaw, I, 53
 Columbia, I, 53
 Duane's Early, IV, 142
 Greengage, I, 140
 Miner, I, 53
 Wild, I, 15, 55
poaeoides, *Eragrostis*, VIII, 122
Podophyllum peltatum, IV, 15
 Poison Ivy, VI, 121
 Pokeweed, VIII, 122
 Poke Milk-weed, III, 144
 Polecat-weed, VI, 121
Polygonum aviculare, Supp., 47
 hydropiper, III, 70, VI, 12
 Poplar, II, 91, III, 72, 73, 120, 127, 135, 153, 168, IV,
 123, 126, V, 136, VI, 105
 Lombardy, I, 150, 157, II, 89, VII, 160
 Silver, III, 156, 168
 Silver-leaf, VII, 160
Populus monilifera, II, 119
arcina, *Carya*, V, 105
 Post Oak, I, 157, IV, 42, 66, 134, V, 132, VI, 64, 115,
 VII, 97
 Post Oak Grape, VI, 79.
 Potato, I, 91, 93, 95, 96, 97, 98, 99, 100, 101, 158, II,
 42, 56, 57, 70, 114, III, 98, 105, 111, IV, 5,
 10, 11, V, 18, 111, 112, 114, VI, 11, VII, 2, 47,
 146, 159, 169, 172, VIII, 1, 25, 37, 119, IX,
 39
 Chili No. 2, III, 101
 Early Goodrich, I, 100, III, 101, V, 51
 Rose, III, 101, IV, 11
 Harrison, I, 110
 King of the Earlies, IV, 11
 Mercer, III, 101
 Peach-blow, I, 97, 98, 99, III, 101
 Peerless, III, 101
 Pink-eye, III, 101
 Quaker Russet, I, 98
 Russet, III, 101
 Shaker, III, 101
Potentilla, III, 82
 verna, I, 143
Poterium sanguisorba, I, 143
 Prickly Mesquit Grass, VII, 192
 Prostrate Currant, IX, 2
prostratum, *Ribes*, IX, 2
 Prune, III, 40
Prunus, VI, 141
 americana, I, 15
 lauro-cerasus, V, 33
 serotina, V, 136
puberula, *Yucca*, V, 153
pumilio, *Pinus*, V, 100
 Pumpkin, I, 79, II, 42, III, 105, VIII, 25
 Purple-fringe, VI, 119
purpurascens, *Asclepias*, III, 144
 Purslane, II, 10, III, 112, 141, V, 69, VI, 158, VII,
 46, 47, VIII, 122, 123
 Speedwell, II, 35, VII, 48
 Pyrenaian Pine V, 100
pyrenaica, *Pinus*, V, 100
- Q.**
- Quaker Russet Potato, I, 98
Quercus coccifera, VII, 96
 falcata, VIII, 182
 incana, IV, 114
 infectoria, V, 18
 serrata, IV, 130
 Quince, I, 43, 65, 150, II, 35, 114, III, 30, 35, 36, 38,
 IV, 39, 41, 126, 132, VII, 48, Supp., 79
quinquefolia, *Ampelopsis*, I, 132, II, 74
- R.**
- racemosus*, *Symphoricarpus*, II, 113
radicans, *Bignonia*, III, 64, VI, 87
 Radish, I, 156, IV, 36, V, 111, VII, 159
 Rambo Apple, V, 86
 Raspberry, I, 70, 139, II, 34, III, 72, V, 120, 123,
 VI, 111, VII, 48
 Doolittle, VI, 111
ravenelii, *Cordyceps*, Supp., 53
 Rawles Janet Apple, III, 34
 Rebecca Grape, V, 59, VI, 48

Red Astrachan Apple, V, 86
 Bud, III, 72
 Cedar, I, 150, IV, 51, VII, 159, VIII, 119, IX, 30
 Currant, II, 8, 9, VI, 12, IX, 2
 -flowered Currant, IX, 2
 June Apple, IV, 52
 Mulberry, IV, 100
 Oak, I, 14, V, 132, VI, 115
 Pine, V, 100
 Romanite Apple, V, 86
 -root, II, 35, IV, 104, VII, 48
 Rentz Grape, V, 64, VI, 81, VII, 115
resinosa, *Pinus*, V, 100
retroflexus, *Amaranthus*, VI, 12
Rhamnus jujuba, IV, 138
 Rhubarb, II, 123, III, 51
Rhus aromatica, II, 58, VI, 121, IX, 6
 copallina, II, 58, VI, 119
 coriaria, IV, 118
 cotinus, VI, 119
 glabra, II, 58, VI, 119
 toxicodendron, V, 127, VI, 121
 typhina, VI, 19
Ribes aureum, IX, 2
 floridum, IX, 2
 grossularia, IX, 2
 nigrum, IX, 2
 prostratum, IX, 2
 rubrum, IX, 2
 sanguineum, IX, 2
 speciosum, IX, 2
Ricinus communis, IV, 112
riparia, *Vitis*, IV, 60, 61, 63, V, 62, 65, 116, 118, VI, 36, 47, 48, 58, 72, 73, 74, 75, 95, VII, 96
 River Bank Grape, IV, 60, 63, V, 116, 117, VI, 36, 47
 Riverside Grape, VI, 72
robustum, *Solanum*, IV, 10
 Rogers' Grape, I, 30, II, 92, V, 66
 Hybrid Grape, I, 30, V, 117, VI, 80
 Rome Beauty Apple, I, 71
Rosaceæ, V, 86
 Rose, I, 70, 146, III, 120, 124, 127, IV, 126, V, 109, 123, 127, VI, 127, VII, 159, Supp., 57
 Wild, V, 126
rostratum, *Solanum*, I, 102, 108, IV, 10, VII, 1
rubra, *Morus*, IV, 100
rubrum, *Ribes*, IX, 2
Rubus, V, 154, VI, 113, Supp., 72
 Rulander European Grape, VI, 77, 78
 Grape, V, 65, VI, 47, 48, VII, 111
rupestris, *Vitis*, IV, 60, VI, 73, 74
rupicola, *Yucca*, V, 157
 Russet Potato, III, 101
 Rutabaga, II, 113, VII, 159, VIII, 143
 Rye, I, 160, II, 29, 44, 54, III, 111, VII, 38, 146, 168, VIII, 27, 49, IX, 51

S.

saccharinum, *Acer*, IV, 108
 Salem Grape, V, 65, 66, VI, 48, 79
Salix, VI, 136
 cordata, IX, 20
 humilis, V, 132
Salvia trichostemmoides, VIII, 119

Sand Bur, VIII, 9, 122
 Grape, IV, 60, VI, 73
sanguinale, *Panicum*, VIII, 122
sanguineum, *Ribes*, IX, 2
sanguisorba, *Poterium*, I, 143
Sarracenia, III, 155
 Sassafras, IV, 123, V, 122, 134, 136, VI, 127
sativa, *Cichorium*, I, 79
 Madia, VI, 55
scandens, *Senecio*, Supp., 78
 Scegety Grape, V, 66
 Scotch Pine, III, 92, V, 100, IX, 30, 32, 33
 Screw-bean, I, 65
 Scrub Oak, III, 163
 Willow, II, 90, III, 168, V, 132
 Scuppernong Grape, III, 77, IV, 62, 64, VI, 50, 76, VII, 106
 Seckel Pear, III, 36
 Segar-box Grape, VII, 107
Senecio scandens, Supp., 78
serotina, *Cerasus*, III, 120
 Prunus, V, 136
serotinum, *Eupatorium*, I, 152
serpentina, *Aristolochia*, II, 116
serrata, *Quercus*, IV, 130
serrulata, *Alnus*, III, 80
sesquipedale, *Angræcum*, V, 153
setigera, *Stipa*, VII, 192
 Shag-bark Hickory, II, 33, IV, 54, VII, 48
 Shaker Potato, III, 101
 Shell-bark Hickory, IV, 66, V, 103, 105
 Shepherd's Purse, II, 10
 Showy Gooseberry, IX, 2
sieglinge, *Solanum*, IV, 10
 Silkweed, III, 144
 Silver Maple, I, 150, V, 137
 Poplar, III, 156, 168
 -leaf Poplar, VII, 160
sipho, *Aristolochia*, II, 116
Sisymbrium officinale, VI, 12
 Smartweed, II, 10, III, 68, 70, VI, 12
 Smoke-tree, VI, 121
 Smooth Sumach, VI, 119
 Snowberry, I, 153, 154, II, 113, Supp., 81, 82
 Snowdrop-tree, IV, 123
 Soft Maple, I, 47, 150, IV, 42, V, 120, 137, VI, 108, Supp., 55, 62, 74
Solanaceæ, IV, 10, VII, 146
Solanum carolinense, I, 103, 107, 108, II, 105, IV, 10, VIII, 122
 cornutum, IV, 10
 discolor, IV, 10
 eleagnifolium, Supp., 54
 nigrum, VI, 12
 robustum, IV, 10
 rostratum, I, 102, 108, IV, 10, VII, 1, VIII, 9, 10, 122
 sieglinge, IV, 10
 tuberosum, II, 56, VIII, 9
 warscewiczii, IV, 10
Solidago, V, 154
 missouriensis, I, 174
 nemoralis, I, 173
 Sonora Grape, IV, 64
 Sorghum, II, 23, 44, 54, VII, 146, 159, VIII, 27
 Soulard Apple, V, 87

- Sour Cherry, VII, 159
 Southern Fox Grape, III, 77, IV, 60, 62, 63, VI, 48, 72, 75
 Spanish Chestnut, IV, 56
speciosum, *Ribes*, IX, 2
 Spice-bush, IV, 123
 Spinach, II, 113
Spiraea, V, 154
 ulmaria, III, 51
 Spruce, III, 112
 Norway, I, 150, VII, 172, VIII, 119, IX, 30
 Squash, II, 64, 66, 70
 Staghorn Sumach, VI, 119
 Standard Pear, I, 128
 St. Augustine Grape, V, 63
Stellaria media, I, 179
 Stickseed, V, 52
 Stink-weed, VI, 121
Stipa setigera, VII, 192
 St. John's Wort, II, 10
stramonium, *Datura*, V, 56
 Strawberry, I, 142, 143, 157, II, 34, III, 11, 42, 43, 82, 83, 105, IV, 34, V, 114, VII, 46, 47, 48, 159, IX, 27
strictum, *Echinosperrum*, V, 52
 String bean, III, 105
strobis, *Pinus*, I, 24
Strombocarpa monoica, I, 65
strumarium, *Xanthium*, I, 92
 Sugar Grape, IV, 61
 Sumach, I, 100, II, 58, III, 130, VI, 118, 127, VII, 160
 Dwarf, VI, 119
 Fragrant, VI, 121, IX, 6
 Smooth, VI, 119
 Staghorn, VI, 119
 Venetian, VI, 119
 Summer Grape, III, 89, 90, IV, 60, 63, VI, 47, 72
 Rose Apple, V, 86
 Sunflower, III, 68, 131, IV, 142, V, 52, Supp., 53
 Swamp Rose-mallow, VI, 92
 Sweet Cherry, VII, 159
 Gum, IV, 123, 124, 140
 June Apple, V, 75
 -potato, I, 100, II, 56, 57, 58, 60, 61, 62, 63, VII, 146
 Bermuda, II, 61
 Brazilian, II, 61
 Nansemond, II, 61
 Sycamore, I, 150, III, 114, 127, IV, 126, 129, VI, 128, IX, 127
sylvestris, *Malva*, VI, 89
 Pinus, I, 24, V, 100
Symphoricarpus racemosus, II, 113
 vulgaris, I, 153
- T.**
- Tallman's Sweet Apple, III, 35
 Taylor Grape, I, 30, III, 87, IV, 63, 64, V, 63, 65, 66, VI, 36, 47, 48, 68, 80, 95, VII, 102, 115
 Bullet Grape, VI, 73
 Teasel, IV, 112
 Telegraph Grape, V, 65, VI, 47, VII, 106
 Thistle, I, 180, II, 10, 112, III, 67
 Canada, II, 10
 Thorn, III, 120, IV, 126
 -apple, IV, 10
 Thoroughwort, VI, 12
Thuja occidentalis, I, 24
thunbergii, *Vitis*, VI, 71
Tilia, IV, 126
 Timothy, III, 111, VII, 38, 146, VIII, 39, 49, 50, 143, IX, 51
 Tinto Grape, VI, 36
 Toad-flax, II, 10
 Tobacco, I, 80, 96, 105, IV, 10, VII, 146, 159
 Tokay Grape, V, 66
 Tomato, I, 80, 92, 95, 107, 108, III, 105, IV, 10, VI, 12, VII, 146
tomentosa, *Aristolochia*, II, 116
 Crataegus, III, 36
 Tonzuru, IV, 136
Torrubia cinerea, VI, 123
 militaris, VI, 123
toxicodendron, *Rhus*, V, 127, VI, 121
 Transcendent Crab, I, 15
 Trees, growth of trunk of, VI, 98
trichostemmoides, *Salvia*, VIII, 119
trifida, *Ambrosia*, Supp., 56
Triosteum perfoliatum, III, 134
 Trumpet Creeper, III, 64, VI, 87
tuberosa, *Asclepias*, III, 144
tuberosum, *Solanum*, II, 56, VIII, 9
tuberosus, *Helianthus*, VIII, 143
tuckeri, *Oidium*, V, 57, 70, VI, 30, 63, 79, 85, IX, 43
tulipifera, *Liriodendron*, Supp., 55
 Tulip tree, IV, 123, Supp., 55
 Turnip, I, 101, II, 105, 114, III, 11, 109, 111, 141, IV, 36, V, 69, 111, 114, VII, 159, VIII, 143
typhina, *Rhus*, VI, 119
- U.**
- ulmaria*, *Spiraea*, III, 51
Ulmus, VI, 137
Urticaceae, IV, 100
- V.**
- vaginæflora*, *Vilfa*, VIII, 122
 Venango Grape, VI, 84
 Venetian Sumach, VI, 119
Verbascum, V, 35, VII, 14
 Verbena, III, 68
verna, *Potentilla*, I, 143
Vernonia, V, 136
 fasciculata, I, 153
 novæboracensis, VIII, 119
Veronica peregrina, II, 35, VII, 48
Vilfa, VIII, 123
Vilfa vaginæflora, VIII, 122
vinifera, *Vitis*, IV, 55, 63, V, 65, VI, 32, 47, 48, 72, 74, 78, 80, 85
 Virginia Creeper, II, 78, 86, III, 77, VI, 88
virginiana, *Juniperus*, I, 24
 Virginia Seedling Grape, IV, 61, VI, 74
Vitis aestivalis, III, 89, 96, IV, 60, 61, 63, VI, 36, 47, 48, 72, 74, 75, 80, VII, 103
 arizonica, IV, 60, VI, 73, 76
 californica, IV, 60, VI, 73
 candicans, IV, 60, VI, 73, 76
 cordifolia, III, 87, 88, 89, 90, IV, 60, V, 66, 118, VI, 36, 73, 74, 75, 76, VII, 96
 labrusca, III, 87, 89, 90, IV, 60, 63, V, 60, 65, 66, 118, VI, 36, 47, 48, 71, 72, 74, 75, 76, VII, 103
 lincecumii, VI, 74
 monticola, VI, 57, 74
 mustangensis, IV, 62, VI, 76

Vitis riparia, IV, 60, 61, 63, V, 62, 65, 116, 118, VI, 36,
47, 48, 58, 72, 73, 74, 75, 95, VII, 96
rupestris, IV, 60, VI, 73, 74
thunbergii, VI, 71
vinifera, IV, 55, 63, V, 65, VI, 32, 47, 48, 72, 74,
78, 80, 85
vulpina, III, 77, 90, IV, 60, 63, V, 66, 118, VI, 48,
49, 72, 75, VII, 106
vulgaris, *Symphoricarpos*, I, 153
vulpina, *Vitis*, III, 77, 90, IV, 60, 63, V, 66, 118, VI,
48, 49, 72, 75, VII, 106

W.

Walter Grape, V, 59, 62, VI, 80
Walnut, I, 153, III, 125, 126, 127, IV, 42, 124, 126
Black, I, 153, III, 131, IV, 82, VII, 160,
Supp., 54
Warren Grape, VII, 102
warscewiczii, *Solanum*, IV, 10
Watermelon, III, 141
Water-weed, II, 11, IX, 43
Weehawken Grape, V, 59
Weeping Willow, II, 109
Wheat, I, 79, 87, 88, 159, 160, II, 16, 17, 23, 29, 30, 42,
44, 54, 55, III, 110, 111, 112, 115, 116, VII, 25, 27, 34,
38, 146, 168, 173, VIII, 25, 27, 49, 59, 143, IX, 51, 52,
Supp., 56
whipplei, *Yucca*, V, 157
White-berry, II, 113
Doyenne Pear, I, 15
Elm, I, 123
Grub fungus, I, 158, VI, 123, 125
Hellebore, IX, 13
Mulberry, I, 72, 73
Oak, III, 120, 127, VI, 64, 115
Pine, III, 92, V, 97, 100, 102, IX, 13, 29, 30, 32
Riesling Grape, VII, 102
Scotch Cluster Grape, VI, 77
Thorn, IV, 132, V, 86
Willow, IV, 72, V, 120
-wood, II, 91
Whortleberry, III, 163, Supp., 83

12 MO

Wild Black Currant, IX, 2
Cherry, IV, 123
Crab, IV, 42, Supp., 80
Endive, I, 79, 83
Plum, I, 15, 55
Rose, V, 126
Wilder Grape, IV, 63, V, 62, 63, VI, 48, 81, VII, 101,
102, 111, 116
Windsor Bean, I, 98, III, 51
Winesap Apple, IV, 45
Winter Grape, IV, 60, VI, 73
Willow, I, 24, III, 120, 153, 153, 163, 168, IV, 104, 112,
124, 126, 142, V, 127, 136, VI, 162, VII, 160
Golden, III, 168
Heart-shaped, IX, 20
Scrub, II, 90, III, 168, V, 132
Twig Apple, V, 87
Weeping, II, 109
White, IV, 72, V, 120
Witch Hazel, III, 120
Wormwood, V, 35

X.

Xanthium, VII, 159
strumarium, I, 92, III, 60

Y.

Yellow Bellflower Apple, IV, 45
Yellow Pine, V, 100
York Madeira Grape, IV, 64, V, 64
Yucca, V, 153, 159, 160, VI, 132, 133, 135, VIII, 169,
IX, 129
Yucca aloifolia, V, 153, VIII, 171, 178, IX, 129
angustifolia, V, 157, 159, VIII, 169
filamentosa, V, 158, 159, VIII, 171
flaccida, V, 159
glauca, V, 153
gloriosa, V, 159, 171
puberula, V, 153
rupicola, V, 157
whipplei, V, 157

ERRATA.

Page III, line 9, for Classified read CLASSIFIED.

Page 60, line 17, for LEUCANEÆ read LEUCANÆ.

Pages 93, 94. In making up these pages several of the names got misplaced. "Orgyia" and "Thyridopteryx," on p. 94 should follow "Ecpantheria" on p. 93. "Hæmatopis," on p. 94, should follow "Eufitchia," on p. 93. "Pronuba" and "Galleria," p. 94, should follow "Carpocapsa," on the same page; "Æstrus" should follow "Pipiza" on the same page.

Page 94. After line 10 add "*Gelechia gallæsolidaginis*, *larva* and *pupa*: I, 173-174."



ERRATA.

Page III, line 9, for Classified read CLASSIFIED.

Page 60, line 17, for LEUCANEÆ read LEUCANLÆ.

Pages 93, 94. In making up these pages several of the names got misplaced. "Orgyia" and "Thyridopteryx," on p. 94 should follow "Ecpantheria" on p. 93. "Hæmatopis," on p. 94, should follow "Eufitchia," on p. 93. "Pronuba" and "Galleria," p. 94, should follow "Carpocapsa," on the same page; "Æstrus" should follow "Pipiza" on the same page.

Page 94. After line 10 add "*Gelechia gallæsolidaginis*, *larva* and *pupa*: I, 173-174."



Bulletin of the United States Entomological Commission Department of the
Interior. U. S. Geolog. & Geograph. Survey of the Territories. F. V.
Hayden, in charge
Bd.: 6

1881

Zool. 84 I-1/7

urn:nbn:de:bvb:12-bsb11906059-6